



OBLIX MT[®]


HERBICIDE

PCS 06286

FOR USE AS AN AGRICULTURAL HERBICIDE
FOR PROFESSIONAL USE ONLY

A suspension concentrate containing 300 g/L (30.0% w/w) metamitron and 150 g/L (13.2% w/w) ethofumesate.
Also contains 1,2 benzisothiazolin-3-one. May produce an allergic reaction.

A contact and residual pre and post-emergence herbicide for use on sugar beet, fodder beet and mangels.



SAFETY INFORMATION

Warning

Harmful if swallowed.

Very toxic to aquatic life with long lasting effects.

When using, do not eat, drink or smoke.

Upon skin wash with plenty of soap and water.

IF SWALLOWED: Call a POISON CENTRE or doctor/physician if you feel unwell.

Collect spillage.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

Contains 1,2 benzisothiazolin-3-one. May produce an allergic reaction.

To avoid risks to human health and the environment, comply with the instructions for use.

PCS 06286

A1.0 OBLIX MT IE PCS 06286

5 Litres
350 g/L METAMITRON
150 g/L ETHOFUMESATE



INSTRUCTIONS FOR USE

Crops	Maximum individual dose (L product/ha)	Maximum total dose (L product/ha/crop)	Latest timing of application
Sugar beet, fodder beet and mangels	2.0	6.0	Before crop leaves meet between the rows

OTHER SPECIFIC RESTRICTIONS

1. The maximum total dose must not exceed 1.0 kg ethofumesate per hectare in any three year period.
2. Fodder beet and mangels must not be grazed by livestock or harvested for animal consumption until at least 103 days following the last application.
3. Beet crops may be sown at any time following the use of Oblix MT. Three months must elapse before any other crop is sown.

READ ALL SAFETY PRECAUTIONS AND DIRECTIONS FOR USE BEFORE USE.

SAFETY PRECAUTIONS

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces

DO NOT BREATHE SPRAY

WASH HANDS AND EXPOSED SKIN before eating, drinking or smoking, and after work.

KEEP OUT OF THE REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

WASH OUT CONTAINER THOROUGHLY empty washings into spray tank and dispose of safely.

DO NOT RE-USE CONTAINER for any purpose.

ENVIRONMENTAL PROTECTION

DO NOT CONTAMINATE WATER WITH THE PRODUCT OR ITS CONTAINER (DO NOT CLEAN APPLICATION EQUIPMENT NEAR SURFACE WATER/AVOID CONTAMINATION VIA DRAINS FROM FARMYARDS AND ROADS).

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Oblix MT is a contact and residual herbicide for the control of annual broadleaved weeds in sugar beet, fodder beet and mangels.

RESTRICTIONS

This product may be safely used on all commonly grown sugar beet varieties under normal conditions. Crop tolerance may be reduced by stress caused by growing conditions, effects of pests, disease or other pesticides, nutrient deficiency etc.

RESISTANCE MANAGEMENT

Strains of some annual grasses, e.g. black-grass, wild-oats, and Italian rye-grass, have developed resistance to herbicides which may lead to poor control. A strategy for preventing and managing such resistance should be adopted. Guidelines have been produced by the Weed Resistance Action Group and copies are available from the HGCA, CPA, your distributor, crop adviser or product manufacturer.

WEED CONTROL

Best results are obtained from a series of treatments applied as an overall spray commencing when earliest germinating weeds are no larger than fully expanded cotyledon and the majority of the crop is at the fully expanded cotyledon stage.

Oblix MT may be used on all soil types but residual activity may be reduced on those with more than 5% organic matter.

The following weeds are **susceptible** at the recommended dose of 2.0 L product/ha:

Common chickweed	Field penny-cress
Common hempnettle	Red deadnettle
Fallopia	Redshank
Field pansy	Shepherd's purse

The following weeds are **moderately susceptible** at the recommended dose of 2.0 L product/ha:

Black bindweed	Corn chamomile
Charlock	Volunteer oilseed rape
Common amaranth	

CROP SPECIFIC INFORMATION

Oblix MT should be used as a split application, where up to 3 applications can be made per crop.

Pre-Emergence Application

The first application can be made pre-emergence at a maximum application rate of 2.0 L product/ha. The application should be applied in 200 L of water/ha. Only 1 application can be made pre-emergence.

Post-Emergence Applications

If a pre-emergence application has already been made, up to 2 post-emergence applications can be made. Alternatively, if no pre-emergence application has been made up to 3 post-emergence applications can be made.

The first post-emergence application can be made on small, emerging weeds and when the majority of the crop is at the fully expanded cotyledon stage. Best effects are achieved when the weeds are at the cotyledon stage. A follow up spray should be made on newly emerged weeds. A minimum of 5 days must elapse between treatments.

A maximum individual dose of 2.0 L product/ha applies and no more than 6.0 L product/ha may be applied in total taking into account pre and post-emergence applications

MIXING AND SPRAYING

Apply as a FINE spray (BCPC quality). Pre-emergence applications should be made in 200L of water/ha; post-emergence applications should be made in 80-200 L of water/ha.

On emptying the container, RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add the washings to the sprayer at the time of filling and dispose of the container safely.

Half fill the sprayer tank with clean water. Begin agitation and add the required quantity of Oblix MT to the sprayer tank. Add the adjuvant or oil and/or other herbicide if appropriate. Fill the spray tank whilst continuing agitation and maintain agitation until spraying is completed.

When spraying take care to avoid drift onto neighbouring crops, buildings or non-target plants.

After spraying, drain the tank completely and thoroughly clean the inside of the tank. Flush the tank, hoses and boom with clean water for at least 5 minutes.

Partially fill the tank with clean water and add a cleaning agent. Completely fill the tank with water then flush the cleaning solution through the hoses, boom and nozzles. Repeat this procedure with clean water. Remove nozzles and screen and clean separately in a bucket containing water (and cleaning agent).

FOLLOWING CROPS

Deep and thorough soil cultivation is necessary before sowing or planting succeeding crops. Plough to a depth of 15-20 cm.

Beet crops may be sown at any time after using Oblix MT. Following normal harvest any other crop may be sown after mould-board ploughing to a minimum depth of 15 cm.

COMPATIBILITY

For information on tank mixes please see the UPL website: ie.uplonline.com.

CONDITIONS OF SUPPLY

All goods supplied by the company are of good quality and we believe them to be fit for purpose. However, as we cannot exercise control over their storage, handling, mixing or use or the weather conditions before, during or after application, which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded, and no responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods. Brand names used in this label may be registered trademarks of UPL Europe Ltd, or other manufacturers in which propriety rights may exist.

To access the Safety Data Sheet for OBLIX MT please use the link below or scan the code.



ie.uplonline.com/productdetails/oblix-mt

Alternatively, contact your local supplier.



OBLIX MT[®]

HERBICIDE


PCS 06286

FOR USE AS AN AGRICULTURAL HERBICIDE
FOR PROFESSIONAL USE ONLY

A suspension concentrate containing 350g/L (30.7%w/w) metamitron and 150g/L (13.2% w/w) ethofumesate.
Also contains 1,2 benzisothiazolin-3-one. May produce an allergic reaction.

A contact and residual pre and post-emergence herbicide for use on sugar beet, fodder beet and mangels.

SPECIMEN -
2019 to date



SAFETY INFORMATION

Warning

Harmful if swallowed.

Very toxic to aquatic life with long lasting effects.

When using, do not eat, drink or smoke.

If on skin wash with plenty of soap and water.

IF SWALLOWED: Call a POISON CENTRE or doctor/physician if you feel unwell.

Collect spillage.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

Contains 1,2 benzisothiazolin-3-one. May produce an allergic reaction.

To avoid risks to human health and the environment, comply with the instructions for use.

PCS 06286

AT 1,0 OBLIX MT IE PCS 06286

5 Litres
350 g/L METAMITRON
150 g/L ETHOFUMESATE

OBLIX MT[®]

PCS 06286

FOR USE AS AN AGRICULTURAL HERBICIDE

A contact and residual pre and post-emergence herbicide for use on sugar beet, fodder beet and mangels.

A suspension concentrate containing 350g/L (30.7%w/w) metamitron and 150g/L (13.2% w/w) ethofumesate. Also contains 1,2 benzisothiazolin-3-one. May produce an allergic reaction.

INSTRUCTIONS FOR USE

Crops	Maximum individual dose (L product/ha)	Maximum total dose (L product/ha/crop)	Latest timing of application
Sugar beet, fodder beet and mangels	2.0	6.0	Before crop leaves meet between the rows

OTHER SPECIFIC RESTRICTIONS:

1. The maximum total dose must not exceed 1.0kg ethofumesate per hectare in any three year period.
2. Fodder beet and mangels must not be grazed by livestock or harvested for animal consumption until at least 103 days following the last application.
3. Beet crops may be sown at any time following the use of Oblix Mt. Three months must elapse before any other crop is sown.

READ ALL SAFETY PRECAUTIONS AND DIRECTIONS FOR USE BEFORE USE.

SAFETY PRECAUTIONS

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

DO NOT BREATHE SPRAY

WASH HANDS AND EXPOSED SKIN before eating, drinking or smoking, and after work.

KEEP OUT OF THE REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

WASH OUT CONTAINER THOROUGHLY empty washings into spray tank and dispose of safely.

DO NOT RE-USE CONTAINER for any purpose.

ENVIRONMENTAL PROTECTION

DO NOT CONTAMINATE WATER WITH THE PRODUCT OR ITS CONTAINER (DO NOT CLEAN APPLICATION EQUIPMENT NEAR SURFACE WATER/AVOID CONTAMINATION VIA DRAINS FROM FARMYARDS AND ROADS).

SHAKE WELL BEFORE USE

PROTECT FROM FROST

For Batch Number and
Manufacturing Date see container

UPL Europe Ltd

The Centre, 1st Floor, Birchwood Park, Warrington, WA3 6YN, UK.

Telephone: +44 (0)1925 819999 Fax: +44 (0)1925 817425

For 24 hour emergency information contact:

CARECHEM24 : +44 (0) 1235 239670