



# Mirror<sup>®</sup>

syngenta.

**GROUP M04 FUNGICIDE**

PCS number 06425  
UF: 5PF0-VOQD-700F-5MGN



A suspension concentrate formulation containing 500 g/l (39.5% w/w) folpet. Contains folpet and hexamethyl tetramine: may produce an allergic reaction.

A fungicide for protection against leaf spot (*Zymoseptoria tritici*) on wheat and triticale, leaf blotch (*Rhynchosporium secalis*) on barley and triticale and net blotch (*Pyrenophora teres*) and ramularia leaf spot (*Ramularia collo-cygni*) on barley.

In case of toxic or transport emergency ring +44 (0) 1484 538444 any time

SHAKE WELL BEFORE USE. PROTECT FROM FROST.

## SAFETY INFORMATION

### FOR PROFESSIONAL USE ONLY

MIRROR is a suspension concentrate formulation containing 500 g/L (39.5% w/w) folpet

#### Warning

**May cause an allergic skin reaction.**

**Suspected of causing cancer.**

**Very toxic to aquatic life with long lasting effects.**

Obtain special instructions before use.

Avoid breathing mist or vapours.

Wear protective gloves/ protective clothing/ eye protection/ face protection/ hearing protection.

IF exposed or concerned: Get medical advice/ attention.

If skin irritation or rash occurs: Get medical advice/attention.

Collect spillage.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for empty triple rinsed clean containers which can be disposed of as non-hazardous waste.

**To avoid risks to human health and the environment, comply with the instructions for use.**

PCS: 06425

UF: 5PF0-VOQD-700F-5MGN



#### Authorisation holder

Adama Agricultural Solutions UK Ltd  
Third Floor East, 1410 Arlington Business  
Park, Theale, Reading, RG7 4SA  
Tel: + 44 (0)1635 860555

#### Marketing Company

Syngenta Ireland Ltd.,  
Block 6, Cleaboy Business Park,  
Old Kilmeaden Road, Waterford, Ireland  
Tel: (051) 377203

L1130394 IREL/12C PPE 4222873

**5 litres**

Product names marked © or ™, the ALLIANCE FRAME the SYNGENTA Logo and the PURPOSE ICON are Trademarks of a Syngenta Group Company

**IMPORTANT INFORMATION**

FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL FUNGICIDE

<b>Crop</b>	<b>Maximum individual dose (litres of product/ha)</b>	<b>Maximum total dose (litres of product/ha/crop)</b>	<b>Latest time of application</b>
Winter wheat, spring wheat, winter barley, spring barley and triticale	1.5	3	Before end of heading/ inflorescence fully emerged (GS 59)

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**

**SAFETY PRECAUTIONS****(a) Operator Protection**

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate or handling contaminated surfaces. WASH CONCENTRATE from skin or eyes immediately. WASH HANDS AND EXPOSED SKIN before eating or drinking and after work. IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible).

**(b) Environmental Protection**

To protect aquatic organisms respect an unsprayed buffer of 5 metres to surface water bodies.

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

**(c) Storage and Disposal**

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.  
DO NOT RE-USE CONTAINER FOR ANY PURPOSE.

## DIRECTIONS FOR USE

**IMPORTANT:** This information is approved as part of the Product Label. All instructions within this section must be carefully read in order to obtain safe and successful use of this product.

### Resistance

MIRROR® contains folpet, a protectant fungicide from the phthalimide chemical group with multi-site contact activity. For resistance management purposes, it's mode of action is classified as M4 under the FRAC\* code which indicates low-risk without any signs of resistance or cross-resistance developing.

For current advice on resistance management contact your agronomist or specialist advisor, and visit the FRAG-UK website.

### Disease control in wheat, barley and triticale

For the reduction of leaf spot (*Zymoseptoria tritici*) in winter and spring wheat and triticale and leaf blotch (*Rhynchosporium secalis*) in winter and spring barley and triticale; and moderate control of net blotch (*Pyrenophora teres*) and ramularia leaf spot (*Ramularia collo-cygni*) in winter and spring barley. MIRROR may be applied in a programme of 2 suitably timed sprays.

### Dose rate

1.5 L/ha per application

### **Septoria leaf spot (*Zymoseptoria tritici*) in winter and spring wheat and triticale**

As MIRROR is a protectant fungicide with contact activity, it is important to apply the first spray before the disease becomes established in the crop (normally GS 30-33). A second spray should be timed to protect new growth from disease and therefore needs to be applied prior to further infection. This is normally 3 weeks after the first application but may be sooner under conditions of rapid growth and high disease pressure. Where disease infection or re infection has occurred prior to application, MIRROR should be mixed with alternative chemistry to provide curative activity (see Compatibility section).

### **Leaf blotch (*Rhynchosporium secalis*) in winter and spring barley and triticale; and net blotch (*Pyrenophora teres*) and ramularia leaf spot (*Ramularia collo-cygni*) in winter and spring barley**

As MIRROR is a protectant fungicide with contact activity, it is important to apply the first spray before the disease becomes established in the crop (normally GS 30-33). A second spray should be timed to protect new growth from disease infection. This is normally 3 weeks after the first application but may be sooner under conditions of rapid growth and high disease pressure. Where disease infection or re infection has

occurred prior to application, MIRROR should be mixed with alternative chemistry to provide curative activity (see Compatibility section).

### **Water volume**

Apply MIRROR in a water volume of 200-400 L/ha.

Qualified recommendation: lower water volumes of 150 L/ha may also be used\*;  
however, these have not been supported by effectiveness or crop safety data.

\*or the equivalent of a 1% v/v dilution.

As MIRROR has contact activity, it is important to achieve a good coverage of the crop when spraying.

### **Mixing and spraying**

For use by tractor mounted/trailed sprayer only.

Before spraying it is important to check all hoses, filters and nozzles, and to ensure that the sprayer is clean and correctly calibrated to give an accurate application at the correct volume. Half fill sprayer tank with clean water. Add the required amount of MIRROR to the water and commence agitation. On emptying the container, RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of container safely. Add the remainder of the water whilst continuing to agitate the mixture until spraying is complete.

Apply MIRROR as a medium spray (as defined by the BCPC) in cereals.

Thoroughly clean the sprayer and measuring equipment after using MIRROR.

Crops should not be re-entered until spray residues are dry.

### **Compatibility**

Consult SYNGENTA IRL Ltd or your distributor for details on the products which may be mixed with MIRROR in Ireland.

Maintain agitation during the mixing process and continue until the spraying is complete. Do not leave in spray tank for long periods without using.

### **Caution**

After spraying a herbicide, sprayer tanks and equipment should be thoroughly cleaned before using MIRROR.

## **DISCLAIMER/CONDITIONS OF SUPPLY**

The specified properties of our products and the mode of application stated on this label have been established on the basis of research and experience. Products conform to specification at the time of delivery but, as we exercise no control over their subsequent storage, handling, mixing or use or the weather conditions before, during and after application, all of which may affect the performance of the products, no responsibility or liability will be accepted by us or our re-sellers for any failure in performance, damage or injury to person or property whatsoever arising from the storage, handling, application or use of the products. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in or make recommendations concerning the use of such products. We recommend you contact your dealer to request advice on the suitability of this product for any new and/or unusual growing methods or for new varieties not listed on this label.

To access the Safety Data Sheet for this product, scan QR code:



Alternatively, contact your supplier

**IMPORTANT INFORMATION**

FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL FUNGICIDE

Crop	Maximum individual dose (litres of product/ha)	Maximum total dose (litres of product/ha/crop)	Latest time of application
Winter wheat, spring wheat, winter barley, spring barley and triticale	1.5	3	Before end of heading/ inflorescence fully emerged (GS 59)

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**

**SAFETY PRECAUTIONS****(a) Operator Protection**

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate or handling contaminated surfaces. WASH CONCENTRATE from skin or eyes immediately. WASH HANDS AND EXPOSED SKIN before eating or drinking and after work. IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible).

**(b) Environmental Protection**

To protect aquatic organisms respect an unsprayed buffer of 5 metres to surface water bodies.

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

**(c) Storage and Disposal**

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place. DO NOT RE-USE CONTAINER FOR ANY PURPOSE.