

BROKPAC

An emulsifiable concentrate containing 250 g/litre (25.50% w/w) trinexapac-ethyl.

**WARNING**

May cause an allergic skin reaction.

May cause damage to organs (Gastrointestinal tract) through prolonged or repeated exposure.

Very toxic to aquatic life with long lasting effects.

Keep out of reach of children.

Avoid breathing dust/fume/gas/mist/vapours/spray.

Wear protective gloves, protective clothing.

Get medical advice/attention if you feel unwell.

IF ON SKIN: Wash with plenty of soap and water.

If skin irritation or rash occurs: Get medical advice/attention.

Take off contaminated clothing and wash it before reuse.

Collect spillage.

Repeated exposure may cause skin dryness and cracking.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

PCS No. 06494

BROKPAC

An emulsifiable concentrate containing 250 g/litre (25.50% w/w) trinexapac-ethyl.

FOR PROFESSIONAL USE ONLY

A growth regulator for winter and spring wheat, winter and spring barley, winter and spring oats, durum wheat, rye, triticale, and grassland (seed crops).

PROTECT FROM FROST SHAKE WELL BEFORE USE**Authorisation holder & Marketing Company:****Rainbow Agrosociences Europe, S.A.U.**

Polígono Industrial Fondo de litera, Autovía A-2 Km. 441,6, 22520 Fraga (Huesca), Spain Telephone number: (+34) 973 737377

E-mail: registration_eu@rainbowagro.com

PROTECT FROM FROST STORE IN A COOL DRY PLACE

In the event of emergency, call the National Poisons

Information Centre, Beaumont

Hospital at 01 809 2166 or 01 837 9964'

UFI: 7390-J0F4-J00J-X2AQ

Batch number:

Date of Manufacture:

Rainbow
all about growing

**SAFETY PRECAUTIONS****Operator Protection**

AVOID CONTACT WITH SKIN AND EYES.

WEAR EYE/FACE PROTECTION when handling the concentrate.

FOR USE BY TRACTOR MOUNTED/TRAILLED SPRAYER ONLY.

Environmental Protection

Do not contaminate water with the product or its container.

Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

Storage and Disposal

RINSE CONTAINER THOROUGHLY, by using an integrated pressure rinsing device or manually rinsing three times. Add washings to the sprayer at the time of filling and dispose of safely.

Do not re-use container for any other purpose and dispose of safely.

Restrictions

Apply BROKPAC only to healthy, actively growing crops.

Do not apply during periods of frosty weather or when frost is imminent.

Do not apply BROKPAC to crops that are stressed by severe weather conditions, drought, frost, disease, insect damage, nutritional deficiency, etc.

Do not apply if rain is expected or if the crop is wet.

Avoid spray drift on to neighbouring crops.

Aprire qui



IMPORTANT INFORMATION

FOR USE ONLY AS A PROFESSIONAL PLANT GROWTH REGULATOR				
Crops	Max individual dose l/ha	Max no. of applications	Max total dose l/ha per crop/year	Latest time of application
Winter wheat	0.4	-	0.4	Before flag leaf sheath extending stage (GS 41)
Winter barley	0.6	-	0.6	
Winter and spring oats	0.4	-	0.4	Before second node detectable stage (GS 32)
Grassland (seed crop)	0.8	-	0.8	
Spring wheat	0.4	-	0.4	Before third node detectable stage (GS 33)
Spring barley	0.5	-	0.5	
Durum wheat, rye, triticale	0.4	-	0.4	

DIRECTION FOR USE

PROPERTIES OF BROKPACK

BROKPACK is a growth regulator for crop height reduction, lodging prevention and yield protection in all varieties of winter and spring wheat, winter and spring barley and winter and spring oats, durum wheat, rye, triticale and grassland (seed crops).

Treatment may lead to ears remaining erect through to harvest.

MIXING AND SPRAYING

Make sure the sprayer is set to give an even application at the correct volume.

Fill the spray tank with half the required volume of clean water and start agitation. Add the required amount of BROKPACK, agitate, and continue agitation whilst adding the rest of the water.

Agitate the mixture thoroughly before use and continue agitation during spraying.

Thoroughly wash all spray and measuring equipment with water and a wetting agent immediately after use.

APPLICATION

Spray volume

Apply BROKPACK in a minimum of 200 l/ha of water. Increased penetration will be obtained with an increase in water volume but the necessity for this will be dependent on crop growth stage and habit.

Spray nozzles

A medium spray quality is preferred for application of BROKPACK. A spray pressure of 2-3 bar is recommended.

Spraying

Take particular care to avoid overlapping of spray swaths. Apply only using a ground sprayer.

RECOMMENDATIONS

Winter Wheat

Timing and dose

Apply at 0.4 l/ha from the leaf sheath erect stage (GS 30) but before the flag leaf extending stage (GS 41).

Winter Barley

Timing and dose

Apply at 0.4 l/ha from the leaf sheath erect stage (GS 30) but before the third node detectable stage (GS 33).

or

Apply at 0.6 l/ha from the flag leaf just visible stage (GS 37) but before the flag leaf extending stage (GS 41).

Spring Barley

Timing and dose

Apply at 0.5 l/ha from the leaf sheath erect stage (GS 30) but before the third node detectable stage (GS 33).

Spring Wheat

Timing and dose

Apply at 0.4 l/ha from the leaf sheath erect stage (GS 30) but before the third node detectable stage (GS 33).

Winter and Spring Oats

Timing and dose

Apply at 0.4 l/ha from the leaf sheath erect stage (GS 30) but before the second node detectable stage (GS 32).

Rye, Triticale and Durum Wheat

Timing and dose

Apply at 0.4 l/ha from the leaf sheath erect stage (GS 30) but before the third node detectable stage (GS 33).

Grassland (seed crops only)

Timing and dose

Apply at 0.8 l/ha from the leaf sheath erect stage (GS 30) but before the second node detectable stage (GS 32).

CROP FAILURE

In the event of crop failure for any reason, cereals and oilseed rape can be planted in soil treated with BROKPA. Due to reduced activity via the root system and to its rapid degradation in soil, no problems with following crops are foreseen for this product.

Specimen-
2026 to date