

# Difure<sup>®</sup> Solo

A systemic fungicide for use on all varieties of sugar beet and fodder beet.



Emulsifiable concentrate containing 250 g/L of difenoconazole (23.6 % w/w). Contains hydrocarbons, C10-C13, aromatics, <1% naphthalene.

#### RISK AND SAFETY INFORMATION

Harmful if swallowed. May be fatal if swallowed and enters airways. Causes serious eye irritation. Suspected of causing cancer. May cause damage to organs through prolonged or repeated exposure. Very toxic to aquatic life with long lasting effects.

Keep out of reach of children. Avoid release to the environment. Wear protective gloves/protective clothing/eye protection/face protection. IF SWALLOWED: Immediately call a POISON CENTER or doctor/physician. IF IN EYES: Rinse continuously with water for several minutes. Remove contact lenses if present and easy to do – continue rinsing. Do NOT induce vomiting. If eye irritation persists: Get medical advice/attention.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty triple rinsed containers which can be disposed of as non-hazardous waste. Repeated exposure may cause skin dryness or cracking. To avoid risks to human health and the environment, comply with the instructions for use.



**DANGER**

FOR PROFESSIONAL USE ONLY | UFI: WC00-509Q-400N-GX5A | PCS no. 06537

Difure<sup>®</sup> is a registered trademark for Globachem N.V.

PROTECT FROM FROST  
SHAKE WELL BEFORE USE  
BATCH NUMBER: SEE BOTTLE

GROUP 3 FUNGICIDE

CONTENT: 1 lit. e

Emergency number: In the event of emergency call the National Poisons Information Centre, Beaumont Hospital at 01 809 2166 or 01 837 9964.

#### Registration holder:

GLOBACHEM NV  
Brustum Industriepark  
Lichtenberglaan 2019  
B-3800 Sint-Truiden • Belgium  
Tel: +32 11 78 57 17  
globachem@globachem.com  
www.globachem.com

  
**Globachem**  
Your Crop Counts

#### SAFETY PRECAUTIONS

##### OPERATOR PROTECTION

Engineering control of operator exposure must be used where reasonably practical in addition to the following personal protective equipment: WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), GLOVES AND FACE PROTECTION (FACESHIELD), when handling the concentrate. WASH HANDS AND EXPOSED SKIN before meals and after work. WHEN USING, DO NOT DRINK, EAT OR SMOKE.

IF YOU FEEL UNWELL, seek medical advice immediately and show this container or label.

IN CASE OF CONTACT WITH EYES, RINSE IMMEDIATELY with plenty of water and seek medical attention.

##### ENVIRONMENTAL PROTECTION

To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies.

DO NOT CONTAMINATE WATER with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

##### STORAGE AND DISPOSAL

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely.

KEEP OUT OF REACH OF CHILDREN.

Do not re-use container for any purpose.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFF.

THIS MATERIAL AND ITS CONTAINER must be disposed of in a safe way. Store in a cool, dry place.

##### DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the product label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

#### GENERAL INFORMATION

DIFURE SOLO is a systemic fungicide containing difenoconazole which is an active ingredient belonging to the chemical family of triazoles (FRAC code 3) for use on fodder and sugar beet. DIFURE SOLO is aimed at controlling rust (*Uromyces betae*). DIFURE SOLO quickly penetrates into the beet leaves and protects them for several weeks.

#### CROP SPECIFIC INFORMATION

Apply DIFURE SOLO at 0.5 L/ha at the first signs of the disease, before the disease becomes established. DIFURE SOLO gives prolonged protection from re-infection, but a second application can be applied where crops are at risk from later attacks. Respect a minimum application interval of 14 days between applications. Do not apply before BBCH 39 (Crop cover complete: leaves cover 90% of ground).

#### MIXING AND SPRAYING

Make sure the sprayer is clean and set to give an even application at the correct volume.

Fill the spray tank with half the required volume of clean water and start agitation. Add the required amount of DIFURE SOLO and continue agitation while adding the rest of the water. Agitate the mixture thoroughly before use and continue agitation during spraying.

Do not leave the spray liquid in the sprayer for long periods (such as during meal breaks or overnight).

Thoroughly wash all spray equipment with water immediately after use.

#### WATER VOLUME

Apply DIFURE SOLO in a recommended 100-1000 litres of water per hectare. A MEDIUM quality spray (BCPC) is preferred for application of DIFURE SOLO. A spray pressure of 2-3 bars is recommended.

#### COMPATIBILITY

The mixtures should be performed in accordance with the relevant regulations and the official recommendations of good agricultural practices. For up to date details of compatible tank-mixes contact your manufacturer/distributor.

#### RESISTANCE

DIFURE SOLO contains difenoconazole, a DMI fungicide (FRAC code 3). Avoid repeated use of fungicides having the same mode of action. Strains of Light Leaf Spot resistant to DMI fungicides are known to exist. To avoid development of resistance, apply this product protectively in response to disease forecasts.

For some further advice on resistance management in DMI's contact your distributor, crop adviser or product manufacturer.

#### COMPANY INFORMATION

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use, or the weather conditions before, during or after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded. No responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

Our company is not liable for any consequences resulting from the mixtures with other products.

#### IMPORTANT INFORMATION FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE

CROPS	MAXIMUM INDIVIDUAL DOSE (L PRODUCT PER HECTARE)	MAX NUMBER OF APPLICATIONS (PER CROP)	LATEST TIME OF APPLICATION
Sugar beet & Fodder beet	0.5	2	21 days before harvest

Other specific restrictions: A minimum interval of 14 days must be observed between applications. Do not apply before BBCH 39.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE.



5 404020 112685

ET13877\_ED2-V1



IE