

PRIEST

An EC formulation containing 144 g/litre (13.85% w/w) fluroxypyr-meptyl (100 g/litre acid equivalent), 80 g/litre (7.69% w/w) clopyralid and 2.5 g/litre (0.24% w/w) florasulam A post-emergence herbicide for use on MANAGED AMENITY TURF, including lawns, and AMENITY GRASSLAND for the control of DAISY, DANDELION, CLOVER, BUTTERCUP, RIBWORT PLANTAIN and other broad-leaved weeds.

5L

**FOR USE ONLY AS AN HORTICULTURAL HERBICIDE
FOR PROFESSIONAL USE ONLY**

SAFETY INFORMATION



DANGER

May be fatal if swallowed and enters airways
Causes skin irritation
May cause an allergic skin reaction
Causes serious eye irritation
Harmful if inhaled
Very toxic to aquatic life with long lasting effects

Wear protective gloves / protective clothing / eye protection

IF SWALLOWED: Immediately call a POISON CENTER or doctor/physician

IF ON SKIN: Wash with soap and water

IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do - continue rinsing

Do NOT induce vomiting

Collect spillage

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

Do not contaminate water with the product or its container. (Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads).

To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies

To avoid risks to human health and the environment, comply with the instructions for use.

PCS No: 06628

Safety precautions

If exposed or if you feel unwell seek medical advice immediately (show the label where possible).

No specific antidote is available. Treat symptomatically.

In case of accident, call the 24-hour National Poisons Information Centre:

Tel.: +353 (0)1 809 2566 or +353 (0) 1 837 9964.

Approval holder:

Sharda Cropchem España S.L.
Edificio Atalayas Business Center
Carril Condomina nº 3, Planta 12,
30006 Murcia, Spain
Tel: +34 868 12 75 89

Marketing company:

Sharda Cropchem España S.L.
Edificio Atalayas Business Center
Carril Condomina nº 3, Planta 12,
30006 Murcia, Spain
Tel: +34 868 12 75 89



Sharda Cropchem

BATCH NUMBER: SEE CONTAINER

UFI: 9F5J-WJQS-NG0X-336J



Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

Wear suitable protective clothing (coveralls), suitable protective gloves and face protection (faceshield) when handling the concentrate.

Wear suitable protective gloves when handling contaminated surfaces.

Wear suitable protective clothing (coveralls) and suitable protective gloves when applying by hand-held equipment.

DO NOT BREATHE SPRAY.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH CONCENTRATE from skin and eyes immediately.

WASH HANDS AND EXPOSED SKIN before eating and drinking, and after work.

Environmental protection

DO NOT CONTAMINATE WATER with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from yards and roads.

To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies

Storage and disposal

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely.

DO NOT RE-USE CONTAINER for any purpose.

SHAKE WELL BEFORE USE. PROTECT FROM FROST.

KEEP OUT OF REACH OF CHILDREN.

General information and general recommendations

PRIEST herbicide has activity against a range of broad-leaved weeds. The ideal timing for application is when the weeds are small and actively growing.

Instructions for use

FOR PROFESSIONAL USE ONLY AS A HORTICULTURAL HERBICIDE

Crop	Max. single dose	Max. no. of applications	Max. total dose	Latest time of application
Grassland - Amenity	2 L/ha	1 per year	2 L/ha	N/A

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

Method of application: Tractor mounted/trailed horizontal boom sprayer. Handheld/knapsack sprayer.

Other specific restrictions: Application to amenity grass should not take place between the 30th of September and the 1st of February.

Important notes:

- Broad-leaved weeds that are not present at application will not be controlled.
- Clippings from grass treated with PRIEST can be safely used for mulch after the third cut. An interval of four weeks must elapse between application of PRIEST and re-seeding turf. Do not apply if turfgrass is wet.
- Do not apply to turf, lawns or grass areas which are under stress.

- Do not apply if night temperatures are low, if ground frost is imminent, or in periods of prolonged cold or dry weather. Ensure weeds are actively growing as after periods of prolonged drought, weeds can take a long time to start actively growing again after soil moisture returns.
- Take extreme care to avoid drift onto crops and non-target plants e.g. trees, shrubs, bedding, outside the target area.

Area of use: PRIEST can be applied to established managed amenity turf, including lawns, and amenity grassland. PRIEST has been tested for selectivity on the following range of turf grass species:

- Annual meadow-grass: Chewings fescue and creeping bent
- Perennial ryegrass: Smooth-stalked meadow-grass and rough stalked bluegrass

In view of the large number of turf grass cultivars grown consult manufacturer for current approved list or test PRIEST for turf safety on a small area of turf before overall application.

Application timing: Apply when weeds are in active growth normally from March to October when the soil is moist. Do not apply in periods of drought unless irrigation is applied. Avoid mowing 3 days before and after spraying to ensure sufficient weed leaf surface is present and to allow uptake and movement of PRIEST within the weed.

Water volume: For overall application apply PRIEST in 200 litres of water per hectare. For knapsack application apply PRIEST in 2 litres of water per 100 m².

Application equipment: PRIEST may be applied through tractor-mounted hydraulic sprayers or knapsack sprayers providing they are in good working order and have been calibrated according to the manufacturers' recommendations. Do not apply through CDA applicators.

Mixing: Half fill the spray tank with water and add the required amount of PRIEST. Fill up the spray tank, agitating continuously to ensure thorough mixing, and maintain agitation until spraying is complete. Use only clean water for mixing.

Spray quality: Apply PRIEST as a MEDIUM spray as defined by the BPCP system.

Tank cleaning: To avoid subsequent injury to crops other than cereals, all spraying equipment must be thoroughly cleaned both inside and out, using All Clear Extra spray cleaner as follows:

1. Immediately after spraying, drain tank completely. Any contamination on the outside of the spraying equipment should be removed by washing with clean water.
2. Rinse inside of tank with clean water and flush through booms and hoses using at least one tenth of the spray tank volume. Drain tank completely.
3. Half fill tank with clean water and add All Clear Extra at the recommended rate. Agitate and then briefly flush the boom and hoses with the cleaning solution. Top up with water making sure the tank is completely full and allow to stand for minutes with agitation. Flush the boom and hoses and drain tank completely
4. Nozzles and filters should be removed and cleaned separately with All Clear Extra solution containing 50 ml of All Clear Extra per 10 litres of water.
5. Rinse the tank with clean water and flush through the boom and hoses using at least one tenth of the spray tank volume. Drain tank completely.

Note: If it is not possible to drain the tank completely, step 3 must be repeated before going onto step 4.

Resistance

PRIEST contains active ingredients with differing modes of action and the risk of resistance building is therefore reduced. However, as florasulam is an ALS inhibitor there is a risk of resistance building to this active ingredient and so precautions should be taken to minimise the risk. Therefore, avoid using single mode of action herbicides, such as ALS inhibitors in the same field over a number of years. Users are advised to apply products containing herbicides with different modes of action or use sequences or tank mixtures where two or more components are active against the target weeds.

Specific weeds controlled

1L/ha	1.5L/ha	2L/ha
Black medick	Common daisy	Creeping buttercup
Bristly oxtongue	Common dandelion	Ribwort plantain
Common cat's ear	Common mouse-ear	Self heal
Creeping cinquefoil	Greater plantain	Slender speedwell
Sorrel		
White clover		
Bird's-foot trefoil (MC)		

MC: Moderate control

Specimen-
2024 to date