



HYGRASS XTRA

FOR THE SELECTIVE CONTROL OF ANNUAL AND PERENNIAL BROADLEAVED WEEDS IN ESTABLISHED AGRICULTURAL GRASSLAND AND AMENITY GRASSLAND.

A SOLUBLE CONCENTRATE CONTAINING 344 G/L 2,4-D AND 120 G/L DICAMBA AS THE DIMETHYLAMINE SALT.
FOR USE ONLY AS AN AGRICULTURAL/HORTICULTURAL HERBICIDE

FOR PROFESSIONAL USE ONLY

MAX DOSE
3.5L/HA
READ LABEL FOR DETAILS

SELECTIVE HERBICIDE FOR BROADLEAVED WEEDS



DANGER

RISK AND SAFETY INFORMATION

Harmful if swallowed.

Causes serious eye damage.

May cause an allergic skin reaction.

Very toxic to aquatic life with long lasting effects.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

IF SWALLOWED: Call a POISON CENTER or doctor/physician if you feel unwell.

IF ON SKIN: Wash with plenty of soap and water.

If skin irritation or rash occurs: Get medical advice/attention.

Immediately call a POISON CENTER or doctor/physician.

Avoid release to the environment.

Wear suitable gloves and eye/face protection

To avoid risks to human health and the environment, comply with the instructions for use.

UFI: 580D-R89J-1GAS-P611 **PCS No. 06765**

IMPORTANT INFORMATION FOR USE ONLY AS AN AGRICULTURAL/HORTICULTURAL HERBICIDE.

CROP	MAXIMUM INDIVIDUAL DOSE	MAXIMUM TOTAL DOSE	LATEST TIME OF APPLICATION
Grassland	3.5 l/ha	3.5 l/ha per year	Before the crop is 25 cm high
Amenity grassland	3.5 l/ha	7.0 l/ha per year	

Method of application: Tractor mounted / trailed sprayer / knapsack

Restriction on category of user: For Professional Use only

Directions for the safe disposal of the plant protection product and of the packaging

Dispose of contents/container to a licensed waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste

OTHER SPECIFIC RESTRICTION: Livestock must be kept out of treated areas for at least two weeks following treatment and until poisonous weeds such as Ragwort have died and become unpalatable.

Y1UR-P0UV-700A-WR8Q

5 Litre e



Hygeia Chemicals Ltd.
Oranmore, Co. Galway, Ireland.
Tel: 091-794722
Email: enquiries@hygeia.ie Web: www.hygeia.ie

Irish 24 hour emergency number:
National Poisons Centre in Ireland: 01-8092166
Authorisation Holder Nufarm UK Limited
Wyke Lane, Wyke, Bradford, West Yorkshire
BD12 9EJ, UK

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

RESTRICTIONS

Do not contaminate water with the product or its container (Do not clean application equipment near surface water / Avoid contamination via drains from farmyards and roads)

DO NOT graze grass for at least 14 days after spraying.
DO NOT mow or roll four days before or after application.
DO NOT treat newly established grass less than one year old.
DO NOT treat grass crops intended for seed production.
DO NOT treat grass suffering from drought, disease or other adverse factors.
DO NOT roll or harrow crops for seven days either before or after application of HYGRASS XTRA.

DO NOT apply when grassland is flowering.
Avoid spray drift onto neighbouring crops and all broad-leaved plants outside the target area.

Do not spray in windy weather. Beets, all brassicas (including oilseed rape, swedes and turnip) lettuce, sunflowers, onions, peas, potatoes, tomatoes, cucumbers, all fruit crops (including vines) and ornamentals are particularly susceptible and may be damaged by spray drift.

Avoid spraying within five metres of the field boundary to reduce effects on non-target insects or other arthropods.

WEEDS CONTROLLED

Apply when the majority of annual weeds are at the seedling stage.

SUSCEPTIBLE

Thisle, creeping (plants up to 50 cm high)

MODERATELY SUSCEPTIBLE

Buttercup, creeping, Dock, broad-leaved, Dock, curled, Ragwort, common

Susceptible Consistently high levels of control obtained under good growing conditions.

Moderately susceptible Top growth usually killed and a useful measure of long-term control obtained under good growing conditions.

Ragwort control

Spray when the majority of plants are in the rosette stage and growing vigorously in the autumn or spring but before the flower spines start to grow. Fields for hay or silage the following spring should be sprayed in the preceding autumn. Fields to be grazed should be treated in the spring.

Common ragwort will require spraying over a minimum of two years for complete control.

CROP SPECIFIC INFORMATION

Grassland:	Maximum dose rate 3.5 l/ha. Maximum total dose 3.5 l/ha.
Timing	April to September.
Water volume	100–400 litres of water per hectare. For use in knapsack sprayer the minimum water volume is 200 litres per hectare.
Amenity grassland	Maximum dose rate 3.5 l/ha. Maximum total dose 7.0 l/ha.
Timing	April to September Interval between applications Six weeks.
Water volume	100–400 litres of water per hectare. For use in knapsack sprayer the minimum water volume is 200 litres of water per hectare.

MIXING AND SPRAYING

Before use ensure that the spraying equipment has been thoroughly cleaned. Half-fill the spray tank with clean water. With the contents of the spray tank under re-circulation, add the measured quantities of HYGRASS XTRA through the filter. Top up the tank with water to the required level and maintain re-circulation until the tank is sprayed out. Apply the recommended quantity of HYGRASS XTRA through a conventional hydraulic sprayer using a MEDIUM spray to cover the weed leaves evenly and thoroughly. After each day's use wash out with water and wetting agent. Wash out again with water, drain and allow to dry. Traces of herbicide left in the sprayer may damage susceptible crops if these are subsequently sprayed using the same equipment. Avoid spray drift onto neighbouring crops and all broad-leaved plants outside the target area. Do not spray in windy weather. Beets, all brassicas including oilseed rape, lettuce, peas, potatoes, tomatoes, all fruit crops and ornamentals are particularly susceptible to 2,4-D and may be damaged by spray drift.

Knapsack application This method is recommended for spot treatment of weeds and patches of Docks. Apply as a MEDIUM spray to just before run-off occurs.

Add 180ml of Hygrass Xtra to a full 20 Litre Knapsack.

WEATHER AND GROWING CONDITIONS

Apply to a dry crop when rain is not forecast for at least 12 hours. Optimum results are obtained when the weeds are actively growing under good soil and weather conditions. Reduced weed control may be obtained during drought or cold weather. If rain falls shortly after application the effect of HYGRASS XTRA may be reduced.

Resistance Management When herbicides with the same mode of action are used repeatedly over several years in the same field, selection of resistant biotypes can take place. These can propagate and may become dominating. A weed species is considered to be resistant to a herbicide if it survives a correctly applied treatment at the recommended dose. A strategy for preventing and managing such resistance should be adopted. This should include integrating herbicides with a programme of cultural control measures. Guidelines have been produced by the Weed Resistance Action Group and copies are available from the HGCA, CPA, your distributor, crop adviser or product manufacturer.