



TRAILER[®]

HERBICIDE

GROUP 2 HERBICIDE

TRAILER is a water dispersible granules (WG) formulation, containing **750 g/kg of tribenuron-methyl** for the post-emergence control of broad-leaved weeds in wheat, barley, oats, winter rye and triticale.

FOR PROFESSIONAL USE ONLY.

Approval Holder and Marketing Company:

Albaugh TKI d.o.o, Grajski trg 21, 2327 Rače, Slovenia
Tel: +386 2 60 90 211, www.albaugh.eu

Emergency Information:

CARECHEM: +44 1235 239 670 (24 hour)
Technical Enquiries: 1800 901 562



**WARNING
SAFETY INFORMATION**

May cause damage to organs through prolonged or repeated exposure. Very toxic to aquatic life with long lasting effects.

Contains tribenuron-methyl. May produce an allergic reaction.

Do not breathe dust/mist/spray. Wear protective gloves/protective clothing.

IF ON SKIN: wash with plenty of water. Get medical advice/attention if you feel unwell.

Collect spillage. Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for triple rinsed empty, clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

UFI: YD40-Q0G5-N00Y-J6UM

PCS No. 06830

TKI-IE_TRAILER_100g_LBL_18-11-25

Batch number and date of manufacture: see packaging

Net Content:

100 g e

[®] [™] trademark or trade name of Albaugh, LLC or an affiliated company

ALBAUGH
your alternative

IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL HERBICIDE

Crops/Situations	Maximum Individual Dose	Maximum Total Dose	Maximum Number of Treatments	Latest Time of Application
Wheat, barley, oats, winter rye and triticale	20 grams product/ha	20 grams product/ha	1 per crop	Third node detectable (GS 33)

Other Specific Restrictions

Application may only be made from 1st February and when crop is at beginning of tillering (GS 21).

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**SAFETY PRECAUTIONS****Operator Protection**

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE GLOVES when handling the concentrate or handling contaminated surfaces.

WASH ALL PROTECTIVE CLOTHING thoroughly after use, especially the inside of gloves.

WASH SPLASHES from skin or eyes immediately.

WASH HANDS AND EXPOSED SKIN before eating, drinking or smoking and after work.

Environmental Protection

Extreme care must be taken to avoid spray drift onto non-target plants.

Spray must be aimed away from water.

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

Storage and Disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

KEEP OUT OF REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place. Store in a cool dry place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose safely.

DO NOT RE-USE CONTAINER FOR ANY PURPOSE.

DIRECTIONS FOR USE

IMPORTANT: This information is authorised as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

GENERAL INFORMATION

TRAILER is a foliar acting herbicide, for the post-emergence control of broad-leaved weeds in cereals. It can be used on all varieties of wheat, barley, oats, winter rye and triticale. Applications should be made to young, actively growing weeds. Activity is faster under good growing conditions and may appear relatively slow under cold or dry weather conditions.

RESTRICTIONS

Do not spray crops under stress or crops suffering from waterlogging, pest attack, disease or frost.

Do not spray crops undersown with grass mixtures.

Do not apply more than once to any cereal crop.

Do not apply within seven days of rolling the crop.

Special care must be taken to avoid damage by drift onto broad-leaved plants outside the target area, or onto ponds, waterways or ditches. Thorough cleansing of equipment is also very important.

WEED RESISTANCE

TRAILER contains tribenuron-methyl, a sulfonylurea (ALS inhibitor), also classified by the Herbicide Resistance Action Committee as 'Group 2'.

To reduce the risk of developing resistance, applications should be made to young actively growing weeds. The use of cultural methods of control and herbicides with non-ALS modes of action within the cropping season and/or throughout the cropping rotation will help reduce the risk of developing resistance. When herbicides with the same mode of action are used repeatedly over several years in the same field, selection of resistant biotypes can take place. These can propagate and may become dominating. A weed species is considered resistant to a herbicide if it survives a correctly applied treatment at the recommended dose. Development of resistance with a weed species can be avoided or delayed by alternating (or tank mixing) with suitable products having a different mode of action. A strategy for preventing and managing resistance should be adopted. The Herbicide Resistance Action Committee (HRAC) has produced guidelines and copies are available from your distributor, crop advisor or product manufacturer.

SUSCEPTIBLE WEEDS

The susceptibility rating of weeds in the following table refer to good spray cover and young, actively growing weeds at the timing of treatment.

Weed species		Level of susceptibility
Common name	Scientific name	
Corn chamomille	<i>Anthemis arvensis</i>	S
Fat hen	<i>Chenopodium album</i>	S
Creeping thistle	<i>Cirsium arvense</i>	S
Henbit	<i>Lamium amplexicaule</i>	S
Common poppy	<i>Papaver rhoeas</i>	S
Sow-thistle	<i>Sonchus olearaceus</i>	S
Common chickweed	<i>Stellaria media</i>	S
Scarlet pimpernel	<i>Anagallis arvensis</i>	MS
Red dead-nettle	<i>Lamium purpureum</i>	MS
Common fumitory	<i>Fumaria officinalis</i>	R
Speedwells	<i>Veronica sp.</i>	R
S = Susceptible; MS = Moderately Susceptible, R = Resistant		

SOIL AND WEATHER

TRAILER can be used on all soil types. Weed control may be reduced in dry conditions.

VOLUME AND APPLICATION

BEFORE USING TRAILER, SPRAYING EQUIPMENT MUST BE CLEAN AND FREE FROM CONTAMINATION WITH OTHER PESTICIDES.

Spray TRAILER in 200-400 litres of water per hectare. Where there are dense weed populations or advanced growth stages the water volume should be 400 litres per hectare.

Use a conventional field crop sprayer at a pressure of 2-3 bars and apply as a MEDIUM spray (as defined by BCPC). Care should be taken not to overlap spray swaths.

Method of application: Tractor mounted/trailed horizontal boom sprayer.

MIXING

Quarter fill the spray tank with clean water, start the agitation and add the required quantity of TRAILER directly to the tank without prior creaming. Continue agitation while topping up the tank and while spraying.

COMPATIBILITY

TRAILER should be mixed with other herbicides to widen and improve performance for the control of weeds. For information on tank-mixing partners contact your distributor or Albaugh TKI d.o.o.

CROPS

CEREALS

CROP SAFETY

TRAILER can be used on all varieties of the following cereals: wheat, barley, oats, winter rye and triticale.

TIMING

TRAILER must only be applied once per crop up to the third node detectable (GS33). Application may only be made from 1st of February and when the crop is at beginning of tillering (GS21).

DOSE

Apply TRAILER at a maximum rate 20 grams product per hectare.

FOLLOWING CROPS

After harvest of a cereal crop treated with TRAILER, only cereals, oilseed rape, sugar beet, field beans or grass may be sown in the same calendar year.

REPLACEMENT CROPS

In the event of crop failure for any reason, only cereal crops can be sown after an interval of 3 months.

WARNINGS

EXTREME CARE MUST BE TAKEN TO AVOID SPRAY DRIFT ONTO NON-CROP PLANTS OUTSIDE OF THE TARGET AREA.

SPRAYING EQUIPMENT SHOULD NOT BE DRAINED OR FLUSHED ONTO LAND PLANTED WITH OR INTENDED FOR PLANTING WITH TREES OR CROPS OTHER THAN CEREALS.

SPRAY TANK CLEAN-OUT

POOR CLEANOUT PRACTICES AND INSUFFICIENT WATER VOLUMES USED FOR THE RINSE PROCEDURE MAY RESULT IN INADEQUATE REMOVAL OF PRODUCT DEPOSITS. SUBSEQUENT USE OF APPLICATION EQUIPMENT IN THESE CIRCUMSTANCES MAY RESULT IN DAMAGE TO NON-CEREAL CROPS.

TO AVOID SUBSEQUENT DAMAGE TO CROPS OTHER THAN CEREALS, IMMEDIATELY AFTER SPRAYING TRAILER THOROUGHLY CLEAN ALL SPRAY EQUIPMENT, INCLUDING INSIDE AND OUTSIDE OF LID, USING A PROPRIETARY SPRAYER CLEANER FOR USE WITH SULFONYLUREAS ACCORDING TO THE FOLLOWING PROCEDURE:

1. Immediately after spraying, drain tank completely. Wash any contamination off the outside of the sprayer with clean water.
2. Rinse the inside of the tank with clean water and flush at least one tenth of the spray tank volume through the boom and hoses. Drain tank completely.
3. Half-fill the tank with clean water and add the correct quantity of a proprietary sprayer cleaner for use with sulfonyleureas. Agitate and then flush the boom and hoses with the solution. Top up completely with water and allow to stand for 15 minutes with agitation. Again flush the booms and hoses and drain tank completely (if it is not possible to drain the tank completely, repeat step 3 before going on).
4. Remove nozzles and filters and soak in a bucket containing a proprietary sprayer cleaner at the same concentration as that used for the sprayer.
5. Rinse tank again with clean water and flush at least one tenth of the tank volume through the booms and hoses. Drain tank completely.

GENEREAL

Consult label tank cleanup procedures for all tank mix partners and be sure to use the most rigorous procedure recommended.

TERMS AND CONDITIONS OF SUPPLY, SALE AND USE

Many factors can affect or influence the activity of this product, including, but not limited to: weather and soil conditions, crop variety, treatment timing, water volume, application rates, spraying techniques, crop rotation, regional factors, and the occurrence and development of weeds resistant to the active ingredient. Under certain circumstances, changes in activity or crop damage can occur. The manufacturer or supplier is unable to accept any liability in these circumstances. All goods supplied by us are of a high grade and we believe them to be suitable for the purpose for which we expressly supply them: but as we cannot exercise any control over their mixing, use or application which may affect the performance of the goods all conditions and warranties statutory or otherwise as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff, our agents or the re-sellers of the product whether or not they supervise or assist in the use of such goods.