



syngenta®

GROUP **3** **11** FUNGICIDES

Product registration number: PCS No. 06993  
UFI: P2A7-20AG-Y00T-Q658



ANGLE® is a suspension concentrate containing 125 g/L of Azoxystrobin, 125 g/L of Difenoconazole. For the control of black leg (*Leptosphaeria maculans*) and Sclerotinia stem rot (*Sclerotinia sclerotiorum*) on winter oilseed rape, and the control of leaf spot (*Cercospora beticola*), powdery mildew (*Erysiphe betae*), rust (*Uromyces betae*) and *Rhizoctonia solani* on fodder beet and sugar beet.

**SAFETY INFORMATION**

**FOR PROFESSIONAL USE ONLY**

To avoid risks to human health and the environment, comply with the instructions for use.

ANGLE is a suspension concentrate containing 125 g/L of Azoxystrobin, 125 g/L of Difenoconazole.

**Warning**

Harmful if swallowed or if inhaled.

Suspected of causing cancer.

Very toxic to aquatic life with long lasting effects.

Obtain special instructions before use.

Avoid breathing mist or vapours.

Wash skin thoroughly after handling.

Wear protective gloves/ protective clothing/ eye protection/ face protection/ hearing protection.

IF INHALED: Remove person to fresh air and keep comfortable for breathing. Call a POISON CENTER/ doctor if you feel unwell.

Collect Spillage.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except triple rinsed empty clean containers which can be disposed of as non-hazardous waste.

Contains 1,2-benzisothiazol-3(2H)-one. May produce an allergic reaction.



PCS: 06993

UFI: P2A7-20AG-Y00T-Q658

**IN CASE OF TOXIC OR TRANSPORT EMERGENCY RING +44 (0) 1484 53844 (24HR)**

Authorisation Holder	Marketing Company
Syngenta UK Limited, Jealott's Hill International Research Centre, Bracknell, Berkshire, RG42 6EY Tel +44 (0) 1223 883400	Syngenta Ireland Ltd Block 6, Cleaboy Business Park, Old Kilmeaden Road, Waterford, Ireland. Tel: (051) 377203

Lxxxxxxx IREL/10D PPE xxxxxxxx

**5 litres**

Product names marked ® or ™, the ALLIANCE FRAME  
the SYNGENTA Logo and the PURPOSE ICON  
are Trademarks of a Syngenta Group Company

**IMPORTANT INFORMATION**

Crop	Maximum individual dose (L/ha)	Maximum number of treatments (per crop)	Minimum spray interval (days)	Latest time of application
Winter Oilseed rape	1	2		Up to and including end of flowering (BBCH GS69)
Sugarbeet and fodder beet	1	2	21	Up to and including beet root has reached harvestable size (BBCH GS49), and not less than 35 days before harvest

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS**

**Additional Safety Information****(a) Operator protection**

WHEN USING DO NOT EAT DRINK OR SMOKE. **EXPIRES**

WASH SPLASHES from skin and eyes immediately.

DO NOT BREATHE SPRAY.

IF YOU FEEL UNWELL, seek medical advice immediately (show label where possible).

WASH HANDS AND EXPOSED SKIN before meals and after work.

**(b) Environmental protection**

To protect non-target aquatic organisms, respect an unsprayed buffer zone of 5m to surface waters.\* Do not contaminate water with the product or its container. Do not clean application equipment near surface water.

Avoid contamination via drains from farmyards and roads.

\* To reduce this buffer zone please refer to PRCD Guidance - STRIPE (Surface water Tool for Reducing the Impact of Pesticides in the Environment).

**(c) Storage and Disposal**

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times.

Add washings to sprayer at time of filling and dispose of safely.

KEEP IN ORIGINAL CONTAINER, tightly closed in a safe place.

KEEP AWAY FROM FOOD DRINK AND ANIMAL FEEDING STUFFS.

KEEP OUT OF REACH OF CHILDREN.

DO NOT RE-USE CONTAINER FOR ANY PURPOSE

This leaflet is part of the approved Product Label.

## DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be carefully read in order to obtain safe and successful use of this product.

## GENERAL INFORMATION

Angle is a suspension concentrate containing 125 g/l azoxystrobin and 125 g/l difenoconazole. Angle has systemic activity, with protectant properties for use in all commercial varieties of winter oilseed rape, sugarbeet and fodder beet.

## RESTRICTIONS

Certain apple varieties are highly sensitive to azoxystrobin. As a precaution Angle should not be applied when there is a risk of spray drift onto neighbouring apple crops. Spray equipment used to apply Angle to other crops should not be used to treat apples.

## DISEASES CONTROLLED

### Winter Oilseed rape

Sclerotinia stem rot (*Sclerotinia sclerotiorum*)

Stem canker (*Phoma lingam*)

### Sugarbeet and fodder beet

Powdery mildew (*Erysiphe betae*)

Rust (*Uromyces betae*)

Ramularia leaf spot (*Ramularia beticola*)

Cercospora leaf spot (*Cercospora beticola*)

*Rhizoctonia solani*

## CROP SPECIFIC INFORMATION

For the control of black leg (*Leptosphaeria maculans*) and Sclerotinia stem rot (*Sclerotinia sclerotiorum*) on oilseed rape, and the control of leaf spot (*Cercospora beticola*), powdery mildew (*Erysiphe betae*), rust (*Uromyces betae*) and *Rhizoctonia solani* on fodder beet and sugar beet.

Before applying ANGLE, ensure the crop is free from any stress caused by environmental or agronomic effects. Best results will be achieved from applications made as a protectant treatment following a disease risk assessment or the use of appropriate decision support systems.

## WINTER OILSEED RAPE

### Rate of Use

Apply 1 liter per hectare

The maximum number of applications to Winter Oilseed rape is 2 per crop.

Latest time of application - Up to and including end of flowering (BBCH GS69)

### Stem canker (*Phoma lingam*)

**Timing:** Spray protectively with 1 litre of ANGLE per hectare in the Autumn for the control of Stem canker (*Phoma lingam*) from the 4 expanded true leaf stage (GS 14). Where appropriate, a second application can be made in the Spring from the start of stem elongation and no later than individual flower bud visible (GS 30-55).

**Rate of Use:** 1 liter per hectare. Two applications per crop.

### **Sclerotinia stem rot (*Sclerotinia sclerotiorum*)**

**Timing:** Spray one application protectively with 1 litre of ANGLE per hectare in the Spring for the control of Sclerotinia stem rot (*Sclerotinia sclerotiorum*) from yellow bud until the end of flowering (GS 59-69).

**Rate of Use:** 1 litre per hectare

### **SUGARBEET AND FODDER BEET**

#### **Rate of Use**

Apply 1 liter per hectare

The maximum number of applications to Sugarbeet and fodder beet is 2 per crop. Latest time of application - Up to and including beet root has reached harvestable size (BBCH GS49), and not less than 35 days before harvest

A minimum interval of 21 days days must be observed between applications.

**Timing:** Apply ANGLE at the first signs of the disease, before the disease becomes established. ANGLE gives prolonged protection from re-infection, but a second application can be applied where crops are at risk from later attacks. Apply ANGLE only between Crop cover complete: leaves cover 90% of ground and beet root has reached harvestable size (BBCH GS39-49).

#### **MIXING AND SPRAYING**

Ensure that the sprayer is clean and correctly set to give an even application at the required volume. Half-fill the spray tank with clean water and start agitation. Shake the container and add the required amount of ANGLE to the sprayer using a filling device (e.g. induction bowl or closed transfer unit) or by direct addition to the sprayer tank.

Wash out containers thoroughly, preferably using an integrated pressure rinsing device, or manually rinse three times. Add washings to the sprayer at the time of filling. Complete filling to the required volume and continue to agitate throughout the spraying operation.

Do not leave the spray liquid in the sprayer for long periods (such as during meal breaks or overnight).

#### **WATER VOLUME AND SPRAYING**

Apply in a recommended 200-400 litres of water per hectare using a medium quality spray (BCPC) at a pressure of at least 2 bar. Apply through conventional crop spraying equipment.

#### **CLEANING**

Thoroughly wash out sprayer according to the manufacturer's guidelines and dispose of washings and clean containers according to DEFRA Code of Practice and local water authority guidelines.

#### **RESISTANCE MANAGEMENT**

ANGLE contains azoxystrobin, a strobilurin belonging to the Quinone outside inhibitors (QoI) (FRAC group 11) and difenoconazole, a demethylation inhibitor (DMI) fungicide (FRAC group 3). The number of QoI containing applications should be no more than ½ (50%) of the total number of fungicide applications per season.

Strains of Light Leaf Spot resistant to DMI fungicides are known to exist. To avoid development of resistance, apply product protectively in response to disease forecasts and should not be relied upon for its curative potential.

Where possible, when Light Leaf Spot is present, avoid the use of DMI based fungicides when targeting other diseases such as Sclerotinia at mid flowering.

Use ANGLE as part of an Integrated Crop Management (ICM) strategy incorporating other methods of control, including where appropriate other fungicides with a different mode of action.

To avoid the likelihood of resistance developing, application of ANGLE should be made with due

regard to current FRAG-UK guidelines for QoI compounds. Do not apply more than a total of two applications, when used as part of a programme.

#### COMPANY ADVISORY INFORMATION SECTION

This product is to be used only in accordance with the recommendations and instructions given on the labels provided with this pack.

To access the Safety Data Sheet for this product, scan QR code:



Alternatively, contact your supplier

Specimen update  
2025 to date

THESE PAGES CAN BE ADDED OR REMOVED AS NECESSARY

**IMPORTANT INFORMATION**

Crop	Maximum individual dose (L/ha)	Maximum number of treatments (per crop)	Minimum spray interval (days)	Latest time of application
Winter Oilseed rape	1	2		Up to and including end of flowering (BBCH GS69)
Sugarbeet and fodder beet	1	2	21	Up to and including beet root has reached harvestable size (BBCH GS49), and not less than 35 days before harvest

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS**

**Additional Safety Information****(a) Operator protection**

- WHEN USING DO NOT EAT DRINK OR SMOKE.
- WASH SPLASHES from skin and eyes immediately.
- DO NOT BREATHE SPRAY.
- IF YOU FEEL UNWELL, seek medical advice immediately (show label where possible).
- WASH HANDS AND EXPOSED SKIN before meals and after work.

**(b) Environmental protection**

- To protect non-target aquatic organisms, respect an unsprayed buffer zone of 5m to surface waters.\*
- Do not contaminate water with the product or its container. Do not clean application equipment near surface water.
- Avoid contamination via drains from farmyards and roads.
- \* To reduce this buffer zone please refer to PRCD Guidance - STRIPE (Surface water Tool for Reducing the Impact of Pesticides in the Environment).

**(c) Storage and Disposal**

- RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times.
- Add washings to sprayer at time of filling and dispose of safely.
- KEEP IN ORIGINAL CONTAINER, tightly closed in a safe place.
- KEEP AWAY FROM FOOD DRINK AND ANIMAL FEEDING STUFFS.
- KEEP OUT OF REACH OF CHILDREN.
- DO NOT RE-USE CONTAINER FOR ANY PURPOSE