



GOLTIX[®] GOLD



ADAMA



HERBICIDES

Suspension concentrate

PCS 07091

FOR PROFESSIONAL USE ONLY



Ingredients

● Metamitron 700g/L (58.3% w/w)

Crops

A selective herbicide with foliar and root activity for use in sugar beet, beetroot, fodder beet and mangels.



SAFETY INFORMATION

Warning

Harmful if swallowed.

Very toxic to aquatic life with long lasting effects.

Contains 1, 2-Benzisothiazolin-3(2H)-one. May produce an allergic reaction.

Keep out of reach of children.

Do not eat, drink or smoke when using this product.


IF SWALLOWED: call a POISON CENTRE or doctor/physician if you feel unwell.

Rinse mouth.

Wash hands thoroughly after handling.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for triple rinsed, empty containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.



Scan for Safety Data Sheet

Or use the weblink :

<https://bit.ly/35ybrRj>

Alternatively, contact your supplier.



R 3.1 Sep 24

LJ225JIPFR_01

10Le





GOLTIX[®] GOLD



ADAMA



HERBICIDES

Suspension
concentrate

PCS 07091

GROUP 5 HERBICIDES

SHAKE WELL BEFORE USE

PROTECT FROM FROST

In the event of emergency, call the National Poisons Information Centre, Beaumont Hospital on 01 809 2166 or 01 837 9964 (8am - 10pm 7 days a week) or the UK National Chemical Emergency Centre on +44 (0)1865 407333 (24 hours, 7 days a week).

IMPORTANT INFORMATION FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL HERBICIDE

Crops	Maximum individual dose (L product/ha)	Maximum total dose (L product/ha/crop)	Latest time of application
Sugar beet, beet root, fodder beet and mangels	3	3	Pre-emergence (BBCH 09)
	OR		
	2	2	First leaf visible (pinhead-size): cotyledons horizontally unfolded (BBCH 10)
	OR		
	1	5	Before crop leaves meet between the rows (BBCH 39)

Other specific restrictions:
 The minimum interval between applications is 5 days.
 Fodder beet and mangels must not be grazed by livestock or harvested for animal consumption until at least 32 days following the last application.
 Apply no more than 3.5kg of metamitron/ha in any 12 month period.

Authorised to & marketed by:
ADAMA Agricultural Solutions UK Ltd,
Third Floor East,
1410 Arlington Business Park,
Theale, Reading, RG7 4SA

Tel: 01635 860555
Technical Helpline: 01635 876622
www.adama.com
Email: ukenquiries@adama.com

Distributed in Ireland by:
Croplink (Ireland) Ltd,
Springhill, Carrigtwohill,
Co. Cork, T45 F685
Tel: 021 4882806

UFI: 2JHQ-9DFT-8V0K-69KJ



5 056325 503325

R 3.1 Sep 24

Directions For
Use

Open here

L12251IPBB_01



SAFETY INFORMATION

Warning

Harmful if swallowed.

Very toxic to aquatic life with long lasting effects.

Contains 1, 2-Benzisothiazolin-3(2H)-one. May produce an allergic reaction.

Keep out of reach of children.

Do not eat, drink or smoke when using this product.

IF SWALLOWED: call a POISON CENTRE or doctor/physician if you feel unwell.

Rinse mouth.

Wash hands thoroughly after handling.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for triple rinsed, empty containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.



PCS 07091

GOLTIX® GOLD (PCS 07091) is suspension concentrate formulation containing 700g/L (58.3% w/w) metamitron. It is a selective herbicide with foliar and root activity for use in sugar beet, beetroot, fodder beet and mangels.

This leaflet is part of the approved label.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

GOLTIX® GOLD is a versatile, crop safe, selective herbicide with contact and residual properties acting by both leaf and root uptake for the control of annual weeds in sugar beet, beet root, fodder beet and mangels. GOLTIX GOLD is absorbed by the roots of any emerging weeds.

RESTRICTIONS OR WARNINGS

For use by tractor mounted/trailed sprayer only.

Pre-emergence application is not recommended on organic soils, as efficacy may be reduced.

Crops should not be re-entered until spray residues are dry.

Extreme care must be taken to avoid spray drift on to non-crop plants outside of the target area.

RESISTANCE

This product contains metamitron, a Group 5 herbicide, based on the mode of action classification system of the Herbicide Resistance Action Committee.

Repeated use of herbicides with the same mode of action can increase the risk of strains of weeds developing resistance to these compounds, leading to poor control. In order to minimise the risk, a strategy for preventing and

managing such resistance should be adopted.

Growers are advised to apply products containing herbicides with different modes of action in sequence or tank mix where two or more components are active against the target weeds.

Use the recommended rate of GOLTIX GOLD and the correct application timing for the hardest to control weed species present in the field.

The above should be used in conjunction with effective cropping rotation and cultivation techniques e.g. stale seedbed, cultivation and use of non-selective herbicide prior to drilling.

Further guidance on weed resistance management is available from the Herbicide Resistance Action Committee (HRAC) and Weed Resistance Action Group (WRAG). Follow WRAG Guidelines.

WEEDS CONTROLLED

For control of annual broad-leaved weeds and annual meadowgrass.

CROP SPECIFIC INFORMATION

GOLTIX GOLD is recommended for use in all varieties of sugar beet grown on mineral and organic soils and beet root, fodder beet, and mangels grown on mineral soils.

TIME OF APPLICATION

GOLTIX GOLD may be applied either as a pre-emergence application, an application during emergence or as a post-emergence program.

RATE OF USE

Pre-emergence (BBCH 00-09)

Apply a maximum total dose of 3.0 L product/ha in 200-400 L water/ha.

During emergence

Apply 2 L product/ha in 200-400 L water/ha. Apply in combination with 2 L/ha mineral or vegetable oil.

Post-emergence (BBCH 10-39)

Apply a maximum individual dose of 1.0 L product/ha in 200-400 L water/ha up to a maximum total dose of 5.0 L product/ha. The minimum interval between applications is 5 days.

FOLLOWING CROPS

Beet crops may be sown at any time following the use of GOLTIX GOLD.

Providing 16 weeks elapse from the last application of GOLTIX GOLD, winter cereals may be sown in the same season. Any spring crop may be sown in the season following use of GOLTIX GOLD or GOLTIX GOLD with mineral or vegetable oil.

Mouldboard ploughing to a depth of 15 cm followed by thorough cultivation is recommended before planting any crop.

MIXING AND SPRAYING

Thoroughly shake the pack before use.

Add the required quantity of GOLTIX GOLD to the half-filled spray tank with the agitation system in operation and then fill to the required level. Continue agitation at all times during spraying and stoppages until the tank is completely empty. Spray immediately after mixing.

When tank mixes are to be used, each product should be added separately to the tank; GOLTIX GOLD should be dispersed first unless otherwise specified under the Compatibility section.

On emptying the container, RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of container safely.

Pre-emergence of the crop use a FINE-MEDIUM spray and post-emergence a FINE spray as defined by the BCPC

system. Avoid spray drift.

Do not use finer than 80 mesh filters in spray lines or nozzles.

Clean spray equipment thoroughly after use using water or a tank cleaning agent.

COMPATIBILITY

For information on tank mixes, consult ADAMA Agricultural Solutions UK Ltd, your agronomist or your distributor.

DISCLAIMER/CONDITIONS OF SUPPLY

The specified properties of our products and the mode of application stated on this label have been established on the basis of research and experience. Products conform to specification at the time of delivery but, as we exercise no control over their subsequent storage, handling, mixing or use or the weather conditions before, during and after application, all of which may affect the performance of the products, no responsibility or liability will be accepted by us or our re-sellers for any failure in performance, damage or injury to person or property whatsoever arising from the storage, handling, application or use of the products. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in or make recommendations concerning the use of such products. We recommend you contact your dealer to request advice on the suitability of this product for any new and/or unusual growing methods or for new varieties not listed on this label.

Authorised to and marketed by:
ADAMA Agricultural Solutions UK Ltd
Third Floor East
1410 Arlington Business Park
Theale
Reading
UK
RG7 4SA

Tel: + 44 1635 860555
Technical helpline: 00 44 1635 876622
www.adama.com
Email: ukenquiries@adama.com

Distributed in Ireland by:
Croplink (Ireland) Ltd
Springhill
Carrigtwohill
Co. Cork
T45 F685
Ireland
Tel: + (0)21 4882806

GOLTIX® is a registered trademark of a company of the ADAMA Group.

© ADAMA Agricultural Solutions UK Ltd