

BOLSTER FLEX

PCS No. 07107

FUNGICIDE

BOLSTER FLEX is a protectant fungicide for control of scab on apples and pears.

A water dispersible granule formulation containing 700 g/kg (70% w/w) dithianon.

FOR PROFESSIONAL USE ONLY

IMPORTANT INFORMATION

FOR USE ONLY AS A PROFESSIONAL FUNGICIDE

Crop	Maximum individual dose (kg product/ha)	Maximum Number of Applications (per crop)	Latest Time of application
Apples and pears	0.5	6	Up to fruit about 90% of final size GS 79 (no less than 42 days before harvest)

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS

SAFETY PRECAUTIONS

Operator Protection

WHEN USING DO NOT EAT, DRINK OR SMOKE.

Wear suitable protective clothing (coveralls), suitable protective gloves and face protection (faceshield) when handling the product.

Wear suitable protective clothing (coveralls) and suitable protective gloves when handling contaminated surfaces.

Wear suitable protective clothing (coveralls) during application.

IN CASE OF CONTACT WITH EYES, rinse immediately with plenty of water and seek medical advice.

IF SWALLOWED, seek medical advice immediately and show this container or label.

In case of accident or if you feel unwell seek medical advice immediately (show label where possible).

Environmental Protection

To protect aquatic organisms respect an unsprayed buffer zone of 30 m to surface water bodies.

Do not contaminate water with product or its container

(Do not clean application equipment near surface water / Avoid contamination via drains from farmyards and roads).

Aim spray away from water.

Storage and Disposal

KEEP OUT OF REACH OF CHILDREN.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.

THIS MATERIAL AND ITS CONTAINER must be disposed of in a safe way.

KEEP IN ORIGINAL CONTAINER, tightly closed in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

Store well away from seeds, fertilizers and animal feedingstuffs in a safe, dry place designated as an agrochemical store.

Protect from frost.

Approval Holder:

UPL Holdings Coöperatief U.A.
Claudius Prinsenlaan 144 A
Blok A, 4818CP Breda
The Netherlands

Marketing Company:

UPL Europe Ltd.
The Engine Rooms (1st Floor)
Birchwood Park, Warrington
Cheshire, WA3 6YN
UK
Tel: 00 44 1925 819999

For 24 hour emergency information contact: CARECHEM24
+44 (0)1235 239670

Manufacturing date and Batch No.: see packaging

SEE ATTACHED LEAFLET FOR DIRECTIONS FOR USE

RISK AND SAFETY INFORMATION

A water dispersible granule containing 70% w/w of dithianon.



WARNING

Toxic if swallowed.

May cause an allergic skin reaction.

Causes serious eye damage.

Suspected of causing cancer.

Very toxic to aquatic life with long lasting effects.

Obtain special instructions before use.

Avoid breathing dust/fume/gas/mist/vapours/spray.

Wear protective gloves/protective clothing/eye protection/face protection.

If in eyes: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

Immediately call a POISON CENTER or doctor/physician.

Store locked up.

Dispose of contents/container to a licensed waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste

To avoid risks to human health and the environment, comply with the instructions for use.

Repeated exposure may cause skin dryness or cracking.

UFI: 0WRM-S34P-0K2K-6TJF

PCS No. 07107

NET CONTENTS:
5kge



DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

BOLSTER FLEX is a fungicide for use on apples and pears.

1. Restrictions/Warnings

BOLSTER FLEX must not be mixed with lime sulphur or highly alkaline products.
On variety Golden Delicious, do not use BOLSTER FLEX after green cluster.

2. Disease Control

BOLSTER FLEX is a protectant fungicide for the reduction of scab (*Venturia spp.*) on apples and pears.

2.1 Resistance

BOLSTER FLEX contains dithianon, a dicarbonitrile fungicide with broad-spectrum preventative properties. Dithianon is a multi-site inhibitor that acts by modification of sulphydryl (SH) groups found in the cysteine residues of many protein. Because of its multi-site mode of action, BOLSTER FLEX has a low risk of resistance development and there have been no reports of resistance or reduced sensitivity to this fungicide. Ensure that BOLSTER FLEX is used according to the label recommendations and in accordance with FRAC (Fungicide Resistance Action Group) guidelines.

3. Crops

Apples and pears

Time of Application

Apply preventatively from the beginning of bud burst. On variety Golden Delicious, do not apply BOLSTER FLEX after green cluster.

Rate of application

Apply 0.5 kg of BOLSTER FLEX in a minimum of 200 litres water/ha.

Interval: 7-14 days, in periods of high disease pressure. Also use a 7-day spray interval on varieties that are susceptible to scab infection, when carryover is high, or when crop growth is rapid.

Repeat applications until danger of scab attack ceases. Water volumes should be based on the size of the trees and leaf area at application, and sufficient to ensure good coverage.

Number of applications

Apply a maximum of six applications per year.

4. Mixing and spraying

Half fill the spray tank with clean water and start the agitation. Slowly add the required amount of BOLSTER FLEX.

Empty container completely and dispose of safely.

Add the remainder of the water and continue agitation until spraying is complete.

Apply by any low or high volume orchard sprayer. Do not apply by hand-held equipment.