

Revitur®

A suspension concentrate containing 25 g/litre (1.8 % w/w) mefenitrifluconazole and 600 g/litre (44.1 % w/w) sulphur for use as a fungicide for the control of a range of foliar stem and ear diseases in winter and spring cereals.

Risk and Safety Information

Warning

Causes skin irritation.

May cause an allergic skin reaction.

Causes serious eye irritation.

Harmful to aquatic life with long lasting effects.

Wear protective gloves/clothing/eye protection/face protection

Do not breath mist/vapour/spray

Wash contaminated body parts thoroughly after handling.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

If eye irritation persists: Get medical advice/attention.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

Contains: 1,2-benzisothiazol-3(2H)-one and post-reaction mass of 5-chloro-2-methyl-2H-isothiazol-3-one and 2-methyl-2H-isothiazol-3-one (3:1). May cause an allergic skin reaction.

To avoid risks to human health and the environment, comply with the instructions for use.

PCS No. : 07120



® = Registered trademark of BASF

UFI: CD17-4K7X-7008-A6HT

10 L

Supplied by:
BASF Ireland DAC, 4th and 5th
Floors, 2 Stockport Exchange,
Railway Road, Stockport, SK1 3GG,
UK Telephone: 00 44 (0)161 475 3000
Emergency Information
(24 hours freephone): 0049 180 227 3112
Technical Enquiries: 0044 (0)845 602 2553
(office hours)

Authorisation holder:
BASF plc, 4th
and 5th Floors,
2 Stockport Exchange,
Railway Road,
Stockport, SK1 3GG,
UK Telephone:
00 44 (0)161 475 3000



FOR USE ONLY AS A PROFESSIONAL FUNGICIDE, as directed below:

Crops	Maximum individual dose	Maximum total dose (per crop)	Latest time of application
Winter wheat, spring wheat, durum wheat, spelt wheat, winter barley, spring barley and triticale	4 litres product/ha	8 litres product/ha	Inflorescence fully emerged (GS 59)

READ ALL PRECAUTIONS BEFORE USE.

PCS No. : 07120

PRECAUTIONS

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

IN CASE OF CONTACT WITH EYES, RINSE IMMEDIATELY with plenty of water and seek medical advice.

IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible).

This product must not be applied via hand-held equipment.

Do not contaminate water with the product or its container.

Do not clean application equipment near surface water.

Avoid contamination via drains from farmyards and roads.

STORAGE

For professional use only.

DO NOT RE-USE CONTAINER FOR ANY PURPOSE.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times.

Add washings to sprayer at time of filling and dispose of container safely.

Store in a suitable pesticide store. Keep dry and protect from frost.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Revytur is a systemic fungicide with protectant and curative properties for disease control in winter wheat, spring wheat, durum wheat, spelt wheat, winter barley, spring barley and triticale.

Restrictions/Warnings

After spraying, wash out sprayer thoroughly according to manufacturer's guideline and dispose of washings and clean containers according to local authority guidelines.

Disease control

Revytur is a systemic fungicide with protectant and curative properties for disease control in winter wheat, spring wheat, durum wheat, spelt wheat, winter barley, spring barley and triticale as summarised below:

	Winter and spring wheat, durum wheat, spelt wheat	Winter and spring barley	Triticale
Zymoseptoria tritici leaf blotch / <i>Septoria</i> spp. Septoria leaf blotch	MC		MC
<i>Puccinia striiformis</i> , Yellow rust	C		
<i>Puccinia triticina</i> , Brown rust	C		C
<i>Blumeria graminis</i> , Powdery mildew	MC		C
<i>Pyrenophora teres</i> , Net blotch		R	
<i>Rhynchosporium secalis</i> Leaf scald		R	
<i>Ramularia collo-cygni</i> Leaf spot		MC	

C- Control

MC- Moderate control

R- Reduction

Resistance Management

GROUP

3

M02

FUNGICIDES

Revytur should be used in accordance with the instructions for use for the target diseases at the specific growth stages indicated.

Use Revytur as part of an Integrated Crop Management (ICM) strategy incorporating other methods of control, including where appropriate other fungicides with a different mode of action.

Revytur should be used preventatively and should not be relied upon for its curative potential.

There is a significant risk of widespread resistance occurring in *Septoria tritici* populations in the UK. Failure to follow resistance management action may result in reduced levels of disease control.

Revytur contains sulphur- a chemical with multi-site activity belonging to FRAC group M02 and mefenentrifluconazole, DMI fungicide belonging to FRAC Code G1. Resistance to some DMI fungicides has been identified in *Septoria tritici* which may seriously affect the performance of some products.

For further advice on resistance management contact your agronomist or specialist advisor and visit the Fungicide Resistance Action Committee (FRAC) and UK Fungicide Resistance Action Group (FRAG) websites.

Crop Specific Information

Revytur is a systemic fungicide with protectant and curative properties for disease control in winter wheat, spring wheat, durum wheat, spelt wheat, winter barley, spring barley, and triticale.

For best results apply Revytur as a protectant treatment. Applications can be made until inflorescence fully emerged (GS59) in wheat, barley and triticale.

Apply Revytur at 4 l/ha in a recommended water volume of 100-300 litres of water per hectare.

Following Crops

After treating a cereal crop with Revytur, broad bean, carrot, maize, oilseed rape, pea, potato, sugar beet, sunflower, winter barley, winter wheat, may be sown as the following crop. The effect of Revytur on other following crops has not been assessed.

Mixing and Application

Never prepare more spray solution than is required.

Three quarters fill the tank with clean water and start the agitation. To ensure thorough mixing of the product, invert the container several times before opening. Add the required quantity of BAS 768 00 F to the spray tank while re-circulating. Add the remainder of the water and continue agitation until spraying is complete.

When tank mixes are to be used, take due note of any instructions given as to the order of mixing. Each product should be added separately to the spray tank and fully dispersed before the addition of any further product(s).

All tank mixes should be used immediately after mixing.

On emptying the container, rinse container thoroughly by using an integrated pressure rinsing device or by manually rinsing three times. Add washings to tank at time of filling and dispose of container safely.

Application

Apply as a MEDIUM spray, as defined by BCPC.

Sprayer cleaning

Wash sprayer thoroughly immediately after use, using clean water and following the sprayer cleaning guidance provided by the equipment manufacturer.

Compatibility

For details of compatibilities contact your distributor, local BASF representative, the BASF Technical Services Hotline: 0845 602 2553 or visit our website on: www.agricentre.bASF.ie

The following does not form part of the product label under S.I. No. 159 of 2012

With many products there is a general risk of resistance developing to the active ingredients. For this reason a change in activity cannot be ruled out. It is generally impossible to predict with certainty how resistance may develop because there are so many crop and use connected ways of influencing this. We therefore have to exclude liability for damage or loss attributable to any such resistance that may develop. To help minimise any loss in activity the BASF recommended rate should in all events be adhered to.

Numerous, particularly regional or regionally attributable, factors can influence the activity of the product. Examples include weather and soil conditions, crop plant varieties, crop rotation, treatment times, application amounts, admixture with other products, appearance of organisms resistant to active ingredients and spraying techniques. Under particular conditions a change in activity or damage to plants cannot be ruled out. The manufacturer or supplier is therefore unable to accept any liability in such circumstances. All goods supplied by us are of high grade and we believe them to be suitable, but as we cannot exercise control over their mixing or use or the weather conditions during and after application, which may affect the performance of the material, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use; but nothing should be deemed to exclude or restrict any liability upon us which cannot be excluded or restricted under the provisions of any applicable law.

Additional Product Safety Information

The information on this label is based on the best available information including data from test results.

Safety Data Sheet

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



agricentre.bASF.ie/RevYTUR-IE/MSDS

Alternatively, contact your supplier.

Specimen-
2025 to date