TM

FUNGICIDE

Suspension concentrate PCS 07240

PCS 07240

FOR PROFESSIONAL USE ONLY

Ingredients

Also contains: Contains folget and hexamethyl tetramine: may produce an alleraic reaction.

Crops

A fungicide for protection against leaf spot (Zymoseptoria tritici) on wheat and triticale; leaf blotch (Rhynchosporium secalis) on barley and triticale and net blotch (Pyrenophora teres) and ramularia leaf spot (Ramularia collo-cygni) on barley.

SAFETY INFORMATION

Warnina

May cause an allergic skin reaction. Suspected of causing cancer. Very toxic to aquatic life with long lasting effects.

Keep out of reach of children. Obtain special instructions before use. Keep only in original packaging. Wear protective gloves & protective clothing. IF ON SKIN: wash with plenty of soap and water. If exposed or if you feel unwell: call a POISON CENTRE or doctor/physician. Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed, empty containers which can be disposed of as non-hazardous waste. To protect aquatic organisms, respect an unsprayed buffer of 5 metres to surface water bodies.

To avoid risks to human health and the environment, comply with the instructions for use.

Scan for Safety Data Sheet

Or use the weblink : https://bit.ly/48G5g3e Alternatively, contact your supplie

R 3.0 Feb 24



5Le

ADAMA





TM COLORADO

ADAMA

FUNGICIDE

Suspension concentrate

PCS 07240



SHAKE WELL BEFORE USE



SAFETY PRECAUTIONS

Operator Protection

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate or handling contaminated surfaces.

WASH CONCENTRATE from skin or eyes immediately.

WASH HANDS AND EXPOSED SKIN before eating or drinking and after work.

IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible).

Environmental Protection

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmvards and roads.

Storage and Disposal

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

DO NOT RE-USE CONTAINER FOR ANY PURPOSE.

In the event of emergency, call the National Poisons Information Centre, Beaumont Hospital on 01 809 2166 or 01 837 9964 (8am - 10pm 7 days a week) or the UK National Chemical Emergency Centre on +44 (0) 1865 407333 (24 hours, 7 days a week).

IMPORTANT INFORMATION FOR PROFESSIONAL LISE ONLY AS ANIACDICI

FOR PROFESSIONAL USE ONLY AS AN AGRICULI URAL FUNDICIDE			
Crops:	Winter wheat, spring wheat, winter barley, spring barley and triticale		
Maximum individual dose:	1.5 litres per hectare		
Maximum total dose:	3 litres per hectare per crop		
Latest time of application:	Before end of heading/inflorescence fully emerged (GS 59)]	

Authorised to & marketed by: ADAMA Agricultural Solutions UK Ltd Third Floor East, 1410 Arlington Business Park, Theale, Reading, RG7 4SA Tel: +44 (0)1635 860555 Technical Helpline: +44 (0)1635 876622 www.adama.com Email: ukenguiries@adama.com

Distributed in Ireland by: Croplink (Ireland) Ltd Springhill, Carrigtwohill, Co. Cork, T45 F685 Tel: 021 4882806

UFI: P3TI-WEI4-9Q09-FXP6





COLORADO ΔΠΔΜΔ **Suspension** PCS 07240 **FUNGICIDE** 67 concentrate PROTECT FROM FROST FUNGICIDE GROUP SHAKE WELL BEFORE USE **Environmental Protection** SAFETY PRECAUTIONS Do not contaminate water with the product or its container. Do not clean application equipment near **Operator Protection** surface water. Avoid contamination via drains from WEAR SUITABLE PROTECTIVE CLOTHING farmvards and roads. (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate or handling Storage and Disposal contaminated surfaces. KEEP IN ORIGINAL CONTAINER, tightly closed, in a WASH CONCENTRATE from skin or eves safe place. immediately. DO NOT RE-USE CONTAINER FOR ANY PURPOSE. WASH HANDS AND EXPOSED SKIN before eating or

drinking and after work. IF YOU FEEL UNWELL, seek medical advice

immediately (show the label where possible).

In the event of emergency, call the National Poisons Information Centre, Beaumont Hospital on 01 809 2166 or 01 837 9964 (8am - 10pm 7 days a week) or the UK National Chemical Emergency Centre on +44 (0) 1865 407333 (24 hours, 7 days a week).

IMPORTANT INFORMATION

FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL FUNGICIDE

Winter wheat, spring wheat, winter barley, spring barley and triticale Maximum individual dose: 1.5 litres per hectare

. Ó

Use

Maximum total dose: 3 litres per hectare per crop

Latest time of application:

Crops:

Before end of heading/inflorescence fully emerged (GS 59)

Authorised to & marketed by: ADAMA Agricultural Solutions UK Ltd Third Floor East, 1410 Arlington Business Park, Theale, Reading, RG7 4SA Tel: +44 (0)1635 860555 Technical Helpline: +44 (0)1635 876622 www.adama.com Email: ukenguiries@adama.com

Distributed in Ireland by: Croplink (Ireland) Ltd Springhill, Carrigtwohill, Co. Cork, T45 F685 Tel: 021 4882806

R 3.0 Feb 24 **Directions For**



SAFETY INFORMATION

Warning

May cause an allergic skin reaction. Suspected of causing cancer. Very toxic to aquatic life with long lasting effects.

Keep out of reach of children. Obtain special instructions before use.

Keep only in original packaging.

Wear protective gloves & protective clothing.

IF ON SKIN: wash with plenty of soap and water.

If exposed or if you feel unwell: call a POISON CENTRE or doctor/physician.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed, empty containers which can be disposed of as non-hazardous waste.

To protect aquatic organisms, respect an unsprayed buffer of 5 metres to surface water bodies.

To avoid risks to human health and the environment, comply with the instructions for use.

PCS 07240

COLORADO™ (PCS 07240) is a suspension concentrate formulation containing 500 g/l (39.5% w/w) folpet. It contains folpet and hexamethyl tetramine: may produce an allergic reaction.

COLORADO is a fungicide for protection against leaf spot (Zymoseptoria tritici) on wheat and triticale; leaf blotch (Rhynchosporium secalis) on barley and triticale and net blotch (Pyrenophora teres) and ramularia leaf spot (Ramularia collo-cygni) on barley.

This leaflet is part of the approved labe

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Resistance

COLORADO contains folpet, a protectant fungicide from the phthalimide chemical group with multi-site contact activity. For resistance management purposes, its mode of action is classified as M4 under the FRAC* code which indicates low-risk without any signs of resistance or cross-resistance developing.

*FRAC: Fungicide Resistance Action Committee

For current advice on resistance management contact your agronomist or specialist advisor and visit the FRAG-UK website.

Disease control in wheat, barley and triticale

For the reduction of leaf spot (Zymoseptoria tritici) in winter and spring wheat and triticale and leaf blotch (Rhynchosporium secalis) in winter and spring barley and triticale; and moderate control of net blotch (Pyrenophora teres) and ramularia leaf spot (Ramularia collo-cygni) in winter and spring barley. COLORADO may be applied in a programme of 2 suitably timed sprays.

Dose rate

1.5 litre per hectare per application

Septoria leaf spot (Zymoseptoria tritici) in winter and spring wheat and triticale

As COLORADO is a protectant fungicide with contact activity, it is important to apply the first spray before the disease becomes established in the crop (normally GS 30-33). A second spray should be timed to protect new growth from disease and therefore needs to be applied prior to further infection. This is normally 3 weeks after the first application but may be sooner under conditions of rapid growth and high disease pressure. Where disease infection or re infection has occurred prior to application, COLORADO should be mixed with alternative chemistry to provide curative activity (see Compatibility section).

Leaf blotch (Rhynchosporium secalis) in winter and spring barley and triticale; and net blotch (Pyrenophora teres) and ramularia leaf spot (Ramularia collo-cygni) in winter and spring barley

As COLORADO is a protectant fungicide with contact activity, it is important to apply the first spray before the disease becomes established in the crop (normally GS 30-33). A second spray should be timed to protect new growth from disease infection. This is normally 3 weeks after the first application but may be sooner under conditions of rapid growth and high disease pressure. Where disease infection or re infection has occurred prior to application, COLORADO should be mixed with alternative chemistry to provide curative activity (see Compatibility section).

Water volume

Apply COLORADO in a water volume of 200-400 litres per hectare.

Qualified recommendation: lower water volumes of 150 litres per hectare may also be used*; however, these have not been supported by effectiveness or crop safety data.

*or the equivalent of a 1% v/v dilution

As COLORADO has contact activity, it is important to achieve a good coverage of the crop when spraying.

Mixing and spraying

For use by tractor mounted/trailed sprayer only.

Before spraying it is important to check all hoses, filters and nozzles, and to ensure that the sprayer is clean and correctly calibrated to give an accurate application at the correct volume.

Half fill sprayer tank with clean water. Add the required amount of COLORADO to the water and commence agitation. Maintain agitation during the mixing process and continue until the spraying is complete.

On emptying the container, RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of container safely. Add the remainder of the water whilst continuing to agitate the mixture until spraying is complete.

Do not leave in spray tank for long periods without using.

Apply COLORADO as a medium spray (as defined by the BCPC) in cereals.

Thoroughly clean the sprayer and measuring equipment after using COLORADO. Crops should not be re-entered until spray residues are dry.

Compatibility

Consult ADAMA Agricultural Solutions UK Ltd or your distributor for details on the products which may be mixed with COLORADO.

Caution

After spraying a herbicide, sprayer tanks and equipment should be thoroughly cleaned before using COLORADO.

DISCLAIMER/CONDITIONS OF SUPPLY

The specified properties of our products and the mode of application stated on this label have been established on the basis of research and experience. Products conform to specification at the time of delivery but, as we exercise no control over their subsequent storage, handling, mixing or use or the weather conditions before, during and after application, all of which may affect the performance of the products, no responsibility or liability will be accepted by us or our re-sellers for any failure in performance, damage or injury to person or property whatsoever arising from the storage, handling, application or use of the products. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in or make recommendations concerning the use of such products. We recommend you contact your dealer to request advice on the suitability of this product for any new and/or unusual growing methods or for new varieties not listed on this label.

Authorised to & marketed by: Distributed in Ireland by: ADAMA Agricultural Solutions UK Ltd Croplink (Ireland) Ltd Third Floor East Springhill 1410 Arlington Business Park Corrigtwohill Theale Co. Cork Reading T45 F685 RG7 4SA Ireland Tel: 021 4882806 Tel: +44 (0)1635 860555 Technical Helpline: +44 (0)1635 876622 www.adama.com Email: ukenauiries@adama.com

COLORADO[™] is a trademark of a company of the ADAMA Group. © ADAMA Agricultural Solutions UK Ltd