

FEONA

PCS 07314

FEONA contains 300 g/l (25.2% w/w) metrafenone as a suspension concentrate. It is used as a fungicide for mildew for use in winter and spring wheat, winter and spring barley, winter and spring oats, rye, triticale and durum wheat.

A mildew fungicide with protectant and certain curative properties for use in winter and spring wheat, durum wheat, winter and spring barley, winter and spring oats, rye and triticale.

A suspension concentrate containing 300 g/L (25.2% w/w) metrafenone.

Risk and Safety Information
Warning

May cause an allergic skin reaction.

Causes serious eye irritation.

Toxic to aquatic life with long lasting effects.

Wash contaminated body parts with water and soap thoroughly after handling. Wear eye/face protection.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Call a POISON CENTRE or doctor/physician. Collect spillage. Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

PCS 07314



FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE

Crops	Maximum Individual Dose	Maximum Total Dose	Latest Timing of Application
Winter wheat, Spring wheat, Durum wheat, Winter barley, Spring barley, Winter oats, Spring oats, Rye and Triticale	0.5 litre product per hectare	1.0 litre product per hectare	Up to and including beginning of flowering (GS 61)

Do not apply more than two applications of products containing metrafenone to any one crop.

READ ALL PRECAUTIONS BEFORE USE.

PCS 07314

UFI: UU22-90VT-N00U-8KAP

PRECAUTIONS

Operator protection WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES WHEN HANDLING THE CONCENTRATE. WASH HANDS AND EXPOSED SKIN before meals and after work. WHEN USING DO NOT EAT, DRINK OR SMOKE. IF YOU FEEL UNWELL, seek medical advice (show label where possible).

Environmental protection Do not contaminate water with the product or its container. (Do not clean application equipment near surface water / Avoid contamination via drains from farmyards and roads.

Storage KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

KEEP OUT OF REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times.

Add washings to sprayer at time of filling and dispose of safely.

DO NOT RE-USE CONTAINER FOR ANY PURPOSE.

Store in a suitable pesticide store. Keep dry and protect from frost.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

CROPS / DISEASE CONTROL: FEONA is a fungicide for use on all varieties of winter and spring wheat, durum wheat, winter and spring barley, winter and spring oats, rye and triticale.

DISEASE	CROPS	CONTROL
Powdery Mildew	Winter and spring wheat Durum wheat, Rye, Triticale, Oats	Protectant properties Moderate control in the latent phase
Powdery Mildew	Winter and spring barley	Protective properties, Moderate control
Eyespot	Wheat	Offers useful reduction of eyespot when applied at GS30-32

FEONA offers best results when applied at the first signs of the disease. Best time of application for wheat, barley, rye, triticale and oats is at the beginning of flowering, no more than two applications can be made. Apply FEONA at a rate of 0.5 litres of product in a minimum of 200 litres of water per hectare.

The following crops can be planted after treating a cereal crop with FEONA: cereals, oilseed rape, sugar beet, linseed, maize, clover, field beans, peas, turnips, carrots, cauliflower, onions, lettuce and potatoes.

RESISTANCE MANAGEMENT

FEONA belongs to group 50 fungicides. To reduce the risk of developing resistance apply the product as instructed, at the correct timing, rate and for the target diseases. Do not apply more than 2 applications per crop. Avoid consecutive applications except where the product is used in tank mixes or co-formulations with other products for the same disease but with different mode of action.

Use as part of a resistance management strategy which includes both chemical and non-chemical methods.

MIXING AND SPRAYING: Ensure the spray tank is half full with clean water. Begin agitation and add the correct amount of FEONA. Maintain agitation when filling the tank and during spraying. Add the remaining water to the spray tank before spraying.

Use MEDIUM spray quality as stipulated by BCPC.

TANK MIXES

Always read and follow the instructions regarding the order of mixing. Add each product separately and ensure it is fully dispersed before adding a further product. For advice regarding compatibility, contact your agronomist or distributor.

CLEANING AND DISPOSING OF EMPTY CONTAINERS

Once empty, the containers must be rinsed thoroughly by using an integrated pressure rinsing device or manually rinsing three times. The washings must be added to the spray tank at the time of filling. Dispose of container safely and according to the current regulations.

In the event of an emergency contact 01 8092166

This product has been re-labelled / re-bottled at an approved packing facility.

Authorisation Holder:

Pan Agriculture Ltd, 8 Cromwell Mews, Station Road,
St Ives, Cambridgeshire, PE27 5HJ

Tel: 01480 467790 Email: info@panagriculture.co.uk

5 Litres e