



CONCERTINA™

A protectant fungicide with curative properties for use in winter wheat, winter barley, spring wheat, spring barley, winter oats and spring oats, rye and triticale.

An emulsifiable concentrate (EC) containing 200 g/litre (19.2% w/w) pyraclostrobin.

Pyraclostrobin is a strobilurin fungicide.

For professional use only. For use only as an agricultural fungicide.

GROUP

11


FUNGICIDE

Risk and Safety Information

Harmful if swallowed.
May be fatal if swallowed and enters airways.
Causes skin irritation.
May cause an allergic skin reaction.
Causes serious eye irritation.
Harmful if inhaled.
May cause respiratory irritation.
Very toxic to aquatic life with long lasting effects.
Use only outdoors or in a well-ventilated area.
Wear protective gloves/protective clothing/eye protection.
IF IN EYES: Rinse cautiously with water for several minutes.
Remove contact lenses, if present and easy to do. Continue rinsing.
Call a **POISON CENTRE** or doctor/physician if you feel unwell.
Store in a well-ventilated place. Keep container tightly closed.
Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.
To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies.
To avoid risks to human health and the environment, comply with the instructions for use.

In the event of emergency, call the National Poisons Information Centre, Beaumont Hospital at 01 809 2166 or 01 837 9964.

DANGER



UFI: M86R-F1SW-T007-T9U8

PCS No. : 07446

Manufacturer: **Barclay Chemicals Manufacturing Ltd.**,
Damastown Way, Damastown Industrial Park, Mulhuddart, Dublin 15, Ireland
Tel: +353 1 811 2900 Fax: +353 1 822 4678
E-mail: info@barclay.ie Website: www.barclay.ie

Approval Holder: **Barclay Chemicals (R&D) Ltd.** Contact details as above.

Copyright © Barclay Chemicals (R&D) Ltd. 2025.

CONCERTINA is a trademark of Barclay Chemicals (R&D) Ltd

5 Litres e

**FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE,
As directed below:**

Crops	Max single dose	Max. no. of applications	Max. total dose	Latest time of application
Winter wheat, spring wheat, rye and triticale	1.25 L/ha	2 per season	2.5 L/ha	Before GS 71 (Watery ripe stage)
Winter barley, spring barley, winter oats and spring oats	1.25 L/ha	2 per season	2.5 L/ha	Before GS 59 (End of heading)

Specific diseases controlled:

Wheat, rye & triticale: Control of brown rust and yellow rust

Barley: Control of brown rust, yellow rust and net blotch; moderate control of Rhynchosporium

Oats: Control of crown rust

PRECAUTIONS

For professional use only.

WEAR SUITABLE PROTECTIVE GLOVES when handling the concentrate. AVOID ALL CONTACT WITH SKIN.

WASH HANDS AND EXPOSED SKIN before meals and after work. WHEN USING DO NOT EAT, DRINK OR SMOKE.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS. KEEP OUT OF REACH OF CHILDREN.

VERY TOXIC TO AQUATIC ORGANISMS, may cause long-term adverse effects in the aquatic environment. Do not contaminate surface waters or ditches with chemical or used container.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely. IF YOU FEEL UNWELL, seek medical advice (show label where possible).

Do not contaminate water with the product or its container. (Do not clean application equipment near surface water / Avoid contamination via drains from farmyards and roads). To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies.

STORAGE

Store in a suitable pesticide store. Keep dry and protect from frost.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Apply as a MEDIUM spray, as defined by BCPC.

CONCERTINA is a fungicide with protectant and curative properties for disease control in winter and spring wheat, winter and spring barley, winter and spring oats, rye and triticale, as summarised below:

	Winter Wheat	Spring Wheat	Winter Barley	Spring Barley	Winter and Spring Oats	Rye	Triticale
Yellow rust	C	C	C	C		C	C
Brown rust	C	C	C	C		C	C
Crown rust					C		
Net blotch			C	C			
<i>Rhynchosporium</i>			MC	MC		MC	

C = Control

MC = Moderate Control

Yield response may be obtained in the absence of visual disease.

Time of Application

Apply **CONCERTINA** at the start of foliar disease attack. A maximum of two applications can be made up to and including flowering just complete in winter and spring wheat, rye and triticale and up to and including emergence of ear just complete in winter and spring barley and winter and spring oats.

Rate of Application

Apply 1.25 litres **CONCERTINA** in a minimum of 200 litres of water per hectare.

Resistance Management

- A maximum of 2 applications of any strobilurin type (QoI) product can be made to any cereal crop.
- In cereals this product must always be applied in a mixture with an approved fungicide that has a different mode of action. Both products must be applied at recommended rates.

Important Notes

1. Avoid spray drift on to neighboring crops.
2. Wash equipment thoroughly after use.

Mixing and Application

Method of application: Tractor mounted/trailed horizontal sprayer

Mixing: Half fill the spray tank with clean water and start the agitation. SHAKE THE CONTAINER WELL before use and pour in the required amount of product. Rinse any empty containers thoroughly and add rinsings to the spray tank. Add the remainder of the water and continue agitation until spraying is completed.

When tank mixes are to be used, each product should be added separately to the spray tank.