

LOSKA® FUNGICIDE

GROUP 3 FUNGICIDE

F

For the control of certain diseases in approved cereal crops. See detailed label text for specific information.

WINTER AND SPRING WHEAT
WINTER AND SPRING TRITICALE
DURUM WHEAT
RYE AND SPELT

PCS No. 07556

5L

NET CONTENT:

life scientific®

FIRST TO MARKET

FOR USE ONLY AS A PROFESSIONAL FUNGICIDE

DANGER



Harmful if swallowed.
Causes skin irritation.
May cause an allergic skin reaction.
Causes serious eye damage.
May cause respiratory irritation.
Suspected of damaging the unborn child.
Toxic to aquatic life with long lasting effects.
Keep out of reach of children.
Avoid breathing spray.



Wash hands thoroughly after handling.
Wear protective gloves/protective clothing/eye protection/face protection.
IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
Immediately call a POISON CENTRE/doctor.
IF exposed or concerned: Get medical advice/attention.



Collect spillage.
Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.
To avoid risks to human health and the environment, comply with the instructions for use.

PCS No. 07556

UFI Code: 9JDD-7AG2-120V-K3X7

xxxxxxxxxx_V1

PEEL BACK FOR DIRECTIONS FOR USE LEAFLET

LOSKA® – PCS No. 07556

Contains PROTHIOCONAZOLE and METCONAZOLE 100 + 48 g/L formulated as an emulsifiable concentrate (EC).

Fungicide for selective use in winter and spring wheat, winter and spring triticale, durum wheat, rye and spelt.

Important Information - FOR USE ONLY AS A PROFESSIONAL FUNGICIDE

Crops	Maximum individual dose (L product per hectare)	Maximum number of treatments	Latest time of application
Winter and spring wheat	1.0	1	BBCH 69
Winter and spring triticale	1.0	1	BBCH 69
Durum wheat	1.0	1	BBCH 69
Spelt	1.0	1	BBCH 69
Rye	1.0	1	BBCH 69

READ THE LABEL BEFORE USE.



APPROVAL HOLDER AND MARKETING COMPANY:

Life Scientific Limited,
Block 4, Belfield Office Park, Beech Hill Road, Dublin 4, Ireland.
Tel: +353 2832024

In the event of emergency, call the National Poisons Information Centre,
Beaumont Hospital at 01 809 2166 or 01 837 9964

Production date / Batch number: see packaging

PROTECT FROM FROST

MADE IN EU

SHAKE WELL BEFORE USE



1/xxxxxx

SAFETY PRECAUTIONS

Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND EYE AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

Avoid contact during pregnancy and while nursing.

However, engineering controls may replace personal protective equipment if an assessment shows that they provide an equal or higher standard of protection.

WHEN USING DO NOT EAT DRINK OR SMOKE.

WASH HANDS AND EXPOSED SKIN before eating and drinking and after work.

Environmental protection

DO NOT CONTAMINATE SURFACE WATERS OR DITCHES with chemical or used container.

Do not contaminate water with the product or its container. (Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads).

Storage and disposal

KEEP OUT OF REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of container safely.

Do not reuse container for any other purpose.

PROTECT FROM FROST.

DO NOT STORE IN DIRECT SUNLIGHT.

Resistance strategy

LOSKA® contains active substances in the Fungicide Resistance Action Committee mode of action Group 3.

Fungicide resistance in *Septoria* to some fungicides with this mode of action has been observed. Repeated applications of products with this mode of action should not be used against medium risk pathogens such as *Septoria*. Tank-mixtures or alternations with products with other modes of action have been demonstrated to protect against the development of disease resistance.

SPECIFIC FRAC RECOMMENDATIONS:

Repeated applications of DMI fungicides alone should not be used on the same crop in one season against a high-risk pathogen in areas of high disease pressure.

Split and reduced rate programmes using multiple repeated applications at dose rates below the manufacturers' recommendations must be avoided.

Curative applications should be avoided.

Restrictions:

Ensure the sprayer is free from residues from previous applications.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

LOSKA® is a selective systemic fungicide, absorbed by the foliage, with penetrant, local and acropetal systemicity. One application may be made to any individual crop as part of a planned disease control programme.

Application

Reduction of Septoria

For reduction of Septoria, apply as soon as the disease is seen in the crop but before disease is well established. Application may be made any time from crop BBCH 30 to BBCH 69. To protect the yield-producing upper leaves spray from BBCH 37-BBCH 65, as part of a planned programme.

Control of Brown Rust

For optimum control of Brown rust, apply as soon as the disease is seen in the crop but before disease is well established. Application may be made any time from crop BBCH 30 to BBCH 69.

Reduction of Fusarium

Reduction of Fusarium ear blight can be achieved by applying LOSKA® after ear emergence but before the end of flowering (crop BBCH 55-69).

Diseases Controlled

The following diseases are susceptible to LOSKA®. The susceptibility ratings of diseases in the following table refer to good spray coverage and good growing conditions.

Disease controlled	Common name	Susceptibility
<i>Fusarium sp.</i>	Fusarium	R
<i>Septoria tritici</i>	Septoria leaf spot	R
<i>Puccinia recondita</i>	Brown rust	S

R = Reduction

S = Susceptible

PREPARATION OF THE SPRAY SOLUTION AND APPLICATION

LOSKA® is to be applied through tractor mounted/ trailed horizontal boom sprayers. It is not to be applied through backpack or hand-held sprayers.

Shake the container before use.

Half-fill the spray tank with clean water, add the necessary amount of LOSKA® and agitate while filling the tank.

Rinse container thoroughly by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of container safely.

Do not leave spray solution in the tank overnight.

Water volume

Apply LOSKA® in 200-400 litres of water per hectare.

Good spray solution is necessary for effective control.

Apply at 2-3 bar, medium spray quality, through properly calibrated equipment.

COMPATIBILITY

LOSKA® is a fully optimised formulation; no additional surfactant is required.

Please contact your distributor for further information regarding tank-mix partners.

SPRAY TANK CLEANING

Wash all equipment immediately after use, using clean water and a suitable cleaning agent, according to the manufacturer's instructions.

CONDITIONS OF SUPPLY

All goods supplied by the company are of good quality and we believe them to be fit for purpose. However, as we cannot exercise control over their storage, handling, mixing or use or the weather conditions before, during or after application, which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded, and no responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

LOSKA® – PCS No. 07556

Contains PROTHIOCONAZOLE and METCONAZOLE 100 + 48 g/L formulated as an emulsifiable concentrate (EC).

Fungicide for selective use in winter and spring wheat, winter and spring triticale, durum wheat, rye and spelt.

Important Information - FOR USE ONLY AS A PROFESSIONAL FUNGICIDE

Crops	Maximum individual dose (L product per hectare)	Maximum number of treatments	Latest time of application
Winter and spring wheat	1.0	1	BBCH 69
Winter and spring triticale	1.0	1	BBCH 69
Durum wheat	1.0	1	BBCH 69
Spelt	1.0	1	BBCH 69
Rye	1.0	1	BBCH 69

READ THE LABEL BEFORE USE.



APPROVAL HOLDER AND MARKETING COMPANY:

Life Scientific Limited,
Block 4, Belfield Office Park, Beech Hill Road, Dublin 4, Ireland.
Tel: +353 2832024

In the event of emergency, call the National Poisons Information Centre,
Beaumont Hospital at 01 809 2166 or 01 837 9964

Production date / Batch number: see packaging

PROTECT FROM FROST

MADE IN EU

SHAKE WELL BEFORE USE



xxxxxxxxxx_V1