

EXTENSION OF AUTHORISATION FOR MINOR USES (EAMU)**INSTRUCTIONS FOR USE****PPP: Benevia 10 OD (PCS No. 05354)**

CROPS: Outdoor grown: Swede & turnip, protected strawberry

This approval provides for the use of the product **Benevia 10 OD (PCS No. 05354)** in respect of, the crops and situations as listed above, which are extra to those specified on the product label.

The conditions of application outlined below are statutory. They must be complied with when availing of this extension for minor uses.

The instructions provided in this extension for minor uses must be used in conjunction with information provided on the approved product label.

Failure to abide by the conditions of approval specified below or on the product label may constitute a breach of the product approval.

Reference Date: 12.07.18.

[Includes latest changes. This version should be used until updated]

| | |
|------------------------------|--------------------------------------|
| <i>Product name:</i> | <u>Benevia 10 OD</u> |
| <i>Active Substance:</i> | Cyantraniliprole |
| <i>Formulation:</i> | 100g/L oil dispersion OD |
| <i>Authorisation Holder:</i> | Headland Agrochemicals Ltd |
| <i>Marketing Company:</i> | Headland Agrochemicals Ltd |
| <i>PCS No:</i> | 05354 |
| <i>User type/Function:</i> | Professional Use Only / Insecticide. |

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

Application:

| Crop | Max. individual dose | Max. number of applications | Max. total dose | Minimum interval between applications | Latest time of application | Target |
|----------------------------------|----------------------|-----------------------------|-----------------|---------------------------------------|----------------------------|---|
| Outdoor grown: Swede & turnip | 0.75 l/ha | 2 | 1.5 l/ha | 14 days | 14 days before harvest | Cabbage root fly, whitefly, Silver Y, Cabbage & Diamondback moths. Cabbage white & Garden Pebble butterflies. |
| Protected strawberry | 0.75 l/ha | 2 | 1.5 l/ha | 7 days | 1 day | Spotted wing drosophila |

Other specific restrictions:

1. To avoid risks to human health & the environment comply with the terms of this approval in conjunction with the directions for use on the product label.
2. For any individual crop the maximum total dose refers to the total dose of any product containing the same active ingredient.
3. Application to outdoor crops by tractor mounted/trailed sprayer only. Application to protected crops may also be by way of hand-held lance.
4. Post application, do not allow unprotected persons to enter treated areas until the crop is completely dry & has been well ventilated.
5. A maximum in use concentration of 5 g cyantraniliprole /hL must not be exceeded
6. For environmental information refer to the product label but specifically: To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies from tractor mounted/trailed sprayers. Aim spray away from water. Growers may be eligible to reduce the buffer zone under the PRCD STRIPE Scheme.
7. This approval extends to professional use only.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

8. For operator protection refer to the product label in conjunction with the specific restrictions above.
9. Adverse effects: Approval holders must immediately submit any new information on the potentially dangerous effects of a product or residues of an active substance contained in a product, on human or animal health, or the environment.

Advisory information:

This approval relates to use of Benevia 10 OD as a horticultural insecticide for use on the crops & situations listed above.

Resistance Management:

Use this product as part of an Integrated Crop Management (ICM) strategy incorporating other methods of control where possible.

Total reliance on one pesticide will hasten the development of resistance. Pesticides of different chemical groups or alternative control measures should be included in the planned anti-resistance strategy. This product should be used in conjunction with other insecticides from different cross-resistant groups [different mode of action, IRAC group] with satisfactory efficacy against the target.

Cyantraniliprole is classed IRAC group 28.

IMPORTANT PLEASE NOTE:

Prior to the use of this product, growers should take advice from their crop advisors particularly in relation to the use of this product within an integrated crop management system of production.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.