# **EXTENSION OF AUTHORISATION FOR MINOR USES (EAMU)**

### **INSTRUCTIONS FOR USE**

PPP: Frupica SC (PCS No. 02128)

**CROPS:** 

[Indoor grown: Tomato, Cucumber, Courgette & Aubergine]

For this EAMU the following crop definitions apply:

Tomatoes includes vine and cherry types.

This approval provides for the use of the product <u>Frupica SC (PCS No. 02128)</u> in respect of, the crops and situations as listed above, which are extra to those specified on the product label.

The conditions of application outlined below are statutory. They must be complied with when availing of this extension for minor uses.

The instructions provided in this extension for minor uses must be used in conjunction with information provided on the approved product label.

Failure to abide by the conditions of approval specified below or on the product label may constitute a breach of the product approval.

**Reference Date: 04.08.2021.** [Includes latest changes. This version should be used until updated]

Product name: Frupica SC

Active Substance: Mepanipyrim

Formulation: 440 g/L SC

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

Authorisation Holder: Certis Belchim b.v

Marketing Company: Certis Belchim b.v

*PCS No:* 02128

*User type/Function:* Professional Use Only / Fungicide.

### Application:

| Crop        | Max.       | Max.         | Max.       | Minimum      | Latest      | Target  |
|-------------|------------|--------------|------------|--------------|-------------|---------|
|             | individual | number of    | total dose | interval     | time of     |         |
|             | dose       | applications |            | between      | application |         |
|             |            |              |            | applications |             |         |
| Protected   | 900 ml/ha  | 2 per crop   | 1.8 L/ha   | 7 days       | 1 day       | Powdery |
| crops of:   | *          |              |            |              | before      | mildew  |
| tomato,     |            |              |            |              | harvest     |         |
| cucumber,   |            |              |            |              |             |         |
| aubergine & |            |              |            |              |             |         |
| courgette.  |            |              |            |              |             |         |

<sup>\*</sup> Product should be used with suitable wetting agent.

## Other specific restrictions:

- (1) To avoid risks to human health & the environment comply with the terms of this approval in conjunction with the directions for use on the product label.
- (2) Use in a water volume of 200 to 1000L except when applying by handheld sprayer where a maximum concentration of 90ml product per 100L water is permitted
- (3) For any individual crop the maximum total dose refers to the total dose of any product containing the same active ingredient.
- (4) Application by tractor mounted/trailed/air assisted/ gantry glasshouse or handheld sprayer only.
- (5) Post application do not allow unprotected persons to enter treated areas until the crop is completely dry & has been well ventilated.
- (6) For environmental information refer to the product label.
- (7) This approval extends to professional use only.
- (8) For operator protection refer to the product label in conjunction with the specific restrictions above.
- (9) Adverse effects: Approval holders must immediately submit any new information on the potentially dangerous effects of a product or residues of an active substance contained in a product, on human or animal health, or the environment.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

# Advisory information:

This approval relates to the use of Frupica SC as a horticultural fungicide for the control of powdery mildew on the above mentioned crops and situations.

## Resistance Management:

Total reliance on one pesticide will hasten the development of resistance. Pesticides of different chemical groups or alternative control measures should be included in the planned anti-resistance strategy. This product should be used in conjunction with other fungicides from different cross-resistant groups [different mode of action, FRAC group] with satisfactory efficacy against the target.

Frupica SC is an anilino-pyrimidine FRAC Group 9 fungicide. Do not use more than two applications of Frupica SC or other anilino-pyrimidine containing products per crop.

#### IMPORTANT PLEASE NOTE:

Prior to the use of this product, growers should take advice from their crop advisors particularly in relation to the use of this product within an integrated crop management system of production.