

EXTENSION OF AUTHORISATION FOR MINOR USES (EAMU)

INSTRUCTIONS FOR USE

PPP: Gazelle SG (PCS No. 03248)

CROPS:

[Field & protected crops of: Lettuce*, Parsley**]

[Field crops of: Carrot, Parsnip, Jerusalem Artichoke, Parsley Root, Radish, Beetroot, Swede, Turnip, Horseradish*** and Celeriac]

For this EAMU the following crop definitions apply:

- * Lettuce includes: Head & Leaf varieties of *Lactuca sativa*: Butterhead, Oak, Lolla Rossa, Cos-Romaine, Batavia, Iceberg, Lambs lettuce (*Valerianella locusta*), Endives, Escarole, Multileaf types, Radicchio and Rocket: (all rocket varieties of *Eruca sativa*: herb, salad, wild & baby leaf rockets) Baby leaf lettuce & baby leaf brassica production (up to 8 true leaves).
- ** Parsley: Leaf parsley only.
- *** Horseradish crops include: Dandelion roots, gentiana roots, East Indian galangal/aromatic ginger/sand galangal, finger root, galangal roots, ginger roots, greater galangal, lesser galangal/smaller galangal, orris roots, wasabi roots.

This approval provides for the use of the product **Gazelle SG (PCS No. 03248)** in respect of, the crops and situations as listed above, which are extra to those specified on the product label.

The conditions of application outlined below are statutory. They must be complied with when availing of this extension for minor uses.

The instructions provided in this extension for minor uses must be used in conjunction with information provided on the approved product label.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

Failure to abide by the conditions of approval specified below or on the product label may constitute a breach of the product approval.

Reference Date: 18.07.23. [Includes latest changes. This version should be used until updated]

Product name: Gazelle SG.
Active Substance: (20%w/w Acetamiprid)
Formulation: Water soluble granule.
Authorisation Holder: Nisso Chemical Europe GmbH
Marketing Company: Certis Belchim nv
PCS No: 03248.
User type/Function: Professional Use / Insecticide.
Application:

Crop	Max. individual dose	Max. number of applications	Max. total dose	Minimum interval between applications	Latest time of application	Target pest
Lettuce (field & protected), parsley (field and protected)	250g/ha	2 per crop	500g/ha	-	3 days pre-harvest	Lettuce currant aphid <i>Nasonovia ribisnigri</i>
Turnips, swedes	250 g/ha	2 per crop	500 g/ha	-	14 days pre-harvest	See target pest table below
Carrot, parsnip, horseradish, celeriac, Jerusalem artichoke, parsley root, radish & beetroot	200 g/ha	1 per crop	200 g/ha		28 days pre-harvest, GS 41 & before 31 st July	See target pest table below

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

Target pests in root crops

Melon cotton aphid (*Aphis gossypii*), peach potato aphid (*Myzus persicae*), Black bean aphid (*Aphis fabae*), turnip aphid (*Hyadaphis pseudobrassicae*) and mealy cabbage aphid (*Brevicoryne brassicae*) on turnip, swede, beetroot, parsley root, Jerusalem artichoke and celeriac
Willow-carrot aphid (*Cavariella aegopodii*) and peach potato aphid (*Myzus persicae*) on carrots
Willow-parsnip aphid (*Cavariella Theobaldi*) and parsnip aphid (*Cavariella pastinacae*) on parsnips
Cabbage root fly (*Delia radicum*) on swede, turnip, radish and horseradish

Other specific restrictions:

- (1) To avoid risks to human health & the environment comply with the terms of this approval in conjunction with the directions for use on the product label.
- (2) For any individual crop the maximum total dose refers to the total dose of any product containing the same active ingredient.
- (3) Application by tractor mounted/trailed horizontal boom sprayer or knapsack.
- (4) Post application, do not allow unprotected persons to enter treated areas until the crop is completely dry & has been well ventilated.
- (5) For environmental information refer to the product label but specifically: To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies from tractor mounted/trailed sprayers. To protect aquatic organisms respect an unsprayed buffer zone of 1m to surface water bodies from knapsack sprayers. Aim spray away from water. Growers may be eligible to reduce the buffer zone under the PRCD STRIPE Scheme.
- (6) This approval extends to professional use only.
- (7) For operator protection refer to the product label in conjunction with the specific restrictions above.
- (8) Adverse effects: Approval holders must immediately submit any new information on the potentially dangerous effects of a product or residues of an active substance contained in a product, on human or animal health, or the environment.

Advisory information:

This approval relates to the use of “Gazelle SG” as a horticultural insecticide for the uses listed in the application table above.

Applications should be made using conventional hydraulic sprayers or hand held applicators using a minimum of 200 litres of water per ha.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

Resistance Management:

For resistance purposes *Acetamiprid* is classed within the group: Neonicotinoid IRAC group 4A. Use no more than two applications of any neonicotinoid insecticide on any crop.

Total reliance on one pesticide will hasten the development of resistance. Pesticides of different chemical types or alternative control measures should be included in the planned programme. Alternating insecticides with different modes of action is a recognised anti-resistance strategy and Gazelle SG should always be used in conjunction with other insecticides with a different mode of action. This includes consideration of sequences involving seed treatments followed by foliar sprays.

IMPORTANT PLEASE NOTE:

Prior to the use of this product, growers should take advice from their crop advisors particularly in relation to the use of this product within an integrated crop management system of production.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.