EXTENSION OF AUTHORISATION FOR MINOR USES (EAMU)

INSTRUCTIONS FOR USE

PPP: Infinito (PCS No. 04309)

CROPS:

[Indoor and Outdoor crops of lettuce*, cabbage*, broccoli, calabrese, Brussel sprout, kale, collard, cauliflower, bulb onion, garlic, shallot, leek & salad onion] [Outdoor crops of spinach, baby leaf production*, herbs*]

For this EAMU the following crop definitions apply:

*Cabbage includes: Round, savoy, pointed, red & white head cabbage.

* Lettuce includes: butterhead, iceberg, oaks, little gem, apollo, lolla rossa, verde, batavia, salanova, endive/frise, escarole, cress, lambs lettuce, red mustard, multileaf types, rocket and baby leaf lettuce up to 8 true leaf stage.

*Baby leaf production includes: baby leaf lettuce and baby leaf brassicas up to 8 true leaves

*Herbs includes: angelica, balm, basil, bay, caraway leaves, celery leaves, chervil, chives, clary sage, coriander leaves, dill leaves, edible flowers, fennel leaves, hyssop, lemon grass, lovage, marjoram, mint, oregano, parsley, rosemary, sage, savory, sweet cicely, tarragon, thyme.

This approval provides for the use of the product **Infinito (PCS No. 04309)** in respect of, the crops and situations as listed above, which are extra to those specified on the product label.

The conditions of application outlined below are statutory. They must be complied with when availing of this extension for minor uses.

The instructions provided in this extension for minor uses must be used in conjunction with information provided on the approved product label.

Failure to abide by the conditions of approval specified below or on the product label may constitute a breach of the product approval.

Reference Date: 09.03.22. [Includes latest changes. This version should be used until

updated]

Product name:	Infinito
Active Substance:	Fluopicolide + Propamocarb hydrochloride
Formulation:	62.5g/l + 625g/l SC
Authorisation Holder:	Bayer CropScience Ltd.
Marketing Company:	Bayer CropScience Ltd.
PCS No:	04309
User type/Function:	Professional Use Only / Fungicide.

Application:

Crop	Max.	Max.	Max.	Minimum	Latest	Target
	individual	number of	total	interval	time of	J
	dose	applications	dose	between	application	
				applications		
Indoor crops (prior	1.6 l/ha	1 per year	1.6 l/ha		BBCH 14	Downy
to transplanting) of:					[before 4	mildew
cabbage, broccoli,					true leaf	
calabrese, Brussel					growth	
sprout, kale, collard					stage]	
& cauliflower.						
Outdoor crops of:	1.6 l/ha	1 per year	1.6 l/ha		14 days	Downy
cabbage, broccoli,					before	mildew
calabrese, Brussel					harvest.	
sprout, kale, collard,						
cauliflower, lettuce,						
spinach, baby leaf						
production and						
herbs.						
Outdoor & indoor	1.6 l/ha	1 per year	1.6 l/ha		7 days	Downy
bulb onion, garlic &					before	mildew
shallot.	4.61/1		1.61/1		harvest.	
Outdoor & indoor	1.6 l/ha	1 per year	1.6 l/ha		14 days	White
leek & salad onion.					before	tip
	4.61/1		1.61/1		harvest.	
Indoor crops of	1.6 l/ha	1 per year	1.6 l/ha		21 days	Downy
lettuce.					before	mildew
					harvest	

Other specific restrictions:

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

- (1) To avoid risks to human health & the environment comply with the terms of this approval in conjunction with the directions for use on the product label.
- (2) For any individual crop the maximum total dose refers to the total dose of any product containing the same active ingredient.
- (3) Application may only be made using a tractor mounted/trailer or handheld sprayer only.
- (4) Post application no unprotected person should enter the treated area until the application had dried & the area has been well ventilated.
- (5) This approval extends to professional use only.
- (6) For environmental information refer to the product label but specifically: To protect groundwater do not apply this or any other product containing fluopicolide at a rate of more than 400 g/ha in any three year period on the same area of land. [For example an application of 1.6 litres of Infinito applies 1.6 L/ha x 62.5g of fluopicolide = 100g fluopicolide/ha]
 A maximum of 1 application per year on the same area of land must not be exceeded.
- (7) Applications must only be made between 1st April and 1st October.
- (8) For operator protection refer to the product label in conjunction with the specific restrictions above and operators must wear suitable protective gloves when handling contaminated surfaces or applying by hand held equipment.
- (9) For: cabbage, broccoli, calabrese, Brussel sprout, kale, collard & cauliflower.
 - Application to outdoor crops must only be made between growth stages BBCH 20 [10 true leaves] and BBCH 49 [typical head size, firmness or leaf mass has been reached]
 - Applications made between BBCH 40 [start of formation bulking of the edible part of the plant] and BBCH 49 must be made in a minimum water volume of 500 litres /ha.
- (10) For outdoor lettuce, spinach, baby leaf production and herbs:
 - Applications must only be made between growth stages BBCH 20 and BBCH 49.
 - Apply in a minimum of 200 litres of water /ha.

- (11) For leek, salad onion, bulb onion, garlic & shallot: Apply in a minimum of 200 litres of water/ha.
- (12) For indoor lettuce, applications must only be made from growth stage BBCH13
- (13) Adverse effects: Approval holders must immediately submit any new information on the potentially dangerous effects of a product or residues of an active substance contained in a product, on human or animal health, or the environment.

Advisory information:

This approval relates to the use of Infinito as a horticultural fungicide for the control white tip on leek & downy mildew in all the other crops & situations listed above.

Infinito has protectant/systemic fungicide properties and should not be used as a curative application.

Growers are advised not to apply this product in situations of crop stress especially drought stress where the crop is not actively growing. In this situation wait until it has rained or irrigation has been applied.

Resistance Management:

Use Infinito as part of an Integrated Crop Management (ICM) strategy. Total reliance on one pesticide will hasten the development of resistance. Pesticides of different chemical groups or alternative control measures should be included in the planned anti-resistance strategy. This product should be used in conjunction with other fungicides from different cross-resistant groups [different mode of action, FRAC group] with satisfactory efficacy against the target.

Infinito contains Fluopicolide FRAC class: 43

Infinito also contains Propamocarb hydrochloride FRAC class: 28 Carbamates

IMPORTANT PLEASE NOTE:

Prior to the use of this product, growers should take advice from their crop advisors particularly in relation to the use of this product within an integrated crop management system of production.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.