

EXTENSION OF AUTHORISATION FOR MINOR USES (EAMU)

INSTRUCTIONS FOR USE

PPP: Starane Hi-Load HL (PCS No. 04549)

CROPS:

(Outdoor grown crops of: Bulb onions, shallot, garlic & leek)

For this EAMU the following crop definitions apply:

Bulb onions includes crops grown from seeds or sets.

This approval provides for the use of the product **Starane Hi-Load HL (PCS No. 04549)** in respect of, the crops and situations as listed above, which are extra to those specified on the product label.

The conditions of application outlined below are statutory. They must be complied with when availing of this extension for minor uses.

The instructions provided in this extension for minor uses must be used in conjunction with information provided on the approved product label.

Failure to abide by the conditions of approval specified below or on the product label may constitute a breach of the product approval.

Reference Date: 26.01.18. [Includes latest changes. This version should be used until updated]

<i>Product name:</i>	Starane Hi-Load HL
<i>Active Substance:</i>	Fluroxypyr
<i>Formulation:</i>	333g/L EC
<i>Authorisation Holder:</i>	Dow AgroSciences Ltd
<i>Marketing Company:</i>	Dow AgroSciences Ltd
<i>PCS No:</i>	04549
<i>User type/Function:</i>	Professional Use Only / Herbicide.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

Application:

Crop	Max. Individual dose	Max. Number of applications	Max. Total dose	Minimum interval between applications	Latest time of application, PHI	Target
Outdoor grown: Onion, garlic & shallot.	0.3 l/ha	n/a	0.3 l/ha	n/a	Before BBCH 15 (5 th leaf clearly visible)	Volunteer potato, cleaver & chickweed
Outdoor grown: Leek	0.18 l/ha	n/a	0.9 l/ha	n/a	28 days before harvest	Volunteer potato, cleaver & chickweed

Other specific restrictions:

- (1) To avoid risks to human health & the environment comply with the terms of this approval in conjunction with the directions for use on the product label.
- (2) For any individual crop the maximum total dose refers to the total dose of any product containing the same active ingredient.
- (3) Post application no unprotected person should enter the treated area until the application had dried & the area has been well ventilated.
- (4) This approval extends to professional use only.
- (5) Application may be made via tractor mounted/trailed sprayer only.
- (6) For environmental information refer to the product label and specifically: To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies from tractor mounted/trailed sprayers. Direct spray away from water. Growers may be eligible to reduce the buffer zone under the PRCD STRIPE Scheme.
- (7) For operator protection refer to the product label in conjunction with the specific restrictions above and specifically: Operators must wear suitable protective coveralls and gloves when handling the product. Operators must wear suitable protective gloves when handling contaminated surfaces.
- (8) Adverse effects: Approval holders must immediately submit any new information on the potentially dangerous effects of a product or residues of an active substance contained in a product, on human or animal health, or the environment.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

Advisory information:

This approval relates to the use of Starane Hi-Load HL as a horticultural herbicide for use on the crops and in the situations listed above. Due to the range of crops on this off label & the numerous different varieties growers are advised to test a small area of the crop first to ensure application will not cause damage to the crop especially if the crop is under any stress.

Apply in a minimum of 200 l/water/ha

Anti-resistance Strategy:

This product should be applied in accordance with an anti-resistance strategy. Weed control may be reduced if resistant strains develop. It is recommended that Starane Hi-Load HL is used in an alternating programme of products with different modes of action (HRAC group) & also with efficacy against the target.

Fluroxypyr is classed HRAC group O.

IMPORTANT PLEASE NOTE:

Prior to the use of this product, growers should take advice from their crop advisors particularly in relation to the use of this product within an integrated crop management system of production.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.