EXTENSION OF AUTHORISATION FOR MINOR USES (EAMU)

INSTRUCTIONS FOR USE

PPP: Teppeki (PCS No. 04399)

CROPS: (Outdoor grown) Head cabbage, broad beans, field beans, French beans, peas-combining, peas-vining, lentils, lupins, Brussels sprout, cherries, carrots, parsnips, celeriac, horseradish, Jerusalem artichoke, parsley root, radish, salsify, turnip and swede. (Protected crops) peppers, raspberries, blackberries

This approval provides for the use of the product <u>Teppeki (PCS No. 04399)</u> in respect of, the crops and situations as listed above, which are extra to those specified on the product label.

The conditions of application outlined below are statutory. They must be complied with when availing of this extension for minor uses.

The instructions provided in this extension for minor uses must be used in conjunction with information provided on the approved product label.

Failure to abide by the conditions of approval specified below or on the product label may constitute a breach of the product approval.

Reference Date: 07.05.25. [Includes latest changes. This version should be used until updated]

Product name: Teppeki

Active Substance: Flonicamid

Formulation: 500 g/kg WG

Authorisation Holder: ISK Biosciences Europe N.V.

Marketing Company: Belchim Crop Protection Ltd

PCS No: 04399

User type/Function: Professional Use Only / Insecticide.

Application:

чррисатон.						
Crop	Max. Individual dose	Max. Number of applications	Max. Total dose	Minimum interval between applications	Latest time of application, PHI	Target
Head cabbage, Brussels sprout	0.14 kg/ha	2 per crop	0.28 kg/ha	14 days	14 days before harvest	Aphid
Broad beans, field beans, French beans, Peas- combining, Peas-vining, lentils, lupins	0.14 kg/ha	1 per crop	0.14 kg/ha	-	GS 71 (10% of pods final length)	Aphid
Cherries (outdoor)	0.14 kg/ha	2 per year	0.28 kg/ha	21 days	14 days before harvest	Aphid
Peppers (protected)	10g/100L water	2 per crop	-		1 day before harvest	Aphid
Raspberry, blackberry (protected)	0.14 kg/ha	2 per year	0.28 kg/ha	7 days	3 days before harvest	Aphid
Carrots, parsnips, celeriac, horseradish, Jerusalem artichoke, parsley root, radish, salsify, turnip and swede	0.14 kg/ha	2 per crop	0.28 kg/ha	14 days	21 days before harvest	Aphid

Other specific restrictions:

- (1) To avoid risks to human health & the environment comply with the terms of this approval in conjunction with the directions for use on the product label.
- (2) For any individual crop the maximum total dose refers to the total dose of any product containing the same active ingredient.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

- (3) Post application no unprotected person should enter the treated area until the application had dried.
- (4) This approval extends to professional use only.
- (5) Application may be made via tractor mounted/trailed boom sprayer only, orchard sprayer.
- (6) DANGEROUS TO BEES. To protect bees and pollinating insects do not apply to crop plants when in flower. Do not use where bees are actively foraging. Do not apply when flowering weeds are present.
- (7) For operator protection refer to the product label.
- (8) Adverse effects: Approval holders must immediately submit any new information on the potentially dangerous effects of a product or residues of an active substance contained in a product, on human or animal health, or the environment.

Advisory information:

This approval relates to use of Teppeki as a horticultural insecticide for the control of a range of aphid species on the crops and situations listed above.

Resistance Management:

Teppeki containing Flonicamid is classed as follows: IRAC group 29

Use as part of an Integrated Resistance Management strategy. Rotate the use between insecticides with different modes of action (IRAC groups) showing good efficacy on the target.

IMPORTANT PLEASE NOTE:

Prior to the use of this product, growers should take advice from their crop advisors particularly in relation to the use of this product within an integrated crop management system of production.