

EXTENSION OF AUTHORISATION FOR MINOR USES (EAMU)

INSTRUCTIONS FOR USE

PPP: Zorvec Entecta (PCS No. 06766)

CROPS:

[Outdoor grown: Bulb onion, garlic and shallot]

For this EAMU the following crop definitions apply:

Onions include Bulb onions grown from seed or sets only. This approval does not include salad onion.

This approval provides for the use of the product **Zorvec Entecta (PCS No. 06766)** in respect of, the crops and situations as listed above, which are extra to those specified on the product label.

The conditions of application outlined below are statutory. They must be complied with when availing of this extension for minor uses.

The instructions provided in this extension for minor uses must be used in conjunction with information provided on the approved product label.

Failure to abide by the conditions of approval specified below or on the product label may constitute a breach of the product approval.

Reference Date: 24.04.25. [Includes latest changes. This version should be used until updated]

<i>Product name:</i>	Zorvec Entecta
<i>Active Substance:</i>	Oxathiapiprolin + Amisulbrom
<i>Formulation:</i>	48 g/L + 240 g/L (SE)
<i>Authorisation Holder:</i>	Corteva Agriscience UK Ltd
<i>Marketing Company:</i>	Corteva Agriscience UK Ltd
<i>PCS No:</i>	06766
<i>User type/Function:</i>	Professional Use Only / Fungicide.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

Application:

Crop	Max. individual dose	Max. number of applications	Max. total dose	Minimum interval between applications	Latest time of application	Target
Outdoor grown bulb onion, garlic & shallot	0.3 L/ha	2 per crop	0.6 L/ha	5 days	28 days before harvest	<i>Peronospora destructor</i>

Other specific restrictions:

- (1) To avoid risks to human health & the environment comply with the terms of this approval in conjunction with the directions for use on the product label.
- (2) For any individual crop the maximum total dose refers to the total dose of any product containing the same active ingredient.
- (3) Application by tractor mounted/trailed horizontal boom sprayer only.
- (4) Post application, do not allow unprotected persons to enter treated areas until the crop is completely dry & has been well ventilated.
- (5) For environmental information refer to the product label but specifically: To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies from tractor mounted/trailed sprayers. Aim spray away from water. Growers may be eligible to reduce the buffer zone under the PRCD STRIPE Scheme.
- (6) This approval extends to professional use only.
- (7) For operator protection refer to the product label in conjunction with the specific restrictions above.
- (8) Adverse effects: Approval holders must immediately submit any new information on the potentially dangerous effects of a product or residues of an active substance contained in a product, on human or animal health, or the environment.

Advisory information:

This approval relates to the use of Zorvec Entecta as a horticultural fungicide for the control of downy mildew (*Peronospora destructor*)

Apply Zorvec Entecta as part of a protectant programme, from 3rd leaf fully visible stage (BBCH 13) until leaves dead, bulb top dry; dormancy Growth complete (BBCH 49). Apply in 200 to 1500 litres water/ha.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

Resistance Management:

Total reliance on one pesticide will hasten the development of resistance. Pesticides of different chemical groups or alternative control measures should be included in the planned anti-resistance strategy. This product should be used in conjunction with other fungicides from different cross-resistant groups [different mode of action, FRAC group] with satisfactory efficacy against the target.

Zorvec Entecta contains oxathiapiprolin and amisulbrom

Oxathiapiprolin is classed FRAC group F9

Amisulbrom is classed within the FRAC group C4

Zorvec Entecta should be used preventatively and should not be relied on for its curative potential.

IMPORTANT PLEASE NOTE:

Prior to the use of this product, growers should take advice from their crop advisors particularly in relation to the use of this product within an integrated crop management system of production.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.