



Department of
**Agriculture,
Food and the Marine**
An Roinn
**Talmhaíochta,
Bia agus Mara**

Pesticide Usage in Ireland

Arable Crops Survey Report 2016

Pesticide Usage in Ireland

ARABLE CROPS SURVEY REPORT 2016

**Pesticide Control Division
DAFM
Backweston Campus
Celbridge
Co. Kildare**

Tel: 01 6157578

Fax: 01 6157574

Email: pcs@agriculture.gov.ie

Contents

| | |
|----------------------------------------------------------|--------------|
| List of tables | 3 |
| Arable survey report summary | 5 |
| Background | 6 |
| Methods | 8 |
| Definitions | 10 |
| Crops | 11 |
| Regional distribution of crops | 12 |
| Pesticide usage | 15 |
| Arable crop areas 2012-2016 | 20 |
| Quantity of pesticide applied per crop, 2012-2016 | 20 |
| Pesticide applied on crop growing area, 2012-2016 | 22 |
| Pesticide usage survey results 2016 | 27 |
| Spring barley | 27 |
| Winter barley | 29 |
| Spring wheat | 31 |
| Winter wheat | 33 |
| Spring oats | 35 |
| Winter oats | 37 |
| Oilseed rape | 39 |
| Peas and Beans | 41 |
| Seed potato crops | 43 |
| Early Potato crops | 45 |
| Maincrop (ware) potato crops | 47 |
| Tables | 49-85 |
| Acknowledgements | 86 |
| References | 86 |

List of tables

| Table | | Page |
|-------|----------------------------------------------------------------------------------------------------------------------------------|------|
| 1 | Estimated area (hectares) of arable crops grown regionally in Ireland, 2016. | 49 |
| 2a | Estimated area (spray-hectares) of arable crops treated regionally with each pesticide type in Ireland, 2016. | 49 |
| 2b | Estimated weight (kg) applied to arable crops regionally with each pesticide type in Ireland, 2016. | 50 |
| 3 | The total area (spray hectares) and the basic area (hectares), of arable crops in Ireland 2016 treated with each pesticide type. | 51 |
| 4 | The total quantities (kilograms) of each pesticide type used on arable crops in Ireland, 2016. | 52 |
| 5 | Estimated area (spray-hectares) of arable crops treated with pesticide formulations in Ireland, 2016. | 53 |
| 6 | Estimated quantities (kilograms) of pesticide formulations used on arable crops in Ireland, 2016. | 60 |
| 7 | The active ingredients most extensively used on arable crops in Ireland in 2016, ranked by area treated (spray-hectares). | 67 |
| 8 | The active ingredients most extensively used on arable crops in Ireland in 2016, ranked by weight (kilograms). | 68 |
| 9 | Estimated quantity (kg), spray area (spha) and basic area (ha) of active substance for Spring barley 2016. | 69 |
| 10 | Estimated quantity (kg), spray area (spha) and basic area (ha) of active substance for Winter barley 2016. | 71 |

List of tables (contd.)

| Table | | Page |
|-------|----------------------------------------------------------------------------------------------------------------|------|
| 11 | Estimated quantity (kg), spray area (spha) and basic area (ha) of active substance for spring wheat 2016. | 73 |
| 12 | Estimated quantity (kg), spray area (spha) and basic area (ha) of active substance for winter wheat 2016. | 75 |
| 13 | Estimated quantity (kg), spray area (spha) and basic area (ha) of active substance for spring oats 2016. | 77 |
| 14 | Estimated quantity (kg), spray area (spha) and basic area (ha) of active substance for winter oats 2016. | 79 |
| 15 | Estimated quantity (kg), spray area (spha) and basic area (ha) of active substance for oilseed rape 2016. | 81 |
| 16 | Estimated quantity (kg), spray area (spha) and basic area (ha) of active substance for peas and beans 2016. | 82 |
| 17 | Estimated quantity (kg), spray area (spha) and basic area (ha) of active substance for seed potatoes 2016. | 83 |
| 18 | Estimated quantity (kg), spray area (spha) and basic area (ha) of active substance for early potatoes 2016. | 84 |
| 19 | Estimated quantity (kg), spray area (spha) and basic area (ha) of active substance for maincrop potatoes 2016. | 85 |

Arable survey report summary

This is the third survey of pesticide usage on arable crops in Ireland carried out by the Department of Agriculture, Food and the Marine (DAFM), providing comparative data to that obtained in the previous survey in 2012.

Information on all aspects of pesticide usage was collected from 260 holdings across Ireland representing 7.58% of the total area of arable crops grown. Quantitative data have been adjusted to provide estimates of total pesticide usage.

**Pesticide is an over-arching term that includes both plant protection products (including, for the purpose of this report, fungicides, herbicides, insecticides, molluscicides, biological controls and seed treatments) and biocides.*

In 2016 an estimated 306,092 hectares of arable crops were grown which when adjusted for crops not surveyed (triticale) represents a 10% decrease compared to total estimated area in 2012. An estimated 1,058,461 kgs of active substance was applied to arable crops in 2016 which represents a 7.2% decrease in overall weight of pesticides applied compared to the equivalent figure in 2012. The total pesticide treated area of the crops surveyed declined by 11.2% from 2012 to 2016.

A total of 120 active substances were recorded in use on arable crops in the survey compared to 114 in 2012.

Winter barley comprised 25% of the area of arable crops, accounting for 27% of the total pesticide treated area and 29% of the total weight of pesticides used on all arable crops in 2016.

Spring barley comprised 37% of the area of arable crops and accounted for 30% of the total pesticide treated area and 22% of the total weight of pesticides used on all arable crops in 2016.

Winter wheat comprised 17% of the area of arable crops and accounted for 23% of the total pesticide treated area and 25% of the total weight of pesticides used on all arable crops in 2016.

Spring wheat comprised of 4% of the area of arable crops, accounting for 4% of the total pesticide treated area and 4% of the total weight of pesticides used on all arable crops in 2016.

Winter oats comprised of 4% of the total area of arable crops, accounting for 4% of the total pesticide treated area and 4% of the total weight of pesticides applied on arable crops in 2016.

Spring oats comprised of 3% of the total area of arable crops and accounted for 3% of the total pesticide treated area and 3% of total weight of pesticides used on all arable crops in 2016.

Oilseed rape (spring & winter) comprised of 3% of the total area of arable crops and accounted for 2% of the total pesticide treated area and 2% of the total weight of pesticides applied.

Beans and peas comprised 3% of the total area of arable crops, accounting for 2% of the total pesticide treated area and 3% of the total weight of pesticides applied.

Potato crops compromised 3% of the area of arable crops grown in Ireland in 2016, accounting for 5% of the total pesticide-treated area and 9% of the total weight of pesticides used on all arable crops. The total area of potatoes grown compromised 88% main crop, 4% seed and 8% early potato crops.

Background

The regulatory system for PPPs in Ireland is based directly on EU legislation which provides a very high level of protection for man, animals and the environment. The hazard of an active substance is an inherent property which can cause a harmful effect and cannot be altered or mitigated.

Legislation has been put in place at both EU and national level to minimise the risks associated with the use of PPPs while ensuring necessary crop protection. The Sustainable Use of Pesticides Directive based on the EU ‘Thematic strategy on the sustainable use of pesticides’ aims to achieve a balance between ensuring human and environmental safety while maintaining continued viability of the farming and amenity sectors. This involves training and registration of advisers, distributors, operators and inspectors of pesticide application equipment, controls on storage, supply and use, adoption of the principles of IPM and improved statistics on PPP use.

To address the requirement for improved statistics, Regulation (EC) No 1185/2009 was adopted on 25 November 2009 which requires each member state to collect statistics on PPP use. It is the area identified above as “improved statistics on PPP use” that this survey and future surveys will be addressing.

While sales data can provide information on the overall quantity of PPPs used in the country, surveys at farm/grower/producer level are required to quantify the amounts used on different crops and to identify where and how they are being used. This type of information is required to clearly identify the risks involved and to develop and defend a strategy for the sustainable use of PPPs. Some of the specific outputs of a usage survey are as follows:

1. Provision of reliable factual data to inform policy makers.
2. Provision of information for the on-going review process of existing PPPs by providing data regarding national and regional usage of PPPs and use patterns for particular crops.
3. Monitoring farm practices to highlight areas where PPP use might be reduced by supplementation with or replacement by alternative pest control strategies e.g. use of resistant varieties, cultivation practices etc.
4. Provision of data to assess likely operator exposure to PPPs and to predict environmental impact of PPP use.
5. Monitoring changes in patterns of PPP use over time in response to government policy or economic factors.
6. Provision of information for residue monitoring programmes to assist with identifying particular areas of risk and to validate findings.

Methods

The sample of holdings to be surveyed was selected from each of the 26 counties, on the basis of the total area of arable crops grown, using data from DAFM. For the purpose of the survey the country was divided into three geographical regions namely the East, South and the North/West as per Table A. The sample was stratified into six size groups, according to the total area of arable crops grown in each region. Holdings were selected at random within each of the size groups and the number of holdings selected was proportional to the total area of arable crops grown.

Table A: Regions selected for survey and respective counties.

| Regions | East | South | North/West |
|----------|---------|-----------|------------|
| Counties | Louth | Wexford | Donegal |
| | Meath | Kilkenny | Leitrim |
| | Dublin | Waterford | Monaghan |
| | Kildare | Tipperary | Cavan |
| | Offaly | Limerick | Westmeath |
| | Laois | Cork | Longford |
| | Carlow | Kerry | Sligo |
| | Wicklow | | Roscommon |
| | | | Mayo |
| | | | Galway |
| | | | Clare |

The purpose of the survey was explained to the occupiers of selected holdings in preliminary correspondence. A total of 260 holdings were visited during the period May to July 2017 and data collected by personal interview for arable crops harvested in 2016. The data collected included; the area of crops grown, area treated, target crop, pesticide used and number of treatments applied. Holdings selected in the original sample which were unable to provide data were replaced with ones from the same county and size group held on a reserve list. The total number of farms in each size group and the number of farms sampled are shown in Table B. The collected data were entered using Oracle, a relational database programme. Validated data were downloaded for analysis using SPSS software.

Table B: The total number of arable farms in each size group (A) and the number of samples from each size group (B).

| Region | Size group (hectares) | | | | | | | | | | | | | | Total |
|----------------|-----------------------|------------------|---------------|------------------|---------------|------------------|---------------|------------------|---------------|------------------|---------------|------------------|---------------|------------------|-------|
| | <10 | | 10<20 | | 20<40 | | 40<100 | | 100-200 | | >200 | | Holdings | Holdings | |
| | in size group | Holdings sampled | in size group | Holdings sampled | in size group | Holdings sampled | in size group | Holdings sampled | in size group | Holdings sampled | in size group | Holdings sampled | in size group | Holdings sampled | |
| A | B | A | B | A | B | A | B | A | B | A | B | A | B | A | B |
| East | 1,457 | 7 | 857 | 8 | 897 | 19 | 803 | 40 | 229 | 27 | 86 | 19 | 4,329 | 120 | |
| South | 2,343 | 8 | 1,287 | 15 | 1,073 | 27 | 712 | 34 | 140 | 14 | 57 | 14 | 5,612 | 112 | |
| North/West | 795 | 7 | 203 | 3 | 118 | 7 | 102 | 7 | 10 | 4 | 2 | 0 | 1,230 | 28 | |
| <i>Ireland</i> | <i>4,595</i> | <i>22</i> | <i>2,347</i> | <i>26</i> | <i>2,088</i> | <i>53</i> | <i>1,617</i> | <i>81</i> | <i>379</i> | <i>45</i> | <i>145</i> | <i>33</i> | <i>11,171</i> | <i>260</i> | |

Definitions

- ‘Basic area’; refers to the actual planted area of crop treated with a given pesticide.
- ‘Biocides’; are defined as chemicals that are used to control and / or prevent various types of harmful or unwanted organisms, including disinfectants, preservatives, insect repellents, rodenticides and insecticides.
- ‘Fungicides’; are defined as PPPs used to control and / or prevent harmful fungal disease.
- ‘Growth regulators’; are defined as PPPs used to control/ regulate the growth of the plant.
- ‘Herbicides’; are defined as PPPs used to control and / or prevent unwanted vegetation.
- ‘Insecticides’; are defined as PPPs used to control and / or prevent harmful insects.
- ‘Molluscicides’; are defined as PPPs used to control and / or prevent harmful slugs and snails.
- ‘PPP’; refers to plant protection product.
- ‘Rounding’; due to rounding of figures there may be slight differences in totals both within and between tables.
- ‘Seed treatments’; are defined as PPPs applied to seeds to provide protection and improve the establishment of healthy crops.

- ‘Spray applications’; refers to the number of treatments of any pesticide type to the treated areas.
- ‘Treated area’; refers to the total area treated with a pesticide, which includes all repeated applications to the basic area. This is measured in ‘spray-hectares’ (basic area x number of spray applications = spray hectares (spha)).

Crops


Information was collected for spring barley, winter barley, spring wheat, winter wheat, spring oats, winter oats, oilseed rape, peas & beans, seed potatoes, early potatoes and maincrop potatoes.

The number and areas of crops surveyed are shown in Table C. Data from 260 farms provided information on 860 examples of 11 crop types. The total area of crops sampled in the survey (23,199 ha) was representative of the area of arable crops grown in Ireland in 2016 (306,092 ha).

Table C: The total number and area (hectares) of crops sampled, and the proportion (%) of the total area of crops surveyed in Ireland, 2016.

| Crop | Number of crops surveyed | Survey area (ha) | Proportion of crop area surveyed (%) |
|----------------------|--------------------------|------------------|--------------------------------------|
| Spring barley | 206 | 5,668 | 5.06% |
| Winter barley | 182 | 5,803 | 7.57% |
| Spring wheat | 41 | 527 | 4.74% |
| Winter wheat | 139 | 5,661 | 10.86% |
| Spring oats | 49 | 702 | 6.29% |
| Winter oats | 56 | 940 | 7.15% |
| Oilseed rape (S & W) | 54 | 1,207 | 12.17% |
| Peas & beans | 67 | 935 | 8.49% |
| Seed potatoes | 9 | 143 | 46.56% |
| Early potatoes | 19 | 178 | 23.54% |
| Maincrop potatoes | 38 | 1,436 | 17.98% |
| Total | 860 | 23,199 | 7.58% |

Figure 1: Areas of individual arable crops grown in Ireland (ha), 2016.



Regional distribution of crops

The East region was the largest producer of arable crops during 2016 and accounted for 50% of the total area of arable crops grown and 52% of the total pesticide-treated area. Overall, 50% of the weight of herbicides, 51% of the weight of fungicides, 73% of the weight of insecticides, 52% of the weight of molluscicides, 55% of the weight of growth regulators and 49% of the weight of seed treatments were applied to arable crops in this region.

The South region accounted for 45% of the area of arable crops grown and 44% of the total pesticide-treated area. Overall, 45% of the weight of herbicides, 44% of the weight of fungicides, 25% of the weight of insecticides, 39% of the weight of molluscicides, 43% of the weight of growth regulators and 47% of the weight of seed treatments were applied to arable crops in this region.

The North/West region accounted for 5% of the total arable growing area and 4% of the pesticide treated area. Overall, 5% of the weight of herbicides, 5% of the weight of fungicides, 2% of the weight of insecticides, 9% of the weight of molluscicides, 2% of the weight of growth regulators and 4% of the weight of seed treatments were applied to arable crops in this region.

Figure 2: Regional distribution (ha) of arable crops grown in Ireland, 2016.




Figure 3: Regional distribution (ha) of individual arable crops grown in Ireland, 2016.




Figure 4: Regional distribution (ha) of cereal crops grown in Ireland, 2016.




Figure 5: Regional distribution (ha) of potato crops (seed, early & maincrop) grown in Ireland, 2016.





Figure 6: Regional distribution (ha) of other arable crops (oilseed rape, peas & beans) grown in Ireland, 2016.



Pesticide usage

Fungicides were applied to 44% of the pesticide-treated area, accounting for 46% of the total weight of pesticides used. Herbicides were applied to 28% of the pesticide-treated area, representing 31% of the total weight of pesticides used. Insecticides were applied to 9% of the pesticide treated area of arable crops, representing 2% of the weight of pesticides applied. Molluscicide treatments represented less than 1% of pesticide treated area and less than 1% of the weight of pesticides applied. Growth regulator usage accounted for 11% of the pesticide-treated area and 18% of the weight of active substance applied. Seed treatments were applied to 8% of the pesticide-treated area, representing 2% of the weight of active substances applied.

The use of fungicides on cereal crops accounted for 42% of the cereal pesticide treated area and 44% of the total weight of pesticides applied to cereal crops. The use of fungicides on potato crops accounted for 68% of the potato pesticide treated area and 70% of the total weight of pesticides applied to potato crops. The use of herbicides on other arable crops (Peas, beans and Oil seed rape) accounted for 38% of the treated area and 68% of the total weight of pesticides applied to this crop group.

Figure 7: Pesticide usage (spha) on arable crops treated in Ireland, 2016.




Figure 8: Weight (kgs) of pesticides applied to arable crops treated in Ireland, 2016.




Figure 9: Pesticide usage (spha) on cereal crops grown in Ireland, 2016.




Figure 10: Weight of pesticides (kg) applied to cereal crops grown in Ireland, 2016.




Figure 11: Pesticide usage (spha) on potato crops (seed, early & maincrop) grown in Ireland, 2016.




Figure 12: Weight of pesticides (kg) applied to potato crops (seed, early & maincrop) grown in Ireland, 2016.




Figure 13: Pesticide usage (spha) on other arable crops (oilseed rape, peas & beans) grown in Ireland, 2016.





Figure 14: Weight of pesticides (kg) applied to other arable crops (oilseed rape, peas & beans) grown in Ireland, 2016.



Arable crop areas 2012-2016

Winter barley increased in area by 88% to 76,626 ha when comparing 2016 and 2012. The area of Spring barley decreased by 26%. Winter wheat witnessed a 38% reduction in area from 83,751 ha to 52,134 ha. There was a 34% increase in the area of winter oats in the period 2012 to 2016. Details of changes in crop areas between 2012 and 2016 are outlined in Table D below.


Table D: Arable crops areas (ha) for surveys in 2012 and 2016 and percentage (%) change in areas grown.

| Crop | Ha grown | | % change |
|-----------------------------|----------|----------------|----------|
| | 2012 | 2016 | |
| Spring barley | 150,378 | 111,929 | -26 |
| Winter barley | 40,717 | 76,626 | 88 |
| Spring wheat | 13,314 | 11,125 | -16 |
| Winter wheat | 83,751 | 52,134 | -38 |
| Spring oats | 13,998 | 11,151 | -20 |
| Winter oats | 9,782 | 13,146 | 34 |
| Oilseed rape | 17,282 | 9,912 | -43 |
| Peas & beans | 3,696 | 11,016 | 198 |
| Triticale | 1,445 | (not surveyed) | |
| All Potatoes | 9,035 | 9,052 | 0 |
| All crops (excl. triticale) | 341,953 | 306,092 | -10 |

Quantity of pesticide applied per crop, 2012-2016


The average weight of pesticide applied per hectare of crop grown for each crop is provided in Figure 15. Average weights were calculated as the total weight of pesticides applied divided by the total area of crop grown (whether treated or untreated).

Figure 15: Average weight of pesticides applied per hectare of crop grown in Ireland (kg/ha), 2016.



The highest weight of pesticides applied per hectare was on potato crops (10.4 kg/ha) which results from the relatively high number of pesticide treatments, in particular fungicide treatments, applied to potato crops. The winter cereal crops had the next highest levels of pesticide use with winter wheat (5.0 kg/ha), winter barley (4.0 kg/ha) and winter oats (3.4 kg/ha). The average weight of pesticide applied per hectare of crop grown as per 2012 survey is provided in Figure 16.

Figure 16: Average weight of pesticides applied per hectare of crop grown in Ireland (kg/ha), 2012.



The quantity of pesticides applied to potato crops increased from 10.3 kg/ha to 10.4 kg/ha during the period 2012 to 2016. The quantity of pesticides applied to peas and beans has decreased from 3.4 kg/ha to 2.5 kgs/ha during this period. There has been a decrease in pesticides applied to winter wheat from 5.2 kgs/ha to 5.0 kgs/ha. The level of pesticides applied to spring wheat increased from 2.7 kgs/ha to 4.0kgs ha.

Pesticide applied on crop growing area, 2012-2016.

The average weight of pesticide applied per hectare of crop grown for 2012 and 2016 including percentage change is provided.

Spring barley

Quantities (kg/ha) of fungicides, herbicides and molluscicides applied to spring barley decreased by 3%, 8% and 100% respectively when comparing 2012 and 2016. Quantities of insecticides, growth regulators and seed treatments increased by 74%, 84% and 52% respectively. Overall, there was only a 2% increase in quantities of pesticides (kg/ha) applied to spring barley when comparing 2012 and 2016.

Figure 17: Quantity of pesticide type (kg/ha) and percentage change (%) for spring barley for surveys in 2012 and 2016.

| Pesticide type | Kg/ha on area grown | | |
|-------------------|---------------------|--------|----------|
| | 2012 | 2016 | % change |
| Fungicides | 1.14 | 1.11 | -3 |
| Herbicides | 0.67 | 0.61 | -8 |
| Insecticides | 0.0203 | 0.0354 | 74 |
| Molluscicides | 0.0002 | 0.0000 | -100 |
| Growth Regulators | 0.11 | 0.21 | 84 |
| Seed treatments | 0.05 | 0.07 | 52 |
| All pesticides | 1.99 | 2.04 | 2 |

Winter barley

Quantities (kg/ha) of fungicides & molluscicides reduced by 6% & 65% respectively when comparing 2012 and 2016. Quantities of herbicides, insecticides, growth regulators and seed treatments increased by 12%, 92%, 19% and 51% respectively. Overall, there was a 7% increase in quantities of pesticides applied to winter barley when comparing 2012 and 2016.

Figure 18: Quantity of pesticide type (kg/ha) and percentage change (%) for winter barley for surveys in 2012 and 2016.

| Pesticide type | Kg/ha on area grown | | |
|-----------------------|----------------------------|-------------|-----------------|
| | 2012 | 2016 | % change |
| Fungicides | 1.57 | 1.48 | -6 |
| Herbicides | 1.35 | 1.52 | 12 |
| Insecticides | 0.01 | 0.03 | 92 |
| Molluscicides | 0.0058 | 0.0020 | -65 |
| Growth Regulators | 0.75 | 0.89 | 19 |
| Seed treatments | 0.05 | 0.08 | 51 |
| All pesticides | 3.74 | 3.99 | 7 |

Spring wheat

Overall, there was a 46% increase in quantities of pesticides applied to Spring Wheat when comparing 2012 and 2016.

Figure 19: Quantity of pesticide type (kg/ha) and percentage change (%) for spring wheat for surveys in 2012 and 2016.

| Pesticide type | Kg/ha on area grown | | |
|-----------------------|----------------------------|-------------|-----------------|
| | 2012 | 2016 | % change |
| Fungicides | 1.23 | 1.70 | 38 |
| Herbicides | 0.52 | 0.75 | 46 |
| Insecticides | 0.11 | 0.20 | 81 |
| Molluscicides | 0.00 | 0.00 | 0 |
| Growth Regulators | 0.79 | 1.22 | 55 |
| Seed treatments | 0.06 | 0.09 | 46 |
| All pesticides | 2.71 | 3.96 | 46 |

Winter wheat

Quantities (kg/ha) of fungicides, herbicides & growth regulators applied to winter wheat reduced by 3%, 7% and 7% respectively when comparing 2012 and 2016. Overall, there was a 4% reduction in quantities of pesticides applied when comparing 2012 and 2016.

Figure 20: Quantity of pesticide type (kg/ha) and percentage change (%) for winter wheat for surveys in 2012 and 2016.

| Pesticide type | Kg/ha on area grown | | |
|-----------------------|----------------------------|-------------|-----------------|
| | 2012 | 2016 | % change |
| Fungicides | 2.37 | 2.30 | -3 |
| Herbicides | 1.30 | 1.21 | -7 |
| Insecticides | 0.19 | 0.21 | 9 |
| Molluscicides | 0.01 | 0.02 | 21 |
| Growth Regulators | 1.26 | 1.18 | -7 |
| Seed treatments | 0.07 | 0.08 | 9 |
| All pesticides | 5.21 | 5.00 | -4 |

Spring oats

There was no significant change in overall quantities of pesticides applied to spring oats when comparing 2012 and 2016.

Figure 21: Quantity of pesticide type (kg/ha) and percentage change (%) for spring oats for surveys in 2012 and 2016.

| Pesticide type | Kg/ha on area grown | | |
|-----------------------|----------------------------|-------------|-----------------|
| | 2012 | 2016 | % change |
| Fungicides | 0.98 | 0.98 | 0 |
| Herbicides | 0.6099 | 0.6063 | -1 |
| Insecticides | 0.0242 | 0.0237 | -2 |
| Molluscicides | 0.00 | 0.00 | 0 |
| Growth Regulators | 0.81 | 0.82 | 1 |
| Seed treatments | 0.06 | 0.05 | -6 |
| All pesticides | 2.48 | 2.48 | 0 |

Winter oats

Quantities (kg/ha) of fungicides, herbicides and insecticides applied to winter oats reduced by 8%, 11% and 70% respectively during the period 2012 to 2016. No molluscicide applications were recorded on winter oats. Overall, there was no significant change in total pesticides applied to winter oats when comparing 2012 and 2016.

Figure 22: Quantity of pesticide type (kg/ha) and percentage change (%) for winter oats for surveys in 2012 and 2016.

| Pesticide type | Kg/ha on area grown | | % change |
|-------------------|---------------------|------|----------|
| | 2012 | 2016 | |
| Fungicides | 1.42 | 1.30 | -8 |
| Herbicides | 0.62 | 0.55 | -11 |
| Insecticides | 0.02 | 0.01 | -70 |
| Molluscicides | 0.00 | 0.00 | 0 |
| Growth Regulators | 1.28 | 1.44 | 12 |
| Seed treatments | 0.05 | 0.08 | 47 |
| All pesticides | 3.39 | 3.39 | 0 |

Oilseed rape

Quantities (kg/ha) of molluscicides applied to oilseed rape reduced by 40% when comparing 2012 and 2016. Overall, there was a 10% increase in total quantities of pesticides applied to oilseed rape during the period 2012 to 2016. No growth regulator applications were recorded on oilseed rape.

Figure 23: Quantity of pesticide type (kg/ha) and percentage change (%) for oilseed rape for surveys in 2012 and 2016.

| Pesticide type | Kg/ha on area grown | | % change |
|-------------------|---------------------|--------|----------|
| | 2012 | 2016 | |
| Fungicides | 0.37 | 0.40 | 8 |
| Herbicides | 1.99 | 2.20 | 10 |
| Insecticides | 0.0035 | 0.0145 | 311 |
| Molluscicides | 0.05 | 0.03 | -40 |
| Growth Regulators | 0.00 | 0.00 | 0 |
| Seed treatments | 0.01 | 0.02 | 17 |
| All pesticides | 2.42 | 2.65 | 10 |

Peas and beans

Quantities (kg/ha) of fungicides and herbicides applied to peas and beans reduced by 5% and 41% respectively when comparing 2012 and 2016. Overall, there was a 28% reduction in total pesticides applied to peas and beans during the period 2012 to 2016. No growth regulator or molluscicide applications were noted on peas and beans.

Figure 24: Quantity of pesticide type (kg/ha) and percentage change (%) for peas and beans for surveys in 2012 and 2016.

| Pesticide type | Kg/ha on area grown | | % change |
|------------------------------|----------------------------|-------------|-----------------|
| | 2012 | 2016 | |
| Fungicides | 1.19 | 1.14 | -5 |
| Herbicides | 2.21 | 1.30 | -41 |
| Insecticides | 0.0062 | 0.0128 | 108 |
| Molluscicides | 0.00 | 0.00 | 0 |
| Growth Regulators | 0.00 | 0.00 | 0 |
| Seed treatments | (not recorded) | 0.00 | n/a |
| All pesticides (Ex. sd. tr.) | 3.41 | 2.46 | -28 |

Potatoes (seed, early and maincrop)

Quantities (kg/ha) of fungicides, molluscicides, growth regulators and seed treatments applied to potatoes (as a group) reduced by 6%, 44%, 100% and 29% respectively when comparing 2012 and 2016. Overall, there was only a 1% increase in total pesticides when comparing 2012 and 2016.

Figure 25: Quantity of pesticide type (kg/ha) and percentage change (%) for potatoes for surveys in 2012 and 2016.

| Pesticide type | Kg/ha on area grown | | % change |
|-----------------------|----------------------------|-------------|-----------------|
| | 2012 | 2016 | |
| Fungicides | 7.71 | 7.25 | -6 |
| Herbicides | 2.20 | 2.90 | 32 |
| Insecticides | 0.0188 | 0.0767 | 308 |
| Molluscicides | 0.11 | 0.06 | -44 |
| Growth Regulators | 0.09 | 0.00 | -100 |
| Seed treatments | 0.18 | 0.13 | -29 |
| All pesticides | 10.31 | 10.42 | 1 |

Pesticide usage survey results 2016

Pesticide usage on spring barley

111,929 ha of spring barley grown in Ireland.

978,966 treated hectares.

227,973 kilogrammes applied.

Figure 26: Pesticide usage (spha) on spring barley crops in Ireland, 2016.




Figure 27: Weight of pesticides (kg) applied to spring barley crops in Ireland, 2016.




Figure 28: The top 10 active ingredients most extensively used on spring barley in Ireland in 2016, ranked by area treated (spray-hectares).

| Active substance | Treated area (spha) | Basic area treated (ha) | Quantity applied (kg) | % of the treated area |
|-----------------------|---------------------|-------------------------|-----------------------|-----------------------|
| Prothioconazole | 133,547 | 91,513 | 16,129 | 9.5 |
| Chlorothalonil | 116,017 | 95,771 | 55,179 | 8.2 |
| Fluroxypyr | 83,068 | 81,194 | 11,288 | 5.9 |
| Thifensulfuron-methyl | 75,657 | 73,859 | 2,066 | 5.4 |
| Tribenuron-methyl | 73,950 | 73,479 | 738 | 5.3 |
| Epoxiconazole | 67,661 | 50,186 | 3,946 | 4.8 |
| Pinoxaden | 62,578 | 60,799 | 1,797 | 4.4 |
| Fenpropimorph | 62,018 | 53,784 | 17,101 | 4.4 |
| Metsulfuron-methyl | 59,159 | 56,164 | 245 | 4.2 |
| Lambda-cyhalothrin | 56,757 | 55,142 | 274 | 4.0 |

Figure 29: The top 10 active ingredients most extensively used on spring barley in Ireland in 2016, ranked by weight (kg).

| Active substance | Quantity applied (kg) | Treated area (spha) | Basic area treated (ha) | % of the weight applied |
|----------------------|-----------------------|---------------------|-------------------------|-------------------------|
| Chlorothalonil | 55,179 | 116,017 | 95,771 | 24.2 |
| Glyphosate | 32,230 | 36,952 | 32,924 | 14.1 |
| Chlormequat chloride | 21,575 | 29,810 | 29,458.6 | 9.5 |
| Fenpropimorph | 17,101 | 62,018 | 53,784 | 7.5 |
| Prothioconazole | 16,129 | 133,547 | 91,513 | 7.1 |
| Fluroxypyr | 11,288 | 83,068 | 81,194 | 5.0 |
| Mecoprop-P | 7,192 | 15,989 | 15,168 | 3.2 |
| Carboxin/thiram | 5,360 | 28,313 | 28,313 | 2.4 |
| Pyraclostrobin | 4,945 | 50,629 | 45,982 | 2.2 |
| Cyprodinil | 4,847 | 16,400 | 16,400 | 2.1 |

Pesticide usage on winter barley

76,626 ha of winter barley grown in Ireland.

887,186 treated hectares.

305,992 kilogrammes applied.

Figure 30: Pesticide usage (spha) on winter barley crops in Ireland, 2016.




Figure 31: Weight of pesticides (kg) applied to winter barley crops in Ireland, 2016.




Figure 32: The top 10 active ingredients most extensively used on winter barley in Ireland in 2016, ranked by area treated (spray-hectares).

| Active substance | Treated area (spha) | Basic area treated | Quantity applied (kg) | % of the treated area |
|----------------------|---------------------|--------------------|-----------------------|-----------------------|
| Chlorothalonil | 110,466 | 66,954 | 53,779 | 9.5 |
| Prothioconazole | 107,877 | 64,914 | 13,684 | 9.3 |
| Isoproturon | 63,466 | 61,593 | 69,585 | 5.5 |
| Chlormequat chloride | 63,155 | 57,798 | 51,643 | 5.4 |
| Diflufenican | 56,286 | 53,920 | 6,606 | 4.8 |
| Pyraclostrobin | 55,843 | 40,365 | 6,097 | 4.8 |
| Epoxiconazole | 54,228 | 39,074 | 3,473 | 4.7 |
| Lambda-cyhalothrin | 43,796 | 40,698 | 207 | 3.8 |
| Pinoxaden | 41,473 | 40,910 | 1,124 | 3.6 |
| Trinexapac-ethyl | 41,070 | 35,522 | 2,405 | 3.5 |

Figure 33: The top 10 active ingredients most extensively used on winter barley in Ireland in 2016, ranked by weight (kg).

| Active substance | Quantity applied (kg) | Treated area (spha) | Basic area treated | % of the weight applied |
|----------------------|-----------------------|---------------------|--------------------|-------------------------|
| Isoproturon | 69,585 | 63,466 | 61,593 | 22.7 |
| Chlorothalonil | 53,779 | 110,466 | 66,954 | 17.6 |
| Chlormequat chloride | 51,643 | 63,155 | 57,798 | 16.9 |
| Glyphosate | 29,266 | 34,212 | 27,063 | 9.6 |
| Prothioconazole | 13,684 | 107,877 | 64,914 | 4.5 |
| Fenpropimorph | 8,661 | 31,892 | 25,381 | 2.8 |
| Mepiquat chloride | 7,967 | 23,367 | 22,800 | 2.6 |
| Spiroxamine | 6,941 | 27,468 | 19,983 | 2.3 |
| Diflufenican | 6,606 | 56,286 | 53,920 | 2.2 |
| Ethephon | 6,299 | 29,787 | 29,219 | 2.1 |

Pesticide usage on spring wheat

11,125 ha of spring wheat grown in Ireland.

126,449 treated hectares.

44,050 kilogrammes applied.

Figure 34: Pesticide usage (spha) on spring wheat crops in Ireland, 2016.




Figure 35: Weight of pesticides (kg) applied to spring wheat crops in Ireland, 2016.




Figure 36: The top 10 active ingredients most extensively used on spring wheat in Ireland in 2016, ranked by area treated (spray-hectares).

| Active substance | Treated area (spha) | Basic area treated (ha) | Quantity applied (kg) | % of the treated area |
|----------------------|---------------------|-------------------------|-----------------------|-----------------------|
| Chlorothalonil | 22,010 | 10,396 | 10,927 | 13.6 |
| Chlormequat chloride | 15,510 | 8,972 | 13,212 | 9.6 |
| Epoxiconazole | 10,442 | 8,335 | 867 | 6.4 |
| Fluroxypyr | 8,868 | 8,661 | 1,252 | 5.5 |
| Prothioconazole | 7,315 | 6,549 | 1,183 | 4.5 |
| Metsulfuron-methyl | 6,385 | 6,385 | 28 | 3.9 |
| Glyphosate | 6,282 | 6,143 | 5,296 | 3.9 |
| Tribenuron-methyl | 5,626 | 5,412 | 49 | 3.5 |
| Tebuconazole | 5,195 | 5,195 | 618 | 3.2 |
| Lambda-cyhalothrin | 4,814 | 4,264 | 33 | 3.0 |

Figure 37: The top 10 active ingredients most extensively used on spring wheat in Ireland in 2016, ranked by weight (kg).

| Active substance | Quantity applied (kg) | Treated area (spha) | Basic area treated (ha) | % of the weight applied |
|----------------------|-----------------------|---------------------|-------------------------|-------------------------|
| Chlormequat chloride | 13,212 | 15,510 | 8,972 | 30.0 |
| Chlorothalonil | 10,927 | 22,010 | 10,396 | 24.8 |
| Glyphosate | 5,296 | 6,282 | 6,143 | 12.0 |
| Dimethoate | 1,358 | 4,508 | 4,508 | 3.1 |
| Fluroxypyr | 1,252 | 8,868 | 8,661 | 2.8 |
| Prothioconazole | 1,183 | 7,315 | 6,549 | 2.7 |
| Fenpropimorph | 1,117 | 3,375 | 2,377 | 2.5 |
| Boscalid | 959 | 4,093 | 4,093 | 2.2 |
| Carboxin/thiram | 885 | 4,708 | 4,708 | 2.0 |
| Fenpropidin | 880 | 1,543 | 1,543 | 2.0 |

Pesticide usage on winter wheat

52,134 ha of winter wheat grown in Ireland.

746,636 treated hectares.

260,613 kilogrammes applied.

Figure 38: Pesticide usage (spha) on winter wheat crops in Ireland, 2016.




Figure 39: Weight of pesticides (kg) applied to winter wheat crops in Ireland, 2016.




Figure 40: The top 10 active ingredients most extensively used on winter wheat in Ireland in 2016, ranked by area treated (spray-hectares).

| Active substance | Treated area (spha) | Basic area treated (ha) | Quantity applied (kg) | % of the treated area |
|----------------------|---------------------|-------------------------|-----------------------|-----------------------|
| Chlorothalonil | 138,714 | 50,549 | 70,189 | 13.9 |
| Chlormequat chloride | 68,018 | 44,843 | 57,826 | 6.8 |
| Prothioconazole | 66,201 | 45,072 | 8,516 | 6.6 |
| Epoxiconazole | 63,108 | 37,974 | 5,227 | 6.3 |
| Fluxapyroxad | 53,782 | 38,390 | 5,100 | 5.4 |
| Tebuconazole | 48,860 | 37,254 | 6,422 | 4.9 |
| Metconazole | 45,108 | 36,345 | 2,647 | 4.5 |
| Diflufenican | 40,057 | 33,543 | 4,512 | 4.0 |
| Trinexapac-ethyl | 33,162 | 26,620 | 2,349 | 3.3 |
| Isoproturon | 31,600 | 28,684 | 31,521 | 3.2 |

Figure 41: The top 10 active ingredients most extensively used on winter wheat in Ireland in 2016, ranked by weight (kg).

| Active substance | Quantity applied (kg) | Treated area (spha) | Basic area treated (ha) | % of the weight applied |
|----------------------|-----------------------|---------------------|-------------------------|-------------------------|
| Chlorothalonil | 70,189 | 138,714 | 50,549 | 27 |
| Chlormequat chloride | 57,826 | 68,018 | 44,843 | 22 |
| Isoproturon | 31,521 | 31,600 | 28,684 | 12 |
| Glyphosate | 18,347 | 23,672 | 20,291 | 7 |
| Dimethoate | 9,079 | 30,791 | 30,171 | 3 |
| Prothioconazole | 8,516 | 66,201 | 45,072 | 3 |
| Tebuconazole | 6,422 | 48,860 | 37,254 | 2 |
| Epoxiconazole | 5,227 | 63,108 | 37,974 | 2 |
| Fluxapyroxad | 5,100 | 53,782 | 38,390 | 2 |
| Fenpropimorph | 4,790 | 17,249 | 13,574 | 2 |

Pesticide usage on spring oats.

11,151 ha of spring oats grown in Ireland.

92,122 treated hectares.

27,686 kilogrammes applied.

Figure 42: Pesticide usage (spha) on spring oat crops in Ireland, 2016.




Figure 43: Weight of pesticides (kg) applied to spring oat crops in Ireland, 2016.




Figure 44: The top 10 active ingredients most extensively used on spring oats in Ireland in 2016, ranked by area treated (spray-hectares).

| Active substance | Treated area (spha) | Basic area treated | Quantity applied (kg) | % of the treated area |
|--------------------------|---------------------|--------------------|-----------------------|-----------------------|
| Fenpropimorph | 18,337 | 9,167 | 6,240 | 13.9 |
| Pyraclostrobin | 12,469 | 8,028 | 1,473 | 9.4 |
| Chlormequat chloride | 11,258 | 8,756 | 8,855 | 8.5 |
| Epoxiconazole | 10,860 | 6,920 | 685 | 8.2 |
| Tribenuron-methyl | 10,091 | 10,091 | 91 | 7.6 |
| Fluroxypyr | 8,448 | 7,060 | 1,200 | 6.4 |
| Thifensulfuron-methyl | 6,441 | 6,441 | 72 | 4.9 |
| Trinexapac-ethyl | 5,239 | 4,714 | 260 | 4.0 |
| Prochloraz/triticonazole | 5,155 | 5,155 | 129 | 3.9 |
| Metrafenone | 4,228 | 4,051 | 363 | 3.2 |

Figure 45: The top 10 active ingredients most extensively used on spring oats in Ireland in 2016, ranked by weight (kg).

| Active substance | Quantity applied (kg) | Treated area (spha) | Basic area treated | % of the weight applied |
|----------------------|-----------------------|---------------------|--------------------|-------------------------|
| Chlormequat chloride | 8,855 | 11,258 | 8,756 | 32.0 |
| Fenpropimorph | 6,240 | 18,337 | 9,167 | 22.5 |
| Glyphosate | 3,180 | 3,583 | 3,214 | 11.5 |
| Pyraclostrobin | 1,473 | 12,469 | 8,028 | 5.3 |
| Fluroxypyr | 1,200 | 8,448 | 7,060 | 4.3 |
| Mecoprop-P | 911 | 1,430 | 1,330 | 3.3 |
| Epoxiconazole | 685 | 10,860 | 6,920 | 2.5 |
| Fenpropidin | 463 | 1,215 | 1,134 | 1.7 |
| 2,4-DB | 398 | 237 | 237 | 1.4 |
| Dimethomorph | 397 | 397 | 397 | 1.4 |

Pesticide usage on winter oats.

13,146 ha of winter oats grown in Ireland.

132,247 treated hectares.

44,513 kilogrammes applied.

Figure 46: Pesticide usage (spha) on winter oat crops in Ireland, 2016.




Figure 47: Weight of pesticides (kg) applied to winter oat crops in Ireland, 2016.




Figure 48: The top 10 active ingredients most extensively used on winter oats in Ireland in 2016, ranked by area treated (spray-hectares).

| Active substance | Treated area (spha) | Basic area treated (ha) | Quantity applied (kg) | % of the treated area |
|--------------------------|---------------------|-------------------------|-----------------------|-----------------------|
| Fenpropimorph | 30,239 | 12,342 | 11,027 | 16.4 |
| Chlormequat chloride | 23,506 | 16,908 | 18,407 | 12.8 |
| Epoxiconazole | 16,159 | 10,003 | 1,228 | 8.8 |
| Pyraclostrobin | 14,527 | 10,466 | 1,930 | 7.9 |
| Trinexapac-ethyl | 11,099 | 8,355 | 562 | 6.0 |
| Metrafenone | 9,696 | 7,141 | 900 | 5.3 |
| Tribenuron-methyl | 9,389 | 9,389 | 103 | 5.1 |
| Fluroxypyr | 6,893 | 5,708 | 807 | 3.7 |
| Prochloraz/triticonazole | 6,330 | 6,330 | 153 | 3.4 |
| Proquinazid | 5,860 | 5,377 | 206 | 3.2 |

Figure 49: The top 10 active ingredients most extensively used on winter oats in Ireland in 2016, ranked by weight (kg).

| Active substance | Quantity applied (kg) | Treated area (spha) | Basic area treated (ha) | % of the weight applied |
|----------------------|-----------------------|---------------------|-------------------------|-------------------------|
| Chlormequat chloride | 18,407 | 23,506 | 16,908 | 41.4 |
| Fenpropimorph | 11,027 | 30,239 | 12,342 | 24.8 |
| Glyphosate | 4,941 | 5,423 | 5,256 | 11.1 |
| Pyraclostrobin | 1,930 | 14,527 | 10,466 | 4.3 |
| Epoxiconazole | 1,228 | 16,159 | 10,003 | 2.8 |
| Metrafenone | 900 | 9,696 | 7,141 | 2.0 |
| Fluroxypyr | 807 | 6,893 | 5,708 | 1.8 |
| Carboxin/thiram | 783 | 4,515 | 4,515 | 1.8 |
| Mecoprop-P | 731 | 907 | 907 | 1.6 |
| Tebuconazole | 678 | 3,410 | 3,114 | 1.5 |

Pesticide usage on oilseed rape.

9,912 ha of oilseed rape grown in Ireland.

72,582 treated hectares.

26,283 kilogrammes applied.

Figure 50: Pesticide usage (spha) on oilseed rape crops in Ireland, 2016.




Figure 51: Weight of pesticides (kg) applied to oilseed rape crops in Ireland, 2016.




Figure 52: The top 10 active ingredients most extensively used on oilseed rape in Ireland in 2016, ranked by area treated (spray-hectares).

| Active substance | Treated area (spha) | Basic area treated | Quantity applied (kg) | % of the treated area |
|--------------------|---------------------|--------------------|-----------------------|-----------------------|
| Glyphosate | 10,417 | 8,306 | 12,688 | 12.5 |
| Prothioconazole | 10,294 | 7,026 | 1,222 | 12.4 |
| Prochloraz/thiram | 8,960 | 8,960 | 156 | 10.8 |
| Metazachlor | 6,278 | 5,868 | 4,443 | 7.6 |
| Quinmerac | 6,222 | 5,811 | 1,467 | 7.5 |
| Lambda-cyhalothrin | 6,082 | 4,261 | 39 | 7.3 |
| Azoxystrobin | 5,284 | 3,732 | 804 | 6.4 |
| Metconazole | 5,088 | 4,578 | 226 | 6.1 |
| Propaquizafop | 4,768 | 4,662 | 294 | 5.7 |
| Boscalid | 4,362 | 4,362 | 1,012 | 5.3 |

Figure 53: The top 10 active ingredients most extensively used on oilseed rape in Ireland in 2016, ranked by weight (kg).

| Active substance | Quantity applied (kg) | Treated area (spha) | Basic area treated | % of the weight applied |
|------------------|-----------------------|---------------------|--------------------|-------------------------|
| Glyphosate | 12,688 | 10,417 | 8,306 | 48.3 |
| Metazachlor | 4,443 | 6,278 | 5,868 | 16.9 |
| Propyzamide | 2,322 | 3,245 | 2,934 | 8.8 |
| Quinmerac | 1,467 | 6,222 | 5,811 | 5.6 |
| Prothioconazole | 1,222 | 10,294 | 7,026 | 4.6 |
| Boscalid | 1,012 | 4,362 | 4,362 | 3.9 |
| Azoxystrobin | 804 | 5,284 | 3,732 | 3.1 |
| Tebuconazole | 435 | 3,252 | 2,539 | 1.7 |
| Propaquizafop | 294 | 4,768 | 4,662 | 1.1 |
| Diquat | 279 | 745 | 745 | 1.1 |

Pesticide usage on peas and beans.

11,016 ha of peas and beans grown in Ireland.

58,706 treated hectares.

27,070 kilogrammes applied.

Figure 54: Pesticide usage (spha) on pea and bean crops in Ireland, 2016.




Figure 55: Weight of pesticides (kg) applied to pea and bean crops in Ireland, 2016.




Figure 56: The top 10 active ingredients most extensively used on peas and beans in Ireland in 2016, ranked by area treated (spray-hectares).

| Active substance | Treated area (spha) | Basic area treated | Quantity applied (kg) | % of the treated area |
|--------------------|---------------------|--------------------|-----------------------|-----------------------|
| Pyraclostrobin | 12,540 | 8,479 | 589 | 14.3 |
| Boscalid | 12,439 | 8,479 | 2,299 | 14.2 |
| Chlorothalonil | 10,115 | 6,579 | 6,379 | 11.5 |
| Pendimethalin | 10,086 | 10,086 | 7,828 | 11.5 |
| Imazamox | 9,487 | 9,487 | 489 | 10.8 |
| Linuron | 6,506 | 6,506 | 2,305 | 7.4 |
| Clomazone | 6,305 | 6,305 | 394 | 7.2 |
| Lambda-cyhalothrin | 4,913 | 4,913 | 33 | 5.6 |
| Azoxystrobin | 3,816 | 3,144 | 577 | 4.3 |
| Metalaxy-M | 2,682 | 2,631 | 1,839 | 3.1 |

Figure 57: The top 10 active ingredients most extensively used on peas and beans in Ireland in 2016, ranked by weight (kg).

| Active substance | Quantity applied (kg) | Treated area (spha) | Basic area treated | % of the weight applied |
|------------------|-----------------------|---------------------|--------------------|-------------------------|
| Pendimethalin | 7,828 | 10,086 | 10,086 | 28.9 |
| Chlorothalonil | 6,379 | 10,115 | 6,579 | 23.6 |
| Linuron | 2,305 | 6,506 | 6,506 | 8.5 |
| Boscalid | 2,299 | 12,439 | 8,479 | 8.5 |
| Metalaxy-M | 1,839 | 2,682 | 2,631 | 6.8 |
| Glyphosate | 1,731 | 1,935 | 1,535 | 6.4 |
| Prosulfocarb | 1,215 | 419 | 419 | 4.5 |
| Pyraclostrobin | 589 | 12,540 | 8,479 | 2.2 |
| Azoxystrobin | 577 | 3,816 | 3,144 | 2.1 |
| Mancozeb | 568 | 398 | 398 | 2.1 |

Pesticide usage on seed potato crops.

308 ha of seed potato grown in Ireland.

9,003 treated hectares.

3,870 kilogrammes applied.

Figure 58: Pesticide usage (spha) on seed potato crops in Ireland, 2016.




Figure 59: Weight of pesticides (kg) applied to seed potato crops in Ireland, 2016.




Figure 60: The top 10 active ingredients most extensively used on seed potato crops in Ireland in 2016, ranked by area treated (spray-hectares).

| Active substance | Treated area (spha) | Basic area treated | Quantity applied (kg) | % of the treated area |
|---------------------------|---------------------|--------------------|-----------------------|-----------------------|
| Cymoxanil | 1,735 | 324 | 167 | 16.1 |
| Fluazinam | 1,445 | 324 | 264 | 13.4 |
| Diquat | 1,135 | 329 | 349 | 10.5 |
| Mandipropamid | 991 | 299 | 138 | 9.2 |
| Mancozeb | 950 | 310 | 1243 | 8.8 |
| Propamocarb hydrochloride | 892 | 297 | 870 | 8.3 |
| Fluopicolide | 793 | 263 | 79 | 7.4 |
| Flonicamid | 467 | 237 | 37 | 4.3 |
| Thiacloprid | 450 | 235 | 43 | 4.2 |
| Metribuzin | 308 | 308 | 170 | 2.9 |

Figure 61: The top 10 active ingredients most extensively used on seed potato crops in Ireland in 2016, ranked by weight (kg).

| Active substance | Quantity applied (kg) | Treated area (spha) | Basic area treated | % of the weight applied |
|---------------------------|-----------------------|---------------------|--------------------|-------------------------|
| Mancozeb | 1,243 | 950 | 310 | 32 |
| Propamocarb hydrochloride | 870 | 892 | 297 | 22 |
| Diquat | 349 | 1,135 | 329 | 9 |
| Fluazinam | 264 | 1,445 | 324 | 7 |
| Metribuzin | 170 | 308 | 308 | 4 |
| Cymoxanil | 167 | 1,735 | 324 | 4 |
| Mandipropamid | 138 | 991 | 299 | 4 |
| Linuron | 132 | 256 | 256 | 3 |
| Dimethoate | 90 | 268 | 249 | 2 |
| Fluopicolide | 79 | 793 | 263 | 2 |

Pesticide usage on early potato crops.

757 ha of early potato grown in Ireland.

8,535 treated hectares.

6,457 kilogrammes applied.

Figure 62: Pesticide usage (spha) on early potato crops in Ireland, 2016.




Figure 63: Weight of pesticides (kg) applied to early potato crops in Ireland, 2016.




Figure 64: The top 10 active ingredients most extensively used on early potato crops in Ireland in 2016, ranked by area treated (spray-hectares).

| Active substance | Treated area (spha) | Basic area treated | Quantity applied (kg) | % of the treated area |
|---------------------------|---------------------|--------------------|-----------------------|-----------------------|
| Mancozeb | 1,805 | 784 | 2,622 | 15.6 |
| Cymoxanil | 1,756 | 746 | 187 | 15.2 |
| Propamocarb hydrochloride | 1,561 | 799 | 1,450 | 13.5 |
| Fluazinam | 1,299 | 439 | 220 | 11.2 |
| Fluopicolide | 999 | 556 | 97 | 8.6 |
| Metribuzin | 673 | 673 | 306 | 5.8 |
| Diquat | 635 | 431 | 267 | 5.5 |
| Mandipropamid | 534 | 354 | 81 | 4.6 |
| Linuron | 386 | 386 | 174 | 3.3 |
| Glyphosate | 313 | 313 | 304 | 2.7 |

Figure 65: The top 10 active ingredients most extensively used on early potato crops in Ireland in 2016, ranked by weight (kg).

| Active substance | Quantity applied (kg) | Treated area (spha) | Basic area treated | % of the weight applied |
|---------------------------|-----------------------|---------------------|--------------------|-------------------------|
| Mancozeb | 2,622 | 1,805 | 784 | 40.6 |
| Propamocarb hydrochloride | 1,450 | 1,561 | 799 | 22.5 |
| Prosulfocarb | 391 | 127 | 127 | 6.1 |
| Metribuzin | 306 | 673 | 673 | 4.7 |
| Glyphosate | 304 | 313 | 313 | 4.7 |
| Diquat | 267 | 635 | 431 | 4.1 |
| Fluazinam | 220 | 1,299 | 439 | 3.4 |
| Cymoxanil | 187 | 1,756 | 746 | 2.9 |
| Linuron | 174 | 386 | 386 | 2.7 |
| Pendimethalin | 123 | 93 | 93 | 1.9 |

Pesticide usage on maincrop potato crops.

7,987 ha of maincrop potato grown in Ireland.

158,959 treated hectares.

83,956 kilogrammes applied.

Figure 66: Pesticide usage (spha) on maincrop potato in Ireland, 2016.




Figure 67: Weight of pesticides (kg) applied to maincrop potato in Ireland, 2016.




Figure 68: The top 10 active ingredients most extensively used on maincrop potato in Ireland in 2016, ranked by area treated (spray-hectares).

| Active substance | Treated area (spha) | Basic area treated | Quantity applied (kg) | % of the treated area |
|---------------------------|---------------------|--------------------|-----------------------|-----------------------|
| Fluazinam | 27,694 | 7,231 | 5,202 | 14.2 |
| Cymoxanil | 23,908 | 6,663 | 2,640 | 12.3 |
| Propamocarb hydrochloride | 21,219 | 7,391 | 17,582 | 10.9 |
| Diquat | 18,309 | 7,585 | 6,947 | 9.4 |
| Mandipropamid | 18,101 | 5,570 | 2,612 | 9.3 |
| Mancozeb | 17,790 | 6,021 | 25,624 | 9.1 |
| Fluopicolide | 17,536 | 7,179 | 1,471 | 9.0 |
| Cyazofamid | 11,412 | 4,506 | 1,936 | 5.9 |
| Metribuzin | 7,040 | 6,622 | 3,775 | 3.6 |
| Glyphosate | 4,562 | 4,562 | 4,291 | 2.3 |

Figure 69: The top 10 active ingredients most extensively used on maincrop potato in Ireland in 2016, ranked by weight (kg).

| Active substance | Quantity applied (kg) | Treated area (spha) | Basic area treated | % of the weight applied |
|---------------------------|-----------------------|---------------------|--------------------|-------------------------|
| Mancozeb | 25,624 | 17,790 | 6,021 | 30.5 |
| Propamocarb hydrochloride | 17,582 | 21,219 | 7,391 | 20.9 |
| Diquat | 6,947 | 18,309 | 7,585 | 8.3 |
| Prosulfocarb | 6,083 | 2,852 | 2,852 | 7.2 |
| Fluazinam | 5,202 | 27,694 | 7,231 | 6.2 |
| Glyphosate | 4,291 | 4,562 | 4,562 | 5.1 |
| Metribuzin | 3,775 | 7,040 | 6,622 | 4.5 |
| Cymoxanil | 2,640 | 23,908 | 6,663 | 3.1 |
| Mandipropamid | 2,612 | 18,101 | 5,570 | 3.1 |
| Cyazofamid | 1,936 | 11,412 | 4,506 | 2.3 |

Table 1: Estimated area (hectares) of arable crops grown regionally in Ireland, 2016.

| Crop | Region | | | |
|-------------------|----------------|----------------|---------------|----------------|
| | East | South | North/West | Ireland |
| Spring barley | 42,944 | 62,102 | 6,882 | 111,929 |
| Winter barley | 40,503 | 33,092 | 3,031 | 76,626 |
| Spring wheat | 4,691 | 6,344 | 89 | 11,125 |
| Winter wheat | 35,778 | 14,569 | 1,787 | 52,134 |
| Spring oats | 4,875 | 4,429 | 1,848 | 11,151 |
| Winter oats | 7,164 | 5,822 | 160 | 13,146 |
| Oilseed rape | 6,968 | 2,734 | 210 | 9,912 |
| Peas & beans | 6,682 | 4,334 | 0 | 11,016 |
| Seed potatoes | 216 | 41 | 51 | 308 |
| Early potatoes | 240 | 474 | 43 | 757 |
| Maincrop potatoes | 3,534 | 3,294 | 1,159 | 7,987 |
| Total | 153,597 | 137,235 | 15,260 | 306,092 |

Table 2a: Estimated area (spray-hectares) of arable crops treated regionally with each pesticide type in Ireland, 2016.

| Pesticide type | Region | | | |
|-------------------|------------------|------------------|----------------|------------------|
| | East | South | North/West | Ireland |
| Fungicides | 746,615 | 633,545 | 59,222 | 1,439,382 |
| Herbicides | 469,379 | 408,476 | 35,124 | 912,979 |
| Insecticides | 150,501 | 123,215 | 9,486 | 283,203 |
| Molluscicides | 5,182 | 4,448 | 865 | 10,495 |
| Growth Regulators | 199,404 | 146,582 | 8,866 | 354,852 |
| Seed treatments | 140,754 | 112,837 | 16,888 | 270,479 |
| Total | 1,711,836 | 1,429,103 | 130,452 | 3,271,391 |

Table 2b: Estimated weight (kg) applied to arable crops regionally with each pesticide type in Ireland, 2016.

| Pesticide type | Region | | | |
|-----------------------|----------------|----------------|-------------------|------------------|
| | East | South | North/West | Ireland |
| Fungicides | 248,155 | 212,682 | 25,088 | 485,925 |
| Herbicides | 168,389 | 149,176 | 15,250 | 332,815 |
| Insecticides | 15,091 | 5,165 | 428 | 20,684 |
| Molluscicides | 992 | 756 | 172 | 1,921 |
| Growth Regulators | 107,737 | 83,042 | 4,238 | 195,017 |
| Seed treatments | 10,746 | 10,496 | 858 | 22,100 |
| Total | 551,108 | 461,318 | 46,035 | 1,058,461 |

Table 3: The total area (spray hectares) and the basic area (hectares), of arable crops in Ireland 2016 treated with each pesticide type.

| Crop type | Pesticide Type | | | | | | | | | | | | | | |
|-------------------|------------------|----------------|----------------|----------------|----------------|----------------|---------------|--------------|-------------------|----------------|-----------------|----------------|------------------|----------------|----------------|
| | Fungicides | | Herbicides | | Insecticides | | Molluscicides | | Growth regulators | | Seed treatments | | All Pesticides | | |
| | (sp ha) | (ha) | (sp ha) | (ha) | (sp ha) | (ha) | (sp ha) | (ha) | (sp ha) | (ha) | (sp ha) | (ha) | (sp ha) | (ha) treated | (ha) grown |
| Spring barley | 411,893 | 108,934 | 322,138 | 110,618 | 94,609 | 86,131 | | | 44,987 | 33,786 | 105,338 | 102,501 | 978,966 | 111,917 | 111,929 |
| Winter barley | 363,349 | 76,310 | 249,960 | 75,356 | 67,928 | 57,239 | 1,090 | 1,090 | 134,012 | 64,003 | 70,847 | 69,258 | 887,186 | 76,330 | 76,626 |
| Spring wheat | 51,616 | 11,125 | 29,846 | 11,089 | 17,217 | 11,102 | | | 19,277 | 9,864 | 8,493 | 8,460 | 126,449 | 11,125 | 11,125 |
| Winter wheat | 350,205 | 51,806 | 165,307 | 51,742 | 70,987 | 46,707 | 3,930 | 3,930 | 105,347 | 47,110 | 50,861 | 49,331 | 746,636 | 52,134 | 52,134 |
| Spring oats | 33,961 | 10,868 | 25,722 | 10,512 | 7,347 | 6,689 | | | 16,624 | 8,296 | 8,468 | 7,941 | 92,122 | 11,151 | 11,151 |
| Winter oats | 50,209 | 13,141 | 27,655 | 12,400 | 7,449 | 5,916 | 231 | 231 | 34,605 | 12,213 | 12,098 | 12,098 | 132,247 | 13,146 | 13,146 |
| Oilseed rape | 27,574 | 9,006 | 28,041 | 9,672 | 6,428 | 4,284 | 1,578 | 1,425 | | | 8,960 | 8,807 | 72,582 | 9,874 | 9,912 |
| Peas & beans | 30,626 | 10,720 | 21,589 | 10,861 | 6,046 | 6,046 | 222 | 222 | | | 223 | 223 | 58,706 | 11,016 | 11,016 |
| Seed potatoes | 5,351 | 308 | 1,770 | 308 | 1,570 | 263 | 47 | 47 | | | 265 | 265 | 9,003 | 308 | 308 |
| Early potatoes | 5,697 | 757 | 2,326 | 757 | 288 | 185 | | | | | 223 | 223 | 8,535 | 757 | 757 |
| Maincrop potatoes | 108,900 | 7,831 | 38,624 | 7,831 | 3,333 | 1,969 | 3,396 | 2,130 | | | 4,704 | 4,704 | 158,959 | 7,987 | 7,987 |
| Total | 1,439,382 | 300,806 | 912,979 | 301,146 | 283,203 | 226,532 | 10,495 | 9,076 | 354,852 | 175,272 | 270,479 | 263,811 | 3,271,391 | 305,744 | 306,092 |

Table 4: The total quantities (kilograms) of each pesticide type used on arable crops in Ireland 2016.

| <i>Crop</i> | <i>Pesticide type</i> | | | | | | |
|-------------------------|-----------------------|-------------------|---------------------|----------------------|--------------------------|------------------------|----------------------------------|
| | Fungicides | Herbicides | Insecticides | Molluscicides | Growth regulators | Seed treatments | Total weight applied (kg) |
| Spring barley | 123,970 | 68,360 | 3,959 | | 23,538 | 8,145 | 227,973 |
| Winter barley | 113,059 | 116,563 | 2,125 | 154 | 68,315 | 5,777 | 305,992 |
| Spring wheat | 18,908 | 8,336 | 2,229 | | 13,599 | 977 | 44,050 |
| Winter wheat | 119,862 | 63,194 | 11,047 | 829 | 61,463 | 4,216 | 260,613 |
| Spring oats | 10,938 | 6,761 | 264 | | 9,133 | 589 | 27,686 |
| Winter oats | 17,096 | 7,269 | 81 | 46 | 18,969 | 1,052 | 44,513 |
| Oilseed rape | 3,926 | 21,772 | 143 | 286 | | 156 | 26,283 |
| Peas & beans | 12,534 | 14,328 | 141 | 44 | | 22 | 27,070 |
| Seed potatoes | 2,835 | 716 | 286 | 7 | | 25 | 3,870 |
| Early potatoes | 4,756 | 1,576 | 108 | | | 17 | 6,457 |
| Maincrop potatoes | 58,041 | 23,940 | 300 | 553 | | 1,123 | 83,956 |
| <i>All arable crops</i> | 485,925 | 332,815 | 20,684 | 1,921 | 195,017 | 22,100 | 1,058,461 |

Table 5: Estimated area (spray-hectares) of arable crops treated with pesticide formulations in Ireland, 2016.

| Pesticide type & formulation | Spring barley | Winter barley | Spring wheat | Winter wheat | Crop | | | | | | | Maincrop potatoes | All crops |
|--------------------------------------|---------------|---------------|--------------|--------------|-------------|-------------|--------------|--------------|---------------|----------------|-------------------|-------------------|-----------|
| | | | | | Spring oats | Winter oats | Oilseed rape | Peas & beans | Seed potatoes | Early potatoes | Maincrop potatoes | | |
| Fungicides | | | | | | | | | | | | | |
| Azoxystrobin | 487 | 2,943 | 303 | 6,316 | 190 | 1,919 | 5,105 | 3,178 | | 36 | 904 | 21,380 | |
| Azoxystrobin/chlorothalonil | 1,888 | 2,244 | 1,707 | 12,082 | | | 179 | 222 | | | 161 | 18,484 | |
| Azoxystrobin/difenoconazole | | | | | 441 | | | 416 | | | | 857 | |
| Benthiavalicarb-isopropyl/mancozeb | | | | | | | | | 14 | 288 | 2,752 | 3,055 | |
| Bixafen/prothioconazole | 41,989 | 32,629 | 1,539 | 9,717 | 437 | | | 117 | | | | 86,428 | |
| Bixafen/prothioconazole/spiroxamine | | | | 1,427 | | | | | | | | 1,427 | |
| Bixafen/prothioconazole/tebuconazole | | | 1,852 | 10,259 | | | 714 | | | | | 12,825 | |
| Boscalid | | | | | | | 4,362 | | | | | 4,362 | |
| Boscalid/epoxiconazole | 16,430 | 18,315 | 4,093 | 11,997 | 525 | 403 | | | | | | 51,764 | |
| Boscalid/pyraclostrobin | | | | | | | | 12,439 | | | | 12,439 | |
| Chlorothalonil | 104,392 | 100,223 | 17,019 | 113,393 | 707 | 297 | | 9,893 | | | | 345,923 | |
| Chlorothalonil/penthiopyrad | 6,163 | 5,025 | 1,168 | 7,059 | | | | | | | | 19,415 | |
| Chlorothalonil/picoxytirobin | 3,464 | 2,040 | 290 | 1,615 | | | | | | | | 7,410 | |
| Chlorothalonil/propiconazole | | | 553 | | | | | | | | | 553 | |
| Chlorothalonil/proquinazid | 109 | 935 | 1,273 | 4,565 | | | | | | | | 6,883 | |
| Cyazofamid | | | | | | | | 244 | 181 | 11,412 | | 11,837 | |
| Cymoxanil | | | | | 78 | | | 795 | 262 | 11,326 | | 12,460 | |
| Cymoxanil/mancozeb | | | | | | | | 885 | 1,176 | 11,056 | | 13,118 | |
| Cymoxanil/propamocarb hydrochloride | | | | | | | | 55 | 318 | 1,526 | | 1,900 | |
| Cyproconazole/penthiopyrad | | 132 | | 171 | | | | | | | | 303 | |
| Cyproconazole/picoxytirobin | | | | | 71 | 86 | | | | | | 158 | |
| Cyprodinil/isopyrazam | 16,400 | 13,368 | 806 | 118 | | | | | | | | 30,692 | |
| Difenoconazole | | 212 | | 712 | | | | | | | | 924 | |
| Difenoconazole/tebuconazole | | | | 3,046 | | | | | | | | 3,046 | |
| Dimethomorph | 463 | | | | 397 | | | | 63 | | | 860 | |
| Dimethomorph/mancozeb | | | | | | | | | | | | 63 | |

Table 5 (cont.): Estimated area (spray-hectares) of arable crops treated with pesticide formulations in Ireland, 2016.

| Pesticide type & formulation | Spring barley | Winter barley | Spring wheat | Winter wheat | Spring oats | Winter oats | Oilseed rape | Peas & beans | Seed potatoes | Early potatoes | Maincrop potatoes | All crops |
|-----------------------------------------------|---------------|---------------|--------------|--------------|-------------|-------------|--------------|--------------|---------------|----------------|-------------------|-----------|
| Fungicides (cont.) | | | | | | | | | | | | |
| Epoxiconazole | 3,847 | 1,346 | 2,311 | 3,037 | 1,450 | 2,054 | 93 | | | | | 14,138 |
| Epoxiconazole/fenpropimorph | 7,596 | 3,982 | 142 | 2,840 | 720 | 440 | | | | | | 15,719 |
| Epoxiconazole/fenpropimorph/kresoxim-methyl | 1,151 | 1,854 | | 1,299 | | 2,205 | | | | | | 6,510 |
| Epoxiconazole/fenpropimorph/metrafenone | 326 | | 721 | 4,208 | 4,034 | 8,852 | | | | | | 18,140 |
| Epoxiconazole/fenpropimorph/pyraclostrobin | 10,556 | 5,884 | | | 1,804 | 840 | | | | | | 19,084 |
| Epoxiconazole/fluxapyroxad | 652 | 117 | 1,045 | 16,060 | 289 | | | | | | 90 | 18,253 |
| Epoxiconazole/fluxapyroxad/pyraclostrobin | 21,810 | 21,316 | | 1,193 | 1,909 | 985 | | | | | | 47,213 |
| Epoxiconazole/isopyrazam | | | | 601 | | | | | | | | 601 |
| Epoxiconazole/metconazole | 1,212 | 484 | 1,164 | 21,511 | | | | | | | | 24,372 |
| Epoxiconazole/pyraclostrobin | 4,079 | 930 | 966 | 362 | 129 | 381 | | | | | | 6,847 |
| Fenamidone/propamocarb hydrochloride | | | | | | | | | 43 | 244 | 2,157 | 2,444 |
| Fenpropidin | 4,818 | 341 | 1,543 | 1,073 | 1,215 | 547 | | | | | | 9,536 |
| Fenpropimorph | 35,342 | 10,830 | 2,124 | 4,136 | 4,331 | 7,056 | | | | | | 63,820 |
| Fenpropimorph/flusilazole | | | | 564 | | | | | | | | 564 |
| Fenpropimorph/pyraclostrobin | 7,047 | 9,341 | 388 | 4,202 | 7,448 | 10,847 | | | | | | 39,272 |
| Fluazinam | | | | | | | | | 1,445 | 1,299 | 27,684 | 30,429 |
| Fluazinam/metalaxyl-M | | | | | | | | | | | 9 | 9 |
| Fluopicolide | | | 620 | | | | | | | | | 620 |
| Fluopicolide/propamocarb hydrochloride | | | | | | | | | 793 | 999 | 17,536 | 19,328 |
| Fluoxastrobin | 4,693 | 3,548 | 290 | 352 | | | | | | | | 8,883 |
| Fluoxastrobin/prothioconazole | 1,104 | 751 | | | 50 | 258 | | | | | | 2,162 |
| Fluoxastrobin/prothioconazole/trifloxystrobin | 12,155 | 11,915 | | 402 | | | | | | | | 24,472 |
| Fluxapyroxad | 3,735 | 8,469 | 68 | 13,319 | 100 | | | | | | | 25,692 |
| Fluxapyroxad/metconazole | 652 | 1,087 | 106 | 23,210 | 292 | 297 | | | | | | 25,645 |
| Folpet | 1,608 | 1,678 | 880 | 710 | | | | | | | | 4,876 |
| Iprodione | | | | 626 | | | | | | | | 626 |
| Isopyrazam | 3,503 | 7,244 | | | | | | | | | | 10,748 |
| Mancozeb | | | | 161 | | | | | 335 | 51 | 340 | 3,821 |
| Mancozeb/zoxamide | | | | | | | | | | | 161 | 4,708 |
| | | | | | | | | | | | | 161 |

Table 5 (cont.): Estimated area (spray-hectares) of arable crops treated with pesticide formulations in Ireland, 2016.

| Pesticide type & formulation | Crop | | | | | | | | | | | |
|------------------------------------------|----------------|----------------|---------------|----------------|---------------|---------------|---------------|---------------|---------------|----------------|-------------------|------------------|
| | Spring barley | Winter barley | Spring wheat | Winter wheat | Spring oats | Winter oats | Oilseed rape | Peas & beans | Seed potatoes | Early potatoes | Maincrop potatoes | All crops |
| Fungicides (cont.) | | | | | | | | | | | | |
| Mandipropamid | 586 | | | | | | | | 991 | 534 | 18,101 | 20,212 |
| Metalaxyl-M | | | | | | | | 2,682 | 34 | 21 | 203 | 2,940 |
| Metconazole | 616 | 591 | | 387 | | 112 | 5,088 | | | | | 6,794 |
| Metrafenone | | 1,588 | 161 | 1,926 | 194 | 845 | | | | | | 4,713 |
| Penthiopyrad | 663 | 836 | | 2,503 | | | | | | | | 4,002 |
| Penthiopyrad/picoxystrobin | 2,001 | 4,859 | | 206 | | 86 | 401 | | | | | 7,553 |
| Picoxystrobin | 2,076 | 615 | | | | 478 | | | | | | 3,168 |
| Prochloraz | | | | 896 | | | | | | | | 896 |
| Prochloraz/tebuconazole | | | 880 | | | | | | | | | 880 |
| Proquinazid | 1,999 | 2,498 | 3,274 | 4,605 | 3,607 | 5,860 | | | | | | 21,844 |
| Prothioconazole | 47,468 | 30,560 | 2,011 | 13,333 | | 478 | 9,094 | | | | | 102,942 |
| Prothioconazole/spiroxamine | 24,902 | 26,799 | 142 | 2,181 | 388 | | | | | | | 54,412 |
| Prothioconazole/spiroxamine/tebuconazole | | 669 | 221 | 7,562 | | | | | | | | 8,452 |
| Prothioconazole/tebuconazole | | 183 | 1,551 | 20,890 | 100 | | 487 | | | | | 23,212 |
| Prothioconazole/trifloxystrobin | 5,930 | 4,371 | | 429 | | | | | | | | 10,730 |
| Pyraclostrobin | 7,137 | 18,372 | 334 | 5,639 | 1,179 | 1,475 | | 102 | | | | 34,237 |
| Sulphur | 247 | 276 | | | | | | | | | | 523 |
| Tebuconazole | 594 | 201 | 691 | 7,103 | 1,876 | 3,410 | 2,051 | 1,180 | | | | 17,106 |
| Thiram | | | | 201 | | | | | | | | 201 |
| Tolclofos-methyl | | 1,128 | | | | | | | | | | 1,128 |
| All fungicides | 411,893 | 363,349 | 51,616 | 350,205 | 33,961 | 50,209 | 27,574 | 30,626 | 5,351 | 5,697 | 108,900 | 1,439,382 |

Table 5 (cont.): Estimated area (spray-hectares) of arable crops treated with pesticide formulations in Ireland, 2016.

| Pesticide type & formulation | Spring barley | Winter barley | Spring wheat | Winter wheat | Spring oats | Winter oats | Oilseed rape | Peas & beans | Seed potatoes | Early potatoes | Maincrop potatoes | All crops |
|---------------------------------------------------|---------------|---------------|--------------|--------------|-------------|-------------|--------------|--------------|---------------|----------------|-------------------|-----------|
| Herbicides | | | | | | | | | | | | |
| 2,4-D | | | | 712 | | | | | | | | 712 |
| 2,4-D/triclopyr | 270 | | | | | | | | | | | 270 |
| 2,4-DB/MCPA | 791 | | | 282 | 237 | | | | | | | 1,310 |
| 2,4-DB/mecoprop-P | 101 | | | | | | | | | | | 101 |
| Aminopyralid/propyzamide | | | | | | | 1,433 | | | | | 1,433 |
| Asulam | | | | 146 | | | | | | | | 146 |
| Bentazone | | | | | | | | 88 | | | | 88 |
| Carfentrazone-ethyl | | | | | | | | | | | 1,522 | 1,522 |
| Clodinafop-Propargyl/cloquintocet-mexyl/pinoxaden | | | | 176 | | | | | | | | 176 |
| Clomazone/linuron | | | | | | | | 6,305 | | 487 | | 6,792 |
| Clopyralid | | | | | | | 193 | | | | | 193 |
| Clopyralid/florasulam/fluroxypyr | 37,882 | 4,591 | 2,215 | 3,438 | 1,454 | 2,814 | | | | | | 52,393 |
| Clopyralid/picloram | | | | | | | 263 | | | | | 263 |
| Clopyralid/triclopyr | | 238 | | 429 | | | | | | | | 667 |
| Cycloxydim | | | | | 260 | | 600 | 1,167 | | | | 2,028 |
| Dicamba/MCPA/mecoprop-P | 656 | | | | | | | | | | | 656 |
| Dicamba/mecoprop-P | 1,509 | | 303 | | 88 | 93 | | | | | | 1,993 |
| Dichlorprop-P/MCPA/mecoprop-P | 4,330 | | 295 | | 277 | | | | | | | 4,902 |
| Diflufenican | 2,202 | 51,338 | 140 | 25,313 | 397 | 4,175 | | | | 36 | | 83,599 |
| Diflufenican/flufenacet | | 4,949 | | 4,029 | | | | | | | | 8,978 |
| Diflufenican/iodosulfuron-methyl-sodium | | | | 10,715 | | | | | | | | 10,715 |
| Diquat | 1,145 | | | | | | 745 | 24 | 1,135 | 635 | 18,309 | 21,993 |
| Ethametsulfuron-methyl | | | | | | | 618 | | | | | 618 |
| Ethofumesate | 826 | | | | | | | | | | | 826 |
| Fenoxaprop-P-ethyl | 1,509 | 441 | 457 | 5,917 | | | | | | | | 8,324 |
| Florasulam | 683 | 3,199 | | 2,615 | | 326 | | | | | | 6,822 |
| Florasulam/fluroxypyr | 3,138 | 2,696 | | 495 | | 438 | | | | | | 6,767 |
| Florasulam/pyroxslam | | 952 | | 3,367 | | | | | | | | 4,320 |
| Fluazifop-P-butyl | | | | | 260 | | 140 | | | | | 400 |
| Flumioxazin | | | | 33 | | | | | | | | 33 |

Table 5 (cont.): Estimated area (spray-hectares) of arable crops treated with pesticide formulations in Ireland, 2016.

| Pesticide type & formulation | Spring barley | Winter barley | Spring wheat | Winter wheat | Spring oats | Winter oats | Oilseed rape | Peas & beans | Seed potatoes | Early potatoes | Maincrop potatoes | All crops |
|------------------------------------------------|----------------|----------------|---------------|----------------|---------------|---------------|---------------|---------------|---------------|----------------|-------------------|----------------|
| Herbicides (cont.) | | | | | | | | | | | | |
| Flupyrifluron-methyl | | | | | | 157 | | | | | | 157 |
| Fluroxypyr | 42,048 | 10,829 | 6,653 | 11,743 | 6,994 | 3,641 | | | | | | 81,909 |
| Glufosinate-ammonium | | | | | | | | | | 11 | 183 | 193 |
| Glyphosate | 36,952 | 34,212 | 6,282 | 23,672 | 3,583 | 5,423 | 10,417 | 1,935 | 72 | 313 | 4,562 | 127,423 |
| Imazamox/pendimethalin | | | 130 | | | | 93 | 9,487 | | | | 9,711 |
| Iodosulfuron-methyl-sodium | 1,636 | | | | | | | | | | | 1,636 |
| Iodosulfuron-methyl-sodium/mesosulfuron-methyl | | | | 13,664 | | | | | | | | 13,664 |
| Isoproturon | 4,178 | 63,466 | 932 | 31,600 | | | | | | | | 100,176 |
| Linuron | | | | | 328 | | | | 200 | 256 | 386 | 2,155 |
| MCPA | | | | | | | | | | | | 328 |
| Mecoprop-P | 9,393 | 1,445 | 149 | 3,134 | 1,066 | 814 | | | | | | 16,002 |
| Metazachlor | 791 | | | | 360 | | 57 | | | | | 1,208 |
| Metazachlor/quinmerac | | | | | | 6,222 | | | | | | 6,222 |
| Metribuzin | | | | | | | | | 308 | 673 | 7,040 | 8,021 |
| Metsulfuron-methyl | 7,219 | 1,256 | 4,036 | | 67 | 299 | | | | | | 12,878 |
| Metsulfuron-methyl/thifensulfuron-methyl | 27,140 | 3,082 | 370 | 1,571 | | 86 | | 63 | | | | 32,312 |
| Metsulfuron-methyl/tribenuron-methyl | 24,799 | 3,923 | 1,980 | 608 | 3,209 | 1,646 | | | | | | 36,165 |
| Pendimethalin | 791 | | | | | | 598 | | | 93 | 788 | 2,270 |
| Pendimethalin/picolinafen | 326 | 4,210 | 563 | 346 | | | | | | | | 5,445 |
| Pinoxaden | 62,578 | 41,473 | 1,695 | 9,181 | | | | | | | | 114,927 |
| Propaquizafop | | | | | 260 | | 4,768 | 1,257 | | 17 | 258 | 6,560 |
| Propyzamide | | | | | | | 1,813 | 44 | | | | 1,857 |
| Prosulfocarb | 94 | 241 | | 1,017 | | | | 419 | | 127 | 2,852 | 4,751 |
| Rimsulfuron | | | | | | | | | | 36 | 470 | 506 |
| Tepraloxydim | | 198 | | 185 | | | 681 | | | | | 1,064 |
| Thifensulfuron-methyl/tribenuron-methyl | 48,517 | 11,686 | 3,623 | 8,726 | 6,441 | 5,561 | | | | | | 84,554 |
| Tribenuron-methyl | 634 | 5,118 | 23 | 2,191 | 441 | 2,181 | | | | | | 10,589 |
| Triflusulfuron-methyl | | | 418 | | | | | | | | | 418 |
| All herbicides | 322,138 | 249,960 | 29,846 | 165,307 | 25,722 | 27,655 | 28,041 | 21,589 | 1,770 | 2,326 | 38,624 | 912,979 |

Table 5 (cont.): Estimated area (spray-hectares) of arable crops treated with pesticide formulations in Ireland, 2016.

| Pesticide type & formulation | Crop | | | | | | | | | | | |
|----------------------------------------|---------------|---------------|---------------|---------------|--------------|--------------|--------------|--------------|---------------|----------------|-------------------|----------------|
| | Spring barley | Winter barley | Spring wheat | Winter wheat | Spring oats | Winter oats | Oilseed rape | Peas & beans | Seed potatoes | Early potatoes | Maincrop potatoes | All crops |
| Insecticides | | | | | | | | | | | | |
| Acetamiprid | | | | | | | | | 14 | | | 14 |
| Chlorpyrifos | 1,796 | | | 382 | | | | | | | | 2,178 |
| Clofentezine | | | | | | | | 221 | | | | 221 |
| Cypermethrin | 23,443 | 8,323 | 3,828 | 4,029 | 2,186 | 1,619 | 23 | 343 | | | | 43,794 |
| Deltamethrin | 3,526 | 3,730 | 560 | 1,702 | 306 | 93 | | 128 | | | | 10,044 |
| Dimethoate | 708 | 2,453 | 4,508 | 30,791 | 585 | | 323 | | 268 | | 903 | 40,537 |
| Esfenvalerate | 5,587 | 9,301 | 2,245 | 2,609 | 836 | | | 63 | | | | 20,641 |
| Ethoprophos | | | | | | | | | 19 | | | 19 |
| Flonicamid | | | | | | | | | 467 | | | 467 |
| Garlic Extract | 2,793 | | 880 | | | | | | | | | 3,674 |
| Lambda-cyhalothrin | 56,757 | 43,796 | 4,814 | 31,154 | 3,433 | 5,737 | 6,082 | 4,913 | 80 | 253 | 2,430 | 159,448 |
| Metarhizium anisopliae (BIPESCO 5F/52) | | 327 | | 703 | | | | | | | | 1,030 |
| Oxamyl | | | | | | | | | 36 | | | 36 |
| Pirimicarb | | | | | | | | 378 | | | | 378 |
| Pymetrozine | | | | | | | | | 272 | | | 272 |
| Thiacloprid | | | | | | | | | 450 | | | 450 |
| All Insecticides | 94,609 | 67,928 | 17,217 | 70,987 | 7,347 | 7,449 | 6,428 | 6,046 | 1,570 | 288 | 3,333 | 283,203 |
| Molluscicides | | | | | | | | | | | | |
| Metaldehyde | | 649 | | 3,930 | | 231 | 1,186 | 222 | 40 | | 3,396 | 9,654 |
| Methiocarb | | 442 | | | | | 392 | 0 | 7 | | | 842 |
| All molluscicides | 1,090 | 0 | 3,930 | 0 | 231 | 1,578 | 222 | 47 | 0 | 3,396 | 10,495 | |

Table 5 (cont.): Estimated area (spray-hectares) of arable crops treated with pesticide formulations in Ireland, 2016.

| Pesticide type & formulation | Crop | | | | | | | | | | | |
|-----------------------------------------|----------------|----------------|----------------|----------------|---------------|----------------|---------------|---------------|---------------|----------------|-------------------|------------------|
| | Spring barley | Winter barley | Spring wheat | Winter wheat | Spring oats | Winter oats | Oilseed rape | Peas & beans | Seed potatoes | Early potatoes | Maincrop potatoes | All crops |
| Growth regulators | | | | | | | | | | | | |
| Chlormequat chloride | 29,484 | 63,155 | 15,510 | 65,192 | 11,258 | 23,506 | | | | | | 208,104 |
| Chlormequat/imazaquin | | 326 | | 2,826 | | | | | | | | 3,152 |
| Ethephon (2-chloroethylphosphonic acid) | 792 | 6,420 | | 2,658 | | | | | | | | 9,869 |
| Ethephon/mepiquat chloride | 1,795 | 23,367 | 233 | 1,509 | | | | | | | | 26,905 |
| Mepiquat chloride/prohexadione-calcium | | | | | 127 | | | | | | | 127 |
| Trinexapac-ethyl | 12,589 | 41,070 | 3,534 | 33,162 | 5,239 | 11,099 | | | | | | 106,694 |
| All growth regulators | 44,987 | 134,012 | 19,277 | 105,347 | 16,624 | 34,605 | 0 | 0 | 0 | 0 | 0 | 354,852 |
| Seed treatments | | | | | | | | | | | | |
| Carboxin/thiram | 28,313 | 16,237 | 4,708 | 13,217 | 1,908 | 4,515 | | | | | | 68,898 |
| Clothianidin/prothioconazole | 12,348 | 20,429 | | 13,022 | 1,242 | 1,254 | | | | | | 48,295 |
| Cymoxanil/fludioxonil/metalexyl-M | | | | | | | | | | | | 223 |
| Fludioxonil | 183 | | 161 | 161 | 162 | | | | | | | 668 |
| Fludioxonil/flutriafol | | 166 | | | | | | | | | | 166 |
| Fluopyram/prothioconazole/tebuconazole | 1,658 | 86 | | | | | | | | | | 1,744 |
| Flutolanil | | | | | | | | | 31 | 1,127 | | 1,158 |
| Imazalil | | | | | | | | 213 | 175 | 2,114 | | 2,502 |
| Imazalil/pencycuron | | | | | | | | 41 | 17 | 1,463 | | 1,520 |
| Imazalil/thiabendazole | | | | | | | | 11 | | | | 11 |
| Prochloraz/thiram | | | | | | 8,960 | | | | | | 8,960 |
| Prochloraz/triticonazole | 51,576 | 29,010 | 3,624 | 18,964 | 5,155 | 6,330 | | | | | | 114,660 |
| Silthiofam | 11,260 | 4,919 | | 5,496 | | | | | | | | 21,674 |
| All seed treatments | 105,338 | 70,847 | 8,493 | 50,861 | 8,468 | 12,098 | 8,960 | 0 | 265 | 223 | 4,704 | 270,479 |
| All pesticides | 978,966 | 887,186 | 126,449 | 746,636 | 92,122 | 132,247 | 72,582 | 58,483 | 9,003 | 8,535 | 158,959 | 3,271,391 |

Table 6: Estimated quantities (kilograms) of pesticide formulations used on arable crops in Ireland, 2016.

| Pesticide type & formulation | Spring | Winter | Spring | Winter | Spring | Crop | Oilseed | Peas & | Seed | Early | Maincrop | All crops |
|--------------------------------------|--------|--------|--------|--------|--------|--------|---------|--------|-------|-------|----------|-----------|
| | barley | barley | wheat | wheat | oats | Winter | | | | | potatoes | |
| Fungicides | | | | | | | | | | | | |
| Azoxystrobin | 65 | 474 | 36 | 1,197 | 24 | 245 | 785 | 475 | | 9 | 246 | 3,556 |
| Azoxystrobin/chlorothalonil | 1,270 | 1,338 | 941 | 6,885 | | | 116 | 115 | | | 77 | 10,742 |
| Azoxystrobin/difenoconazole | | | | | 67 | | | 135 | | | | 202 |
| Benthiavalicarb-isopropyl/mancozeb | | | | | | | | | 16 | 331 | 2,514 | 2,861 |
| Bixafen/prothioconazole | 7,462 | 5,676 | 489 | 2,187 | 58 | | | 15 | | | | 15,888 |
| Bixafen/prothioconazole/spiroxamine | | | | 758 | | | | | | | | 758 |
| Bixafen/prothioconazole/tebuconazole | | | 509 | 3,152 | | | 152 | | | | | 3,813 |
| Boscalid | | | | | | | 1,012 | | | | | 1,012 |
| Boscalid/epoxiconazole | 3,705 | 4,667 | 1,301 | 4,226 | 79 | 106 | | | | | | 14,084 |
| Boscalid/pyraclostrobin | | | | | | | 2,876 | | | | | 2,876 |
| Chlorothalonil | 50,876 | 48,819 | 8,424 | 57,946 | 351 | 148 | | 6,283 | | | | 172,848 |
| Chlorothalonil/penthiopyrad | 3,153 | 3,008 | 674 | 4,586 | | | | | | | | 11,421 |
| Chlorothalonil/picoxytrobacin | 1,174 | 1,424 | 348 | 1,065 | | | | | | | | 4,011 |
| Chlorothalonil/propiconazole | | | 173 | | | | | | | | | 173 |
| Chlorothalonil/proquinazid | 11 | 532 | 849 | 2,454 | | | | | | | | 3,847 |
| Cyazofamid | | | | | | | 26 | 24 | 1,936 | 1,936 | 1,986 | |
| Cymoxanil | | | | | 8 | | 87 | 29 | 1,313 | 1,313 | 1,436 | |
| Cymoxanil/mancozeb | | | | | | | | 1,207 | 1,923 | 1,923 | 18,637 | 21,767 |
| Cymoxanil/propamocarb hydrochloride | | | | | | | 50 | 347 | 1,469 | 1,469 | 1,866 | |
| Cyproconazole/penthiopyrad | | 44 | | 47 | | 13 | 16 | | | | | 91 |
| Cyproconazole/picoxytrobacin | | | | | | | | | | | | 30 |
| Cyprodinil/isopyrazam | 6,463 | 4,970 | 302 | 29 | | | | | | | | 11,764 |
| Difenoconazole | | 16 | | 53 | | | | | | | | 69 |
| Difenoconazole/tebuconazole | | | | 936 | | | | | | | | 936 |
| Dimethomorph | 463 | | | | 397 | | | | | | | 860 |

Table 6 (cont.): Estimated quantities (kilograms) of pesticide formulations used on arable crops in Ireland, 2016.

| Pesticide type & formulation | Spring barley | Winter barley | Spring wheat | Winter wheat | Spring oats | Winter oats | Oilseed rape | Peas & beans | Seed potatoes | Early potatoes | Maincrop potatoes | All crops |
|-----------------------------------------------|---------------|---------------|--------------|--------------|-------------|-------------|--------------|--------------|---------------|----------------|-------------------|-----------|
| Fungicides (cont.) | | | | | | | | | | | | |
| Dimethomorph/mancozeb | | | | | | | | 35 | | | | 35 |
| Epoxiconazole | 314 | 75 | 263 | 342 | 127 | 150 | 12 | | | | | 1,283 |
| Epoxiconazole/fenpropimorph | 2,112 | 1,348 | 59 | 1,077 | 209 | 128 | | | | | | 4,933 |
| Epoxiconazole/fenpropimorph/kresoxim-methyl | 361 | 703 | | 423 | | 750 | | | | | | 2,236 |
| Epoxiconazole/fenpropimorph/metrafenone | 110 | | 316 | 1,973 | 1,577 | 3,534 | | | | | | 7,510 |
| Epoxiconazole/fenpropimorph/pyraclostrobin | 3,003 | 2,087 | | | 567 | 422 | | | | | | 6,079 |
| Epoxiconazole/fluxapyroxad | 82 | 29 | 177 | 3,019 | 9 | | | | | 16 | | 3,331 |
| Epoxiconazole/fluxapyroxad/pyraclostrobin | 5,017 | 4,903 | | 429 | 429 | 228 | | | | | | 11,006 |
| Epoxiconazole/isopyrazam | | | | 147 | | | | | | | | 147 |
| Epoxiconazole/metconazole | 147 | 63 | 87 | 2,258 | | | | | | | | 2,555 |
| Epoxiconazole/pyraclostrobin | 567 | 146 | 170 | 51 | 23 | 73 | | | | | | 1,031 |
| Fenamidone/propamocarb hydrochloride | | | | | | | | 39 | 201 | 1,882 | | 2,122 |
| Fenpropidin | 2,659 | 255 | 880 | 409 | 463 | 125 | | | | | | 4,791 |
| Fenpropimorph | 11,232 | 3,176 | 752 | 1,237 | 1,308 | 3,027 | | | | | | 20,733 |
| Fenpropimorph/flusilazole | | | | 272 | | | | | | | | 272 |
| Fenpropimorph/pyraclostrobin | 2,984 | 3,812 | 169 | 1,556 | 4,450 | 6,693 | | | | | | 19,664 |
| Fluazinam | | | | | | | | 264 | 220 | 5,200 | | 5,685 |
| Fluazinam/metalaxyl-M | | | | | | | | | | 3 | | 3 |
| Fluopicolide | | 116 | | | | | | | | | | 116 |
| Fluopicolide/propamocarb hydrochloride | | | | | | | | 873 | 1,071 | 16,178 | | 18,122 |
| Fluoxastrobin | 437 | 363 | 36 | 35 | | | | | | | | 872 |
| Fluoxastrobin/prothioconazole | 205 | 182 | | | 9 | 54 | | | | | | 450 |
| Fluoxastrobin/prothioconazole/trifloxystrobin | 2,202 | 2,442 | | 91 | | | | | | | | 4,734 |
| Fluxapyroxad | 214 | 635 | 3 | 1,155 | 4 | | | | | | | 2,012 |
| Fluxapyroxad/metconazole | 70 | 70 | 11 | 3,983 | 50 | 16 | | | | | | 4,201 |
| Folpet | 804 | 1,112 | 440 | 410 | | | | | | | | 2,765 |

Table 6 (cont.): Estimated quantities (kilograms) of pesticide formulations used on arable crops in Ireland, 2016.

| Pesticide type & formulation | Spring barley | Winter barley | Spring wheat | Winter wheat | Spring oats | Winter oats | Crop | | | | Early potatoes | Maincrop potatoes | All crops |
|------------------------------------------|----------------|----------------|---------------|----------------|---------------|---------------|--------------|---------------|---------------|--------------|----------------|-------------------|-----------|
| | | | | | | | Oilseed rape | Peas & beans | Seed potatoes | | | | |
| Fungicides (cont.) | | | | | | | | | | | | | |
| Iprodione | | | | | | 188 | | | | | | 188 | |
| Isopyrazam | 239 | 478 | | | | 181 | | | 536 | 96 | 496 | 5,504 | |
| Mancozeb | | | | | | | | | | | | 6,813 | |
| Mancozeb/zoxamide | | | | | | | | | | | | 217 | |
| Mandipropamid | 73 | | | | | | | | | 138 | 81 | 2,612 | |
| Metalaxyl-M | | | | | | | | | 1,839 | 40 | 24 | 236 | |
| Metconazole | 18 | 51 | | 26 | | 2 | 226 | | | | | 323 | |
| Metrafenone | | 224 | 12 | 263 | 13 | 115 | | | | | | 627 | |
| Penthiopyrad | 128 | 93 | | 478 | | | | | | | | 699 | |
| Penthiopyrad/picoxystrobin | 492 | 1,168 | | 49 | | 21 | 78 | | | | | 1,808 | |
| Picoxystrobin | 227 | 77 | | | | 60 | | | | | | 364 | |
| Prochloraz | | | | 351 | | | | | | | | 351 | |
| Prochloraz/tebuconazole | | | 242 | | | | | | | | | 242 | |
| Proquinazid | 61 | 81 | 125 | 144 | 130 | 206 | | | | | | 748 | |
| Prothioconazole | 6,086 | 3,779 | 412 | 1,689 | | 60 | 1,107 | | | | | 13,133 | |
| Prothioconazole/spiroxamine | 7,369 | 10,310 | 58 | 907 | 112 | | | | | | | 18,756 | |
| Prothioconazole/spiroxamine.tebuconazole | | 385 | 105 | 4,040 | | | | | | | | 4,530 | |
| Prothioconazole.tebuconazole | | 48 | 341 | 5,425 | 10 | 120 | | | | | | 5,944 | |
| Prothioconazole/trifloxystrobin | 806 | 820 | | 34 | | | | | | | | 1,659 | |
| Pyraclostrobin | 834 | 2,379 | 61 | 600 | 154 | 237 | | 12 | | | | 4,277 | |
| Sulphur | 326 | 220 | | | | | | | | | | 547 | |
| Tebuconazole | 148 | 35 | 142 | 1,017 | 294 | 678 | 320 | 214 | | | | 2,848 | |
| Thiram | | | | 80 | | | | | | | | 80 | |
| Tolclofos-methyl | | 429 | | | | | | | | | | 429 | |
| All fungicides | 123,970 | 113,059 | 18,908 | 119,862 | 10,938 | 17,096 | 3,926 | 12,534 | 2,835 | 4,756 | 58,041 | 485,925 | |

Table 6 (cont.): Estimated quantities (kilograms) of pesticide formulations used on arable crops in Ireland, 2016.

| Pesticide type & formulation | Spring barley | Winter barley | Spring wheat | Winter wheat | Spring oats | Winter oats | Oilseed rape | Peas & beans | Seed potatoes | Early potatoes | Maincrop potatoes | All crops |
|---------------------------------------------------|---------------|---------------|--------------|--------------|-------------|-------------|--------------|--------------|---------------|----------------|-------------------|-----------|
| Herbicides | | | | | | | | | | | | |
| 2,4-D | | | | | | 178 | | | | | | 178 |
| 2,4-D/triclopyr | 42 | | | | | | | | | | | 42 |
| 2,4-DB/MCPA | 1,419 | | | | 79 | 464 | | | | | | 1,963 |
| 2,4-DB/mecoprop-P | 76 | | | | | | | | | | | 76 |
| Aminopyralid/propyzamide | | | | | | | 1,206 | | | | | 1,206 |
| Asulam | | | | | 117 | | | | | | | 117 |
| Bentazone | | | | | | | | 32 | | | | 32 |
| Carfentrazone-ethyl | | | | | | | | | | 41 | | 41 |
| Clodinafop-Propargyl/cloquintocet-mexyl/pinoxaden | | | | 5 | | | | | | | | 5 |
| Clomazone/linuron | | | | | | | 2,582 | | | 215 | | 2,797 |
| Clopyralid | | | | | | | 17 | | | | | 17 |
| Clopyralid/florasulam/fluroxypyr | 7,717 | 812 | 448 | 788 | 273 | 407 | | | | | | 10,444 |
| Clopyralid/picloram | | | | | | | 26 | | | | | 26 |
| Clopyralid/triclopyr | | 21 | | 39 | | | | | | | | 60 |
| Cycloxydim | | | | | 8 | | 90 | 167 | | | | 266 |
| Dicamba/MCPA/mecoprop-P | 296 | | | | | | | | | | | 296 |
| Dicamba/mecoprop-P | 798 | | 68 | | 53 | 38 | | | | | | 957 |
| Dichlorprop-P/MCPA/mecoprop-P | 2,889 | | 186 | | 201 | | | | | | | 3,276 |
| Diflufenican | 192 | 6,313 | 18 | 2,886 | 50 | 454 | | | | 7 | | 9,919 |
| Diflufenican/flufenacet | | 878 | | 723 | | | | | | | | 1,601 |
| Diflufenican/iodosulfuron-methyl-sodium | | | | 1,412 | | | | | | | | 1,412 |
| Diquat | 458 | | | | | | 279 | 13 | 349 | 267 | 6,947 | 8,312 |
| Ethametsulfuron-methyl | | | | | | | 11 | | | | | 11 |
| Ethofumesate | 289 | | | | | | | | | | | 289 |
| Fenoxaprop-P-ethyl | 66 | 25 | 18 | 326 | | | | | | | | 435 |
| Florasulam | 3 | 13 | | 12 | | 1 | | | | | | 30 |
| Florasulam/fluroxypyr | 288 | 241 | | 37 | | 23 | | | | | | 589 |
| Florasulam/pyroxsulam | | 20 | | 69 | | | | | | | | 89 |
| Fluazifop-P-butyl | | | | | 32 | | 9 | | | | | 40 |
| Flumioxazin | | | | 1 | | | | | | | | 1 |

Table 6 (cont.): Estimated quantities (kilograms) of pesticide formulations used on arable crops in Ireland, 2016.

| Pesticide type & formulation | Spring barley | Winter barley | Spring wheat | Winter wheat | Spring oats | Crop Winter oats | Oilseed rape | Peas & beans | Seed potatoes | Early potatoes | Maincrop potatoes | All crops |
|------------------------------------------------|---------------|----------------|--------------|---------------|--------------|---------------------|---------------|---------------|---------------|----------------|-------------------|----------------|
| <i>Herbicides (cont.)</i> | | | | | | | | | | | | |
| Flupyrsulfuron-methyl | | | | | | 4 | | | | | | 4 |
| Fluroxypyr | 6,101 | 1,593 | 967 | 1,804 | 1,027 | 526 | | | | | | 12,019 |
| Glufosinate-ammonium | | | | | | | | | | 2 | 37 | 39 |
| Glyphosate | 32,230 | 29,266 | 5,296 | 18,347 | 3,180 | 4,941 | 12,688 | 1,731 | 66 | 304 | 4,291 | 112,340 |
| Imazamox/pendimethalin | | | 104 | | | | 87 | 7,808 | | | | 7,999 |
| Iodosulfuron-methyl-sodium | 24 | | | | | | | | | | | 24 |
| Iodosulfuron-methyl-sodium/mesosulfuron-methyl | | | | 179 | | | | | | | | 179 |
| Isoproturon | 3,292 | 69,585 | 725 | 31,521 | | | | | | | | 105,122 |
| Linuron | | | | | | 223 | | | 117 | 132 | 174 | 1,466 |
| MCPA | | | | | | | 698 | | | | | 223 |
| Mecoprop-P | 5,665 | 1,189 | 81 | 2,113 | 816 | | | | | | | 10,562 |
| Metazachlor | 395 | | | | 180 | | 42 | | | | | 618 |
| Metazachlor/quinmerac | | | | | | | 5,867 | | | | | 5,867 |
| Metribuzin | | | | | | | | | | 170 | 306 | 3,775 |
| Metsulfuron-methyl | 40 | 6 | 19 | | 0 | 1 | | | | | | 67 |
| Metsulfuron-methyl/thifensulfuron-methyl | 1,201 | 107 | 15 | 36 | | 1 | | 2 | | | | 1,361 |
| Metsulfuron-methyl/tribenuron-methyl | 267 | 41 | 17 | 6 | 32 | 22 | | | | | | 384 |
| Pendimethalin | 791 | | | | | | | 510 | | 123 | 1,039 | 2,462 |
| Pendimethalin/picolinafen | 330 | 4,462 | 238 | 351 | | | | | | | | 5,380 |
| Pinoxaden | 1,797 | 1,124 | 51 | 306 | | | | | | | | 3,278 |
| Propaquizafop | | | | | 78 | | 294 | 118 | | 2 | 41 | 533 |
| Propyzamide | | | | | | | 1,130 | 35 | | | | 1,166 |
| Prosulfocarb | 151 | 429 | | 1,627 | | | | 1,215 | | 391 | 6,083 | 9,895 |
| Rimsulfuron | | | | | | | | | | 0 | 4 | 5 |
| Tepraloxydim | | 10 | | 9 | | | 26 | | | | | 45 |
| Thifensulfuron-methyl/tribenuron-methyl | 1,531 | 356 | 86 | 208 | 138 | 123 | | | | | | 2,442 |
| Tribenuron-methyl | 10 | 64 | 0 | 16 | 7 | 30 | | | | | | 127 |
| Triflusulfuron-methyl | | | 6 | | | | | | | | | 6 |
| All herbicides | 68,360 | 116,563 | 8,336 | 63,194 | 6,761 | 7,269 | 21,772 | 14,328 | 716 | 1,576 | 23,940 | 332,815 |

Table 6 (cont.): Estimated quantities (kilograms) of pesticide formulations used on arable crops in Ireland, 2016.

| Pesticide type & formulation | Spring | Winter | Spring | Winter | Spring | Crop | Oilseed | Peas & | Seed | Early | Maincrop | All crops |
|----------------------------------------|--------------|--------------|--------------|---------------|------------|------------|------------|------------|------------|------------|--------------|---------------|
| | barley | barley | wheat | wheat | oats | winter | | | | | potatoes | |
| Insecticides | | | | | | | | | | | | |
| Acetamiprid | | | | | | | | | 1 | | | 1 |
| Chlorpyrifos | 1,343 | | | 76 | | | | | | | | 1,419 |
| Clofentezine | | | | | | | | 44 | | | | 44 |
| Cypermethrin | 565 | 184 | 92 | 88 | 58 | 52 | 1 | 11 | | | | 1,050 |
| Deltamethrin | 21 | 31 | 3 | 9 | 2 | 1 | | 1 | | | | 67 |
| Dimethoate | 238 | 866 | 1,358 | 9,079 | 184 | | 103 | | 90 | | 286 | 12,204 |
| Esfenvalerate | 24 | 38 | 7 | 10 | 4 | | | | | | | 83 |
| Ethoprophos | | | | | | | | 73 | | | | 73 |
| Flonicamid | | | | | | | | 37 | | | | 37 |
| Garlic Extract | 1,495 | | 660 | | | | | | | | | 2,154 |
| Lambda-cyhalothrin | 274 | 207 | 33 | 147 | 17 | 28 | 39 | 33 | 1 | 2 | 14 | 795 |
| Metarhizium anisopliae (BIPESCO 5F/52) | | 798 | | 1,714 | | | | | | | | 2,513 |
| Oxamyl | | | | | | | | | | 107 | | 107 |
| Pirimicarb | | | | | | | | 52 | | | | 52 |
| Pymetrozine | | | | | | | | | 41 | | | 41 |
| Thiacloprid | | | | | | | | | 43 | | | 43 |
| All insecticides | 3,959 | 2,125 | 2,229 | 11,047 | 264 | 81 | 143 | 141 | 286 | 108 | 300 | 20,684 |
| Molluscicides | | | | | | | | | | | | |
| Metaldehyde | | 136 | | 829 | | 46 | 248 | 44 | 7 | | 553 | 1,865 |
| Methiocarb | | 18 | | | | | 38 | 0 | 0 | | | 56 |
| All molluscicides | 154 | | 829 | | 46 | 286 | 44 | 7 | | 553 | 1,921 | |

Table 6 (cont.): Estimated quantities (kilograms) of pesticide formulations used on arable crops in Ireland, 2016.

| Pesticide type & formulation | Spring barley | Winter barley | Spring wheat | Winter wheat | Spring oats | Winter oats | Oilseed rape | Peas & beans | Seed potatoes | Early potatoes | Maincrop potatoes | All crops |
|-----------------------------------------|----------------|----------------|---------------|----------------|---------------|---------------|---------------|---------------|---------------|----------------|-------------------|------------------|
| Growth Regulators | | | | | | | | | | | | |
| Chlormequat chloride | 21,275 | 51,643 | 13,212 | 55,839 | 8,855 | 18,407 | | | | | | 169,232 |
| Chlormequat/imazaquin | 301 | | | 1,991 | | | | | | | | 2,292 |
| Ethephon (2-chloroethylphosphonic acid) | 234 | 2,049 | | 502 | | | | | | | | 2,785 |
| Ethephon/mepiquat chloride | 964 | 12,216 | 151 | 781 | | | | | | | | 14,113 |
| Mepiquat chloride/prohexadione-calcium | | | | | 18 | | | | | | | 18 |
| Trinexapac-ethyl | 764 | 2,405 | 236 | 2,349 | 260 | 562 | | | | | | 6,577 |
| All growth regulators | 23,538 | 68,315 | 13,599 | 61,463 | 9,133 | 18,969 | | | | | | 195,017 |
| Seed treatments | | | | | | | | | | | | |
| Carboxin/thiram | 5,360 | 3,001 | 885 | 2,269 | 347 | 783 | | | | | | 12,644 |
| Clothianidin/prothioconazole | 1,108 | 1,833 | | 1,251 | 112 | 116 | | | | | | 4,420 |
| Cymoxanil/fludioxonil/metalaxyl-M | | | | | | | 22 | | | | | 22 |
| Fludioxonil | 1 | | 1 | 1 | 1 | | | | | | | 5 |
| Fludioxonil/flutriafol | | 3 | | | | | | | | | | 3 |
| Fluopyram/prothioconazole/tebuconazole | 25 | 1 | | | | | | | | | | 26 |
| Flutolanil | | | | | | | | 7 | 257 | | | 264 |
| Imazalil | | | | | | | | 6 | 5 | 56 | | 67 |
| Imazalil/pencycuron | | | | | | | | 18 | 6 | 809 | | 833 |
| Imazalil/thiabendazole | | | | | | | | 1 | | | | 1 |
| Prochloraz/thiram | | | | | | 156 | | | | | | 156 |
| Prochloraz/triticonazole | 1,243 | 749 | 91 | 476 | 129 | 153 | | | | | | 2,841 |
| Silthiofam | 408 | 190 | | 219 | | | | | | | | 817 |
| All seed treatments | 8,145 | 5,777 | 977 | 4,216 | 589 | 1,052 | 156 | 22 | 25 | 17 | 1,123 | 22,100 |
| All pesticides | 227,973 | 305,992 | 44,050 | 260,613 | 27,686 | 44,513 | 26,283 | 27,070 | 3,870 | 6,457 | 83,956 | 1,058,461 |

Table 7: The active ingredients most extensively used on arable crops in Ireland in 2016, ranked by area treated (spray-hectares).

| No. | Active ingredient | Treated area (sp ha) |
|-----|---------------------------|----------------------|
| 1 | Chlorothalonil | 398,667 |
| 2 | Prothioconazole | 327,063 |
| 3 | Epoxiconazole | 222,641 |
| 4 | Chlormequat chloride | 211,256 |
| 5 | Fenpropimorph | 163,110 |
| 6 | Lambda-cyhalothrin | 159,448 |
| 7 | Pyraclostrobin | 159,092 |
| 8 | Fluroxypyr | 141,069 |
| 9 | Tribenuron-methyl | 131,308 |
| 10 | Glyphosate | 127,423 |
| 11 | Thifensulfuron-methyl | 116,866 |
| 12 | Fluxapyroxad | 116,802 |
| 13 | Pinoxaden | 115,103 |
| 14 | Trinexapac-ethyl | 106,694 |
| 15 | Diflufenican | 103,292 |
| 16 | Bixafen | 100,680 |
| 17 | Isoproturon | 100,176 |
| 18 | Metsulfuron-methyl | 81,355 |
| 19 | Florasulam | 70,302 |
| 20 | Boscalid | 68,565 |
| 21 | Tebuconazole | 65,522 |
| 22 | Spiroxamine | 64,291 |
| 23 | Metconazole | 56,811 |
| 24 | Clopyralid | 53,516 |
| 25 | Cypermethrin | 43,794 |
| 26 | Isopyrazam | 42,040 |
| 27 | Azoxystrobin | 40,721 |
| 28 | Dimethoate | 40,537 |
| 29 | Fluoxastrobin | 35,517 |
| 30 | Trifloxystrobin | 35,202 |
| 31 | Ethephon | 31,782 |
| 32 | Penthiopyrad | 31,274 |
| 33 | Cyprodinil | 30,692 |
| 34 | Fluazinam | 30,438 |
| 35 | Proquinazid | 28,726 |
| 36 | Cymoxanil | 27,477 |
| 37 | Mepiquat chloride | 27,032 |
| 38 | Iodosulphon-methyl-sodium | 26,015 |
| 39 | Propamocarb hydrochloride | 23,672 |
| 40 | Mecoprop-P | 23,654 |

Table 8: The active ingredients most extensively used on arable crops in Ireland in 2016, ranked by weight (kilograms).

| No. | Active ingredient | Quantity (kg) |
|-----|---------------------------|---------------|
| 1 | Chlorothalonil | 197,113 |
| 2 | Chlormequat chloride | 171,519 |
| 3 | Glyphosate | 112,340 |
| 4 | Isoproturon | 105,122 |
| 5 | Fenpropimorph | 48,936 |
| 6 | Prothioconazole | 40,924 |
| 7 | Mancozeb | 30,238 |
| 8 | Propamocarb hydrochloride | 19,902 |
| 9 | Fluroxypyr | 19,227 |
| 10 | Pyraclostrobin | 16,397 |
| 11 | Epoxiconazole | 15,446 |
| 12 | Spiroxamine | 15,267 |
| 13 | Pendimethalin | 15,221 |
| 14 | Boscalid | 13,916 |
| 15 | Mecoprop-P | 12,324 |
| 16 | Dimethoate | 12,204 |
| 17 | Diflufenican | 11,838 |
| 18 | Prosulfocarb | 9,895 |
| 19 | Mepiquat chloride | 9,219 |
| 20 | Fluxapyroxad | 9,176 |
| 21 | Tebuconazole | 8,957 |
| 22 | Cyprodinil | 8,823 |
| 23 | Diquat | 8,312 |
| 24 | Trinexapac-ethyl | 6,577 |
| 25 | Etephenon | 6,123 |
| 26 | Fluazinam | 5,686 |
| 27 | Azoxystrobin | 5,471 |
| 28 | Penthiopyrad | 5,221 |
| 29 | Metazachlor | 5,018 |
| 30 | Bixafen | 4,983 |
| 31 | Fenpropidin | 4,791 |
| 32 | Linuron | 4,260 |
| 33 | Metribuzin | 4,250 |
| 34 | Isopyrazam | 3,740 |
| 35 | Clopyralid | 3,738 |
| 36 | Pinoxaden | 3,280 |
| 37 | Metconazole | 3,161 |
| 38 | Cymoxanil | 3,002 |
| 39 | Mandipropamid | 2,903 |
| 40 | Folpet | 2,765 |

Table 9: Estimated quantity (kg), spray area (spha) and basic area (ha) of active substance for spring barley 2016.

| Crop | Active Substance | Quantity (kg) of | Spray area (spha) of | Basic area (ha) of |
|---------------|--------------------|------------------|----------------------|--------------------|
| | | Total | Total | Total |
| Spring Barley | Fungicides | | | |
| | Azoxystrobin | 277 | 2,375 | 2,375 |
| | Bixafen | 1,722 | 41,989 | 35,711 |
| | Boscalid | 2,841 | 16,430 | 16,430 |
| | Chlorothalonil | 55,179 | 116,017 | 95,771 |
| | Cyprodinil | 4,847 | 16,400 | 16,400 |
| | Dimethomorph | 463 | 463 | 463 |
| | Epoxiconazole | 3,946 | 67,661 | 50,186 |
| | Fenpropidin | 2,659 | 4,818 | 4,818 |
| | Fenpropimorph | 17,101 | 62,018 | 53,784 |
| | Fluxapyroxad | 1,689 | 26,850 | 24,979 |
| | Folpet | 804 | 1,608 | 1,608 |
| | Isopyrazam | 1,855 | 19,903 | 19,903 |
| | Kresoxim-methyl | 113 | 1,151 | 1,151 |
| | Mandipropamid | 73 | 586 | 586 |
| | Metconazole | 110 | 2,481 | 1,829 |
| | Metrafenone | 24 | 326 | 326 |
| | Penthiopyrad | 1,354 | 8,827 | 8,555 |
| | Picoxystrobin | 587 | 7,541 | 6,871 |
| | Proquinazid | 61 | 2,109 | 2,109 |
| | Prothioconazole | 16,129 | 133,547 | 91,513 |
| | Pyraclostrobin | 4,945 | 50,629 | 45,982 |
| | Spiroxamine | 4,806 | 24,902 | 17,766 |
| | Sulphur | 326 | 247 | 247 |
| | Tebuconazole | 148 | 594 | 594 |
| | Trifloxystrobin | 820 | 18,085 | 17,316 |
| | Herbicides | | | |
| | 2,4-D | 20 | 270 | 270 |
| | 2,4-DB | 1,286 | 892 | 892 |
| | Clopyralid | 2,726 | 37,882 | 37,104 |
| | Dicamba | 110 | 2,165 | 2,165 |
| | Dichlorprop-P | 1,456 | 4,330 | 3,509 |
| | Diflufenican | 192 | 2,202 | 2,202 |
| | Diquat | 458 | 1,145 | 1,145 |
| | Ethofumesate | 289 | 826 | 826 |
| | Fenoxaprop-P-ethyl | 66 | 1,509 | 1,509 |
| | Florasulam | 96 | 41,702 | 40,924 |
| | Fluoxastrobin | 1,090 | 17,951 | 16,387 |

Table 9 (cont.): Estimated quantity (kg), spray area (spha) and basic area (ha) of active substance for spring barley 2016.

| Crop | Active Substance | Quantity (kg) of Active Ingredient | Spray area (spha) of Active Ingredient | Basic area (ha) of Active Ingredient |
|---------------|----------------------------------------|------------------------------------|----------------------------------------|--------------------------------------|
| | | Total | Total | Total |
| Spring Barley | Herbicides (cont.) | | | |
| | Fluroxypyr | 11,288 | 83,068 | 81,194 |
| | Glyphosate | 32,230 | 36,952 | 32,924 |
| | Imazaquin | 1 | 326 | 326 |
| | Iodosulfuron-methyl-sodium | 24 | 1,636 | 1,636 |
| | Isoproturon | 3,292 | 4,178 | 4,178 |
| | MCPA | 1,101 | 5,777 | 4,956 |
| | Mecoprop-P | 7,192 | 15,989 | 15,168 |
| | Metazachlor | 395 | 791 | 791 |
| | Metsulfuron-methyl | 245 | 59,159 | 56,164 |
| | Pendimethalin | 1,114 | 1,117 | 1,117 |
| | Picolinafen | 7 | 326 | 326 |
| | Pinoxaden | 1,797 | 62,578 | 60,799 |
| | Prosulfocarb | 151 | 94 | 94 |
| | Thifensulfuron-methyl | 2,066 | 75,657 | 73,859 |
| | Tribenuron-methyl | 738 | 73,950 | 73,479 |
| | Triclopyr | 22 | 270 | 270 |
| | Insecticides | | | |
| | Chlorpyrifos | 1,343 | 1,796 | 1,796 |
| | Cypermethrin | 565 | 23,443 | 21,839 |
| | Deltamethrin | 21 | 3,526 | 3,526 |
| | Dimethoate | 238 | 708 | 708 |
| | Esfenvalerate | 24 | 5,587 | 4,909 |
| | Garlic Extract | 1,495 | 2,793 | 2,793 |
| | Lambda-cyhalothrin | 274 | 56,757 | 55,142 |
| | Growth Regulators | | | |
| | Chlormequat chloride | 21,575 | 29,810 | 29,459 |
| | Ethephon | 569 | 2,587 | 2,280 |
| | Mepiquat chloride | 629 | 1,795 | 1,795 |
| | Trinexapac-ethyl | 764 | 12,589 | 12,396 |
| | Seed Treatments | | | |
| | Carboxin/thiram | 5,360 | 28,313 | 28,313 |
| | Clothianidin/prothioconazol | 1,108 | 12,348 | 12,348 |
| | Fludioxonil | 1 | 183 | 183 |
| | Fluopyram/prothioconazole/tebuconazole | 25 | 1,658 | 1,658 |
| | Prochloraz/triticonazole | 1,243 | 51,576 | 51,576 |
| | Silthiofam | 408 | 11,260 | 11,260 |

Table 10: Estimated quantity (kg), spray area (spha) and basic area (ha) of active substance for winter barley 2016.

| Crop | Active Substance | Quantity (kg) of Active Ingredient | Spray area (spha) of Active Ingredient | Basic area (ha) of Active Ingredient |
|---------------|--------------------|------------------------------------|----------------------------------------|--------------------------------------|
| | | Total | Total | Total |
| Winter barley | Fungicides | | | |
| | Azoxystrobin | 697 | 5,187 | 4,315 |
| | Bixafen | 1,321 | 32,629 | 29,255 |
| | Boscalid | 3,534 | 18,315 | 15,948 |
| | Chlorothalonil | 53,779 | 110,466 | 66,954 |
| | Cyproconazole | 13 | 132 | 132 |
| | Cyprodinil | 3,727 | 13,368 | 13,368 |
| | Difenoconazole | 16 | 212 | 212 |
| | Epoxiconazole | 3,473 | 54,228 | 39,074 |
| | Fenpropidin | 255 | 341 | 341 |
| | Fenpropimorph | 8,661 | 31,892 | 25,381 |
| | Fluopicolide | 116 | 620 | 620 |
| | Fluoxastrobin | 1,065 | 16,214 | 13,342 |
| | Fluxapyroxad | 2,051 | 30,990 | 24,864 |
| | Folpet | 1,112 | 1,678 | 1,678 |
| | Isopyrazam | 1,720 | 20,613 | 20,613 |
| | Kresoxim-methyl | 220 | 1,854 | 1,291 |
| | Metconazole | 107 | 2,162 | 2,162 |
| | Metrafenone | 224 | 1,588 | 1,588 |
| | Penthiopyrad | 1,759 | 10,852 | 10,273 |
| | Picoxystrobin | 703 | 7,514 | 7,514 |
| | Proquinazid | 107 | 3,433 | 3,063 |
| | Prothioconazole | 13,684 | 107,877 | 64,914 |
| | Pyraclostrobin | 6,097 | 55,843 | 40,365 |
| | Spiroxamine | 6,941 | 27,468 | 19,983 |
| | Sulphur | 220 | 276 | 276 |
| | Tebuconazole | 143 | 1,054 | 1,054 |
| | Tolclofos-methyl | 429 | 1,128 | 1,128 |
| | Trifloxystrobin | 885 | 16,286 | 13,804 |
| | Herbicides | | | |
| | Clopyralid | 291 | 4,829 | 4,829 |
| | Diflufenican | 6,606 | 56,286 | 53,920 |
| | Fenoxaprop-P-ethyl | 25 | 441 | 441 |
| | Florasulam | 36 | 11,437 | 10,485 |
| | Flufenacet | 585 | 4,949 | 4,949 |
| | Fluroxypyr | 2,339 | 18,116 | 17,408 |
| | Glyphosate | 29,266 | 34,212 | 27,063 |
| | Isoproturon | 69,585 | 63,466 | 61,593 |

Table 10 (cont.): Estimated quantity (kg), spray area (spha) and basic area (ha) of active substance for winter barley 2016.

| Crop | Active Substance | Quantity (kg) of | Spray area (spha) of | Basic area (ha) of |
|---------------|--------------------------------------------|----------------------------|----------------------------|----------------------------|
| | | Active Ingredient Total | Active Ingredient Total | Active Ingredient Total |
| Winter barley | Herbicides (cont.) | | | |
| | Mecoprop-P | 1,189 | 1,445 | 1,445 |
| | Metsulfuron-methyl | 34 | 8,261 | 8,079 |
| | Pendimethalin | 4,363 | 4,210 | 4,210 |
| | Picolinafen | 99 | 4,210 | 4,210 |
| | Pinoxaden | 1,124 | 41,473 | 40,910 |
| | Prosulfocarb | 429 | 241 | 241 |
| | Pyroxsulam | 17 | 952 | 952 |
| | Tepraloxo.dim | 10 | 198 | 198 |
| | Thifensulfuron-methyl | 315 | 14,768 | 14,768 |
| | Tribenuron-methyl | 225 | 20,727 | 19,213 |
| | Triclopyr | 17 | 238 | 238 |
| | Triflusulfuron-methyl | 6 | 418 | 418 |
| | Insecticides | | | |
| | Cypermethrin | 184 | 8,323 | 7,475 |
| | Deltamethrin | 31 | 3,730 | 3,730 |
| | Dimethoate | 866 | 2,453 | 2,453 |
| | Esfenvalerate | 38 | 9,301 | 8,253 |
| | Lambda-cyhalothrin | 207 | 43,796 | 40,698 |
| | Metarhizium anisopliae (BIPESCO 5F/52) | 798 | 327 | 327 |
| | Molluscicides | | | |
| | Metaldehyde | 136 | 649 | 649 |
| | Methiocarb | 18 | 442 | 442 |
| | Growth Regulators | | | |
| | Chlormequat chloride | 51,643 | 63,155 | 57,798 |
| | Ethephon | 6,299 | 29,787 | 29,219 |
| | Mepiquat chloride | 7,967 | 23,367 | 22,800 |
| | Trinexapac-ethyl | 2,405 | 41,070 | 35,522 |
| | Seed Treatments | | | |
| | Carboxin/thiram | 3,001 | 16,237 | 16,237 |
| | Clothianidin/prothioconazole | 1,833 | 20,429 | 20,429 |
| | Fludioxonil/flutriafol | 3 | 166 | 166 |
| | Fluopyram/prothioconazole/te buconazole | 1 | 86 | 86 |
| | Prochloraz/triticonazole | 749 | 29,010 | 29,010 |
| | Silthiofam | 190 | 4,919 | 4,919 |

Table 11: Estimated quantity (kg), spray area (spha) and basic area (ha) of active substance for spring wheat 2016.

| Crop | Active Substance | Quantity (kg) of Active Ingredient Total | Spray area (spha) of Active Ingredient Total | Basic area (ha) of Active Ingredient Total |
|--------------|------------------|------------------------------------------|----------------------------------------------|--------------------------------------------|
| Spring wheat | Fungicides | | | |
| | Azoxystrobin | 193 | 2,011 | 2,011 |
| | Bixafen | 256 | 3,392 | 3,306 |
| | Boscalid | 959 | 4,093 | 4,093 |
| | Chlorothalonil | 10,927 | 22,010 | 10,396 |
| | Cyprodinil | 227 | 806 | 806 |
| | Epoxiconazole | 867 | 10,442 | 8,335 |
| | Fenpropidin | 880 | 1,543 | 1,543 |
| | Fenpropimorph | 1,117 | 3,375 | 2,377 |
| | Fluoxastrobin | 36 | 290 | 290 |
| | Fluxapyroxad | 98 | 1,218 | 1,218 |
| | Folpet | 440 | 880 | 880 |
| | Isopyrazam | 76 | 806 | 806 |
| | Metconazole | 41 | 1,270 | 1,270 |
| | Metrafenone | 82 | 882 | 882 |
| | Penthiopyrad | 192 | 1,168 | 1,168 |
| | Picoxystrobin | 58 | 290 | 290 |
| | Prochloraz | 145 | 880 | 880 |
| | Propiconazole | 35 | 553 | 553 |
| | Proquinazid | 165 | 4,548 | 4,548 |
| | Prothioconazole | 1,183 | 7,315 | 6,549 |
| | Pyraclostrobin | 216 | 1,688 | 1,645 |
| | Spiroxamine | 97 | 362 | 362 |
| | Tebuconazole | 618 | 5,195 | 5,195 |

Table 11 (cont.): Estimated quantity (kg), spray area (spha) and basic area (ha) of active substance for spring wheat 2016.

| Crop | Active Substance | Quantity (kg) of | Spray area (spha) of | Basic area (ha) of |
|--------------|--------------------------|----------------------------|----------------------------|----------------------------|
| | | Active Ingredient Total | Active Ingredient Total | Active Ingredient Total |
| Spring wheat | Herbicides | | | |
| | Clopyralid | 158 | 2,215 | 2,215 |
| | Dicamba | 8 | 303 | 303 |
| | Dichlorprop-P | 92 | 295 | 295 |
| | Diflufenican | 18 | 140 | 140 |
| | Fenoxaprop-P-ethyl | 18 | 457 | 457 |
| | Florasulam | 5 | 2,215 | 2,215 |
| | Fluroxypyr | 1,252 | 8,868 | 8,661 |
| | Glyphosate | 5,296 | 6,282 | 6,143 |
| | Imazamox | 7 | 130 | 130 |
| | Isoproturon | 725 | 932 | 932 |
| | MCPA | 47 | 295 | 295 |
| | Mecoprop-P | 188 | 748 | 748 |
| | Metsulfuron-methyl | 28 | 6,385 | 6,385 |
| | Pendimethalin | 330 | 693 | 693 |
| | Picolinafen | 5 | 563 | 563 |
| | Pinoxaden | 51 | 1,695 | 1,695 |
| | Thifensulfuron-methyl | 59 | 3,992 | 3,846 |
| | Tribenuron-methyl | 49 | 5,626 | 5,412 |
| | Insecticides | | | |
| | Chlorpyrifos | 76 | 382 | 382 |
| | Cypermethrin | 92 | 3,828 | 3,123 |
| | Deltamethrin | 3 | 560 | 560 |
| | Dimethoate | 1,358 | 4,508 | 4,508 |
| | Esfenvalerate | 7 | 2,245 | 2,245 |
| | Garlic Extract | 660 | 880 | 880 |
| | Lambda-cyhalothrin | 33 | 4,814 | 4,264 |
| | Growth Regulators | | | |
| | Chlormequat chloride | 13,212 | 15,510 | 8,972 |
| | Ethephon | 52 | 233 | 233 |
| | Mepiquat chloride | 98 | 233 | 233 |
| | Trinexapac-ethyl | 236 | 3,534 | 3,534 |
| | Seed Treatments | | | |
| | Carboxin/thiram | 885 | 4,708 | 4,708 |
| | Fludioxonil | 1 | 161 | 161 |
| | Prochloraz/triticonazole | 91 | 3,624 | 3,624 |

Table 12: Estimated quantity (kg), spray area (spha) and basic area (ha) of active substance for winter wheat 2016.

| Crop | Active Substance | Quantity (kg) of | Spray area (spha) of | Basic area (ha) of |
|--------------|----------------------|----------------------------|----------------------------|----------------------------|
| | | Active Ingredient Total | Active Ingredient Total | Active Ingredient Total |
| Winter wheat | Fungicides | | | |
| | Azoxystrobin | 2,345 | 18,397 | 18,397 |
| | Bixafen | 1,625 | 21,403 | 19,721 |
| | Boscalid | 3,129 | 11,997 | 11,442 |
| | Chlorothalonil | 70,189 | 138,714 | 50,549 |
| | Cyproconazole | 13 | 171 | 171 |
| | Cyprodinil | 22 | 118 | 118 |
| | Difenoconazole | 321 | 3,759 | 3,759 |
| | Epoxiconazole | 5,227 | 63,108 | 37,974 |
| | Fenpropidin | 409 | 1,073 | 1,073 |
| | Fenpropimorph | 4,790 | 17,249 | 13,574 |
| | Fluoxastrobin | 58 | 754 | 754 |
| | Flusilazole | 81 | 564 | 282 |
| | Fluxapyroxad | 5,100 | 53,782 | 38,390 |
| | Folpet | 410 | 710 | 710 |
| | Iprodione | 188 | 626 | 626 |
| | Isopyrazam | 90 | 719 | 601 |
| | Kresoxim-methyl | 132 | 1,299 | 1,299 |
| | Mancozeb | 181 | 161 | 161 |
| | Metconazole | 2,647 | 45,108 | 36,345 |
| | Metrafenone | 701 | 6,134 | 4,923 |
| | Penthiopyrad | 1,850 | 9,940 | 9,123 |
| | Picoxystrobin | 194 | 1,822 | 1,615 |
| | Prochloraz | 351 | 896 | 896 |
| | Proquinazid | 261 | 9,170 | 8,625 |
| | Prothioconazole | 8,516 | 66,201 | 45,072 |
| | Pyraclostrobin | 1,147 | 11,395 | 11,214 |
| | Spiroxamine | 3,349 | 11,171 | 10,854 |
| | Tebuconazole | 6,422 | 48,860 | 37,254 |
| | Thiram | 80 | 201 | 201 |
| | Trifloxystrobin | 34 | 831 | 831 |
| | Herbicides | | | |
| | 2,4-D | 178 | 712 | 712 |
| | 2,4-DB | 68 | 282 | 282 |
| | Asulam | 117 | 146 | 146 |
| | Clodinafop-Propargyl | 2 | 176 | 176 |
| | Clopyralid | 286 | 3,867 | 3,867 |
| | Cloquintocet-mexyl | 0 | 176 | 176 |
| | Diflufenican | 4,512 | 40,057 | 33,543 |
| | Fenoxaprop-P-ethyl | 326 | 5,917 | 4,253 |
| | Florasulam | 34 | 9,915 | 9,699 |
| | Flufenacet | 482 | 4,029 | 4,029 |

Table 12 (cont.): Estimated quantity (kg), spray area (spha) and basic area (ha) of active substance for winter wheat 2016.

| Crop | Active Substance | Quantity (kg) of | Spray area (spha) of | Basic area (ha) of |
|--------------|-------------------------------------------|----------------------------|----------------------------|----------------------------|
| | | Active Ingredient Total | Active Ingredient Total | Active Ingredient Total |
| Winter wheat | Herbicides (cont.) | | | |
| | Flumioxazin | 1 | 33 | 33 |
| | Fluroxypyr | 2,340 | 15,676 | 14,789 |
| | Glyphosate | 18,347 | 23,672 | 20,291 |
| | Imazaquin | 4 | 2,826 | 2,826 |
| | Iodosulfuron-methyl-sodium | 72 | 24,379 | 23,561 |
| | Isoproturon | 31,521 | 31,600 | 28,684 |
| | MCPA | 11 | 282 | 282 |
| | Mecoprop-P | 2,113 | 3,134 | 3,134 |
| | Mesosulfuron-methyl | 134 | 13,664 | 12,846 |
| | Metsulfuron-methyl | 6 | 2,179 | 2,179 |
| | Pendimethalin | 343 | 346 | 346 |
| | Picolinafen | 8 | 346 | 346 |
| | Pinoxaden | 309 | 9,357 | 9,357 |
| | Prosulfocarb | 1,627 | 1,017 | 1,017 |
| | Pyroxasulam | 57 | 3,367 | 3,367 |
| | Tepraloxydim | 9 | 185 | 185 |
| | Thifensulfuron-methyl | 151 | 10,297 | 10,297 |
| | Tribenuron-methyl | 109 | 11,524 | 10,313 |
| | Triclopyr | 31 | 429 | 429 |
| | Insecticides | | | |
| | Cypermethrin | 88 | 4,029 | 4,029 |
| | Deltamethrin | 9 | 1,702 | 1,702 |
| | Dimethoate | 9,079 | 30,791 | 30,171 |
| | Esfenvalerate | 10 | 2,609 | 2,609 |
| | Lambda-cyhalothrin | 147 | 31,154 | 26,459 |
| | Metarhizium anisopliae (BIPESCO 5F/52) | 1,714 | 703 | 703 |
| | Molluscicides | | | |
| | Metaldehyde | 829 | 3,930 | 3,930 |
| | Growth Regulators | | | |
| | Chlormequat chloride | 57,826 | 68,018 | 44,843 |
| | Ethephon | 774 | 4,167 | 4,167 |
| | Mepiquat chloride | 510 | 1,509 | 1,509 |
| | Trinexapac-ethyl | 2,349 | 33,162 | 26,620 |
| | Seed Treatments | | | |
| | Carboxin/thiram | 2,269 | 13,217 | 13,217 |
| | Clothianidin/prothioconazole | 1,251 | 13,022 | 13,022 |
| | Fludioxonil | 1 | 161 | 161 |
| | Prochloraz/triticonazole | 476 | 18,964 | 18,964 |
| | Silthiofam | 219 | 5,496 | 5,496 |

Table 13: Estimated quantity (kg), spray area (spha) and basic area (ha) of active substance for spring oats 2016.

| Crop | Active Substance | Quantity (kg) of | Spray area (spha) of | Basic area (ha) of |
|-------------|-------------------|----------------------------|----------------------------|----------------------------|
| | | Active Ingredient Total | Active Ingredient Total | Active Ingredient Total |
| Spring Oats | Fungicides | | | |
| | Azoxystrobin | 66 | 631 | 631 |
| | Bixafen | 13 | 437 | 437 |
| | Boscalid | 61 | 525 | 525 |
| | Chlorothalonil | 351 | 707 | 689 |
| | Cymoxanil | 8 | 78 | 78 |
| | Cyproconazole | 4 | 71 | 71 |
| | Difenoconazole | 26 | 441 | 441 |
| | Dimethomorph | 397 | 397 | 397 |
| | Epoxiconazole | 685 | 10,860 | 6,920 |
| | Fenpropidin | 463 | 1,215 | 1,134 |
| | Fenpropimorph | 6,240 | 18,337 | 9,167 |
| | Fluoxastrobin | 5 | 50 | 50 |
| | Fluxapyroxad | 157 | 2,591 | 2,591 |
| | Metconazole | 21 | 292 | 292 |
| | Metrafenone | 363 | 4,228 | 4,051 |
| | Picoxystrobin | 10 | 71 | 71 |
| | Proquinazid | 130 | 3,607 | 2,645 |
| | Prothioconazole | 93 | 976 | 782 |
| | Pyraclostrobin | 1,473 | 12,469 | 8,028 |
| | Spiroxamine | 73 | 388 | 244 |
| | Tebuconazole | 300 | 1,976 | 1,956 |

Table 13 (cont.): Estimated quantity (kg), spray area (spha) and basic area (ha) of active substance for spring oats 2016.

| Crop | Active Substance | Quantity (kg) of | Spray area (spha) of | Basic area (ha) of |
|-------------|------------------------------|----------------------------|----------------------------|----------------------------|
| | | Active Ingredient Total | Active Ingredient Total | Active Ingredient Total |
| Spring Oats | Herbicides | | | |
| | 2,4-DB | 398 | 237 | 237 |
| | Clopyralid | 96 | 1,454 | 1,454 |
| | Cycloxydim | 8 | 260 | 260 |
| | Dicamba | 6 | 88 | 88 |
| | Dichlorprop-P | 101 | 277 | 176 |
| | Diflufenican | 50 | 397 | 397 |
| | Florasulam | 3 | 1,454 | 1,454 |
| | Fluazifop-P-butyl | 32 | 260 | 260 |
| | Fluroxypyr | 1,200 | 8,448 | 7,060 |
| | Glyphosate | 3,180 | 3,583 | 3,214 |
| | MCPA | 341 | 842 | 741 |
| | Mecoprop-P | 911 | 1,430 | 1,330 |
| | Metazachlor | 180 | 360 | 360 |
| | Metsulfuron-methyl | 14 | 3,276 | 3,276 |
| | Propaquizafop | 78 | 260 | 260 |
| | Thifensulfuron-methyl | 72 | 6,441 | 6,441 |
| | Tribenuron-methyl | 91 | 10,091 | 10,091 |
| | Insecticides | | | |
| | Cypermethrin | 58 | 2,186 | 2,186 |
| | Deltamethrin | 2 | 306 | 306 |
| | Dimethoate | 184 | 585 | 585 |
| | Esfenvalerate | 4 | 836 | 836 |
| | Lambda-cyhalothrin | 17 | 3,433 | 3,068 |
| | Growth Regulators | | | |
| | Chlormequat chloride | 8,855 | 11,258 | 8,756 |
| | Mepiquat chloride | 15 | 127 | 127 |
| | Prohexadione-calcium | 3 | 127 | 127 |
| | Trinexapac-ethyl | 260 | 5,239 | 4,714 |
| | Seed Treatments | | | |
| | Carboxin/thiram | 347 | 1,908 | 1,908 |
| | Clothianidin/prothioconazole | 112 | 1,242 | 1,242 |
| | Fludioxonil | 1 | 162 | 162 |
| | Prochloraz/triticonazole | 129 | 5,155 | 5,155 |

Table 14: Estimated quantity (kg), spray area (spha) and basic area (ha) of active substance for winter oats 2016.

| Crop | Active Substance | Quantity (kg) of | Spray area (spha) of | Basic area (ha) of |
|-------------|-------------------|----------------------------|----------------------------|----------------------------|
| | | Active Ingredient Total | Active Ingredient Total | Active Ingredient Total |
| Winter oats | Fungicides | | | |
| | Azoxystrobin | 245 | 1,919 | 1,710 |
| | Boscalid | 82 | 403 | 403 |
| | Chlorothalonil | 148 | 297 | 297 |
| | Cyproconazole | 5 | 86 | 86 |
| | Epoxiconazole | 1,228 | 16,159 | 10,003 |
| | Fenpropidin | 125 | 547 | 547 |
| | Fenpropimorph | 11,027 | 30,239 | 12,342 |
| | Fluoxastrobin | 27 | 258 | 258 |
| | Fluxapyroxad | 73 | 1,282 | 985 |
| | Kresoxim-methyl | 234 | 2,205 | 1,276 |
| | Metconazole | 9 | 409 | 409 |
| | Metrafenone | 900 | 9,696 | 7,141 |
| | Penthiopyrad | 14 | 86 | 86 |
| | Picoxystrobin | 78 | 650 | 564 |
| | Proquinazid | 206 | 5,860 | 5,377 |
| | Prothioconazole | 87 | 735 | 735 |
| | Pyraclostrobin | 1,930 | 14,527 | 10,466 |
| | Tebuconazole | 678 | 3,410 | 3,114 |

Table 14 (cont.): Estimated quantity (kg), spray area (spha) and basic area (ha) of active substance for winter oats 2016.

| Crop | Active Substance | Quantity (kg) of Active | Spray area (spha) of Active | Basic area (ha) of Active |
|-------------|------------------------------|-------------------------|-----------------------------|---------------------------|
| | | Ingredient | Ingredient | Ingredient |
| | | Total | Total | Total |
| Winter oats | Herbicides | | | |
| | Clopyralid | 144 | 2,814 | 2,126 |
| | Dicamba | 4 | 93 | 93 |
| | Diflufenican | 454 | 4,175 | 3,174 |
| | Florasulam | 7 | 3,578 | 2,890 |
| | Flupyrifluorfen-methyl | 4 | 157 | 157 |
| | Fluroxypyr | 807 | 6,893 | 5,708 |
| | Glyphosate | 4,941 | 5,423 | 5,256 |
| | Mecoprop-P | 731 | 907 | 907 |
| | Metsulfuron-methyl | 12 | 2,031 | 2,031 |
| | Thifensulfuron-methyl | 62 | 5,648 | 5,648 |
| | Tribenuron-methyl | 103 | 9,389 | 9,389 |
| | Insecticides | | | |
| | Cypermethrin | 52 | 1,619 | 1,619 |
| | Deltamethrin | 1 | 93 | 93 |
| | Lambda-cyhalothrin | 28 | 5,737 | 4,892 |
| | Growth Regulators | | | |
| | Chlormequat chloride | 18,407 | 23,506 | 16,908 |
| | Trinexapac-ethyl | 562 | 11,099 | 8,355 |
| | Molluscicides | | | |
| | Metaldehyde | 46 | 231 | 231 |
| | Seed Treatments | | | |
| | Carboxin/thiram | 783 | 4,515 | 4,515 |
| | Clothianidin/prothioconazole | 116 | 1,254 | 1,254 |
| | Prochloraz/triticonazole | 153 | 6,330 | 6,330 |

Table 15: Estimated quantity (kg), spray area (spha) and basic area (ha) of active substance for oilseed rape 2016.

| Crop | Active Substance | Quantity (kg) of | Spray area (spha) of | Basic area (ha) of |
|--------------|------------------------|----------------------------|----------------------------|----------------------------|
| | | Active Ingredient Total | Active Ingredient Total | Active Ingredient Total |
| Oilseed rape | Fungicides | | | |
| | Azoxystrobin | 804 | 5,284 | 3,732 |
| | Bixafen | 41 | 714 | 714 |
| | Boscalid | 1,012 | 4,362 | 4,362 |
| | Chlorothalonil | 96 | 179 | 179 |
| | Epoxiconazole | 12 | 93 | 93 |
| | Metconazole | 226 | 5,088 | 4,578 |
| | Penthiopyrad | 52 | 401 | 401 |
| | Picoxystrobin | 26 | 401 | 401 |
| | Prothioconazole | 1,222 | 10,294 | 7,026 |
| | Tebuconazole | 435 | 3,252 | 2,539 |
| | Herbicides | | | |
| | Aminopyralid | 15 | 1,433 | 1,433 |
| | Clopyralid | 38 | 455 | 455 |
| | Cycloxydim | 90 | 600 | 600 |
| | Diquat | 279 | 745 | 745 |
| | Ethametsulfuron-methyl | 11 | 618 | 618 |
| | Fluazifop-P-butyl | 9 | 140 | 140 |
| | Glyphosate | 12,688 | 10,417 | 8,306 |
| | Imazamox | 5 | 93 | 93 |
| | Metazachlor | 4,443 | 6,278 | 5,868 |
| | Pendimethalin | 82 | 93 | 93 |
| | Picloram | 5 | 263 | 263 |
| | Propaquizafop | 294 | 4,768 | 4,662 |
| | Propyzamide | 2,322 | 3,245 | 2,934 |
| | Quinmerac | 1,467 | 6,222 | 5,811 |
| | Tepraloxydim | 26 | 681 | 681 |
| | Insecticides | | | |
| | Cypermethrin | 1 | 23 | 23 |
| | Dimethoate | 103 | 323 | 323 |
| | Lambda-cyhalothrin | 39 | 6,082 | 4,261 |
| | Molluscicides | | | |
| | Metaldehyde | 248 | 1,186 | 1,139 |
| | Methiocarb | 38 | 392 | 286 |
| | Seed Treatments | | | |
| | Prochloraz/thiram | 156 | 8,960 | 8,960 |

Table 16: Estimated quantity (kg), spray area (spha) and basic area (ha) of active substance for peas and beans 2016.

| Crop | Active Substance | Quantity (kg) of | Spray area (spha) of | Basic area (ha) of |
|--------------|------------------------------|----------------------------|----------------------------|----------------------------|
| | | Active Ingredient Total | Active Ingredient Total | Active Ingredient Total |
| Peas & beans | Fungicides | | | |
| | Azoxystrobin | 577 | 3,816 | 3,144 |
| | Bixafen | 3 | 117 | 117 |
| | Boscalid | 2,299 | 12,439 | 8,479 |
| | Chlorothalonil | 6,379 | 10,115 | 6,579 |
| | Difenoconazole | 52 | 416 | 416 |
| | Dimethomorph | 4 | 63 | 63 |
| | Mancozeb | 568 | 398 | 398 |
| | Metalaxy-M | 1,839 | 2,682 | 2,631 |
| | Prothioconazole | 12 | 117 | 117 |
| | Pyraclostrobin | 589 | 12,540 | 8,479 |
| | Tebuconazole | 214 | 1,180 | 935 |
| | Herbicides | | | |
| | Bentazone | 32 | 88 | 88 |
| | Clomazone | 394 | 6,305 | 6,305 |
| | Cycloxydim | 167 | 1,167 | 1,167 |
| | Diquat | 13 | 24 | 24 |
| | Glyphosate | 1,731 | 1,935 | 1,535 |
| | Imazamox | 489 | 9,487 | 9,487 |
| | Linuron | 2,305 | 6,506 | 6,506 |
| | Metsulfuron-methyl | 0.314 | 63 | 63 |
| | Pendimethalin | 7,828 | 10,086 | 10,086 |
| | Propaquizafop | 118 | 1,257 | 1,257 |
| | Propyzamide | 35 | 44 | 44 |
| | Prosulfocarb | 1,215 | 419 | 419 |
| | Thifensulfuron-methyl | 2 | 63 | 63 |
| | Insecticides | | | |
| | Clofentezine | 44 | 221 | 221 |
| | Cypermethrin | 11 | 343 | 343 |
| | Deltamethrin | 1 | 128 | 128 |
| | Esfenvalerate | 0.235 | 63 | 63 |
| | Lambda-cyhalothrin | 33 | 4,913 | 4,913 |
| | Pirimicarb | 52 | 378 | 378 |
| | Molluscicides | | | |
| | Metaldehyde | 44 | 222 | 222 |
| | Seed Treatments | | | |
| | Cymoxanil/fludioxonil/metala | 22 | 223 | 223 |

Table 17: Estimated quantity (kg), spray area (spha) and basic area (ha) of active substance for seed potatoes 2016.

| Crop | Active Substance | Quantity (kg) of | Spray area (spha) of | Basic area (ha) of |
|---------------|---------------------------|----------------------------|----------------------------|----------------------------|
| | | Active Ingredient Total | Active Ingredient Total | Active Ingredient Total |
| Seed potatoes | Fungicides | | | |
| | Benthiavalicarb-isopropyl | 0.394 | 14 | 14 |
| | Cyazofamid | 26 | 244 | 237 |
| | Cymoxanil | 167 | 1,735 | 324 |
| | Fenamidone | 6 | 43 | 14 |
| | Fluazinam | 264 | 1,445 | 324 |
| | Fluopicolide | 79 | 793 | 263 |
| | Mancozeb | 1,243 | 950 | 310 |
| | Mandipropamid | 138 | 991 | 299 |
| | Metalaxyl-M | 40 | 34 | 34 |
| | Propamocarb hydrochloride | 870 | 892 | 297 |
| | Herbicides | | | |
| | Diquat | 349 | 1,135 | 329 |
| | Glyphosate | 66 | 72 | 53 |
| | Linuron | 132 | 256 | 256 |
| | Metribuzin | 170 | 308 | 308 |
| | Insecticides | | | |
| | Acetamiprid | 1 | 14 | 14 |
| | Dimethoate | 90 | 268 | 249 |
| | Ethoprophos | 73 | 19 | 19 |
| | Flonicamid | 37 | 467 | 237 |
| | Lambda-cyhalothrin | 1 | 80 | 47 |
| | Pymetrozine | 41 | 272 | 235 |
| | Thiacloprid | 43 | 450 | 235 |
| | Molluscicides | | | |
| | Metaldehyde | 7 | 40 | 40 |
| | Methiocarb | 0.437 | 7 | 7 |
| | Seed Treatments | | | |
| | Imazalil | 6 | 213 | 213 |
| | Imazalil/pencycuron | 18 | 41 | 41 |
| | Imazalil/thiabendazole | 1 | 11 | 11 |

Table 18: Estimated quantity (kg), spray area (spha) and basic area (ha) of active substance for early potatoes 2016.

| Crop | Active Substance | Quantity (kg) of Active Ingredient | Spray area (spha) of Active Ingredient | Basic area (ha) of Active Ingredient |
|----------------|---------------------------|---------------------------------------|-------------------------------------------|-----------------------------------------|
| | | Total | Total | Total |
| Early Potatoes | Fungicides | | | |
| | Azoxystrobin | 9 | 36 | 36 |
| | Benthiavalicarb-isopropyl | 8 | 288 | 185 |
| | Cyazofamid | 24 | 181 | 151 |
| | Cymoxanil | 187 | 1,756 | 746 |
| | Fenamidone | 34 | 244 | 100 |
| | Fluazinam | 220 | 1,299 | 439 |
| | Fluopicolide | 97 | 999 | 556 |
| | Mancozeb | 2,622 | 1,805 | 784 |
| | Mandipropamid | 81 | 534 | 354 |
| | Metalaxy-M | 24 | 21 | 16 |
| | Propamocarb hydrochloride | 1,450 | 1,561 | 799 |
| | Herbicides | | | |
| | Diflufenican | 7 | 36 | 36 |
| | Diquat | 267 | 635 | 431 |
| | Glufosinate-ammonium | 2 | 11 | 11 |
| | Glyphosate | 304 | 313 | 313 |
| | Linuron | 174 | 386 | 386 |
| | Metribuzin | 306 | 673 | 673 |
| | Pendimethalin | 123 | 93 | 93 |
| | Propaquizafop | 2 | 17 | 17 |
| | Prosulfocarb | 391 | 127 | 127 |
| | Rimsulfuron | 0.444 | 36 | 36 |
| | Insecticides | | | |
| | Lambda-cyhalothrin | 2 | 253 | 149 |
| | Oxamyl | 107 | 36 | 36 |
| | Seed Treatments | | | |
| | Flutolanil | 7 | 31 | 31 |
| | Imazalil | 5 | 175 | 175 |
| | Imazalil/pencycuron | 6 | 17 | 17 |

Table 19: Estimated quantity (kg), spray area (spha) and basic area (ha) of active substance for maincrop potatoes 2016.

| Crop | Active Substance | Quantity (kg) of | Spray area (spha) of | Basic area (ha) of |
|----------------------|---------------------------|------------------|----------------------|--------------------|
| | | Total | Total | Total |
| Maincrop potatoes | Fungicides | | | |
| | Azoxystrobin | 259 | 1,065 | 904 |
| | Benthiavalicarb-isopropyl | 61 | 2,752 | 1,547 |
| | Chlorothalonil | 64 | 161 | 161 |
| | Cyazofamid | 1,936 | 11,412 | 4,506 |
| | Cymoxanil | 2,640 | 23,908 | 6,663 |
| | Epoxiconazole | 8 | 90 | 90 |
| | Fenamidone | 314 | 2,157 | 833 |
| | Fluazinam | 5,202 | 27,694 | 7,231 |
| | Fluopicolide | 1,471 | 17,536 | 7,179 |
| | Fluxapyroxad | 8 | 90 | 90 |
| | Mancozeb | 25,624 | 17,790 | 6,021 |
| | Mandipropamid | 2,612 | 18,101 | 5,570 |
| | Metalaxyl-M | 237 | 212 | 202 |
| | Propamocarb hydrochloride | 17,582 | 21,219 | 7,391 |
| | Zoxamide | 24 | 161 | 161 |
| | Herbicides | | | |
| | Carfentrazone-ethyl | 41 | 1,522 | 1,522 |
| | Clomazone | 33 | 487 | 487 |
| | Diquat | 6,947 | 18,309 | 7,585 |
| | Glufosinate-ammonium | 37 | 183 | 183 |
| | Glyphosate | 4,291 | 4,562 | 4,562 |
| | Linuron | 1,649 | 2,641 | 2,641 |
| | Metribuzin | 3,775 | 7,040 | 6,622 |
| | Pendimethalin | 1,039 | 788 | 788 |
| | Propaquizafop | 41 | 258 | 258 |
| | Prosulfocarb | 6,083 | 2,852 | 2,852 |
| | Rimsulfuron | 4 | 470 | 470 |
| | Insecticides | | | |
| | Dimethoate | 286 | 903 | 903 |
| | Lambda-cyhalothrin | 14 | 2,430 | 1,969 |
| | Molluscicides | | | |
| | Metaldehyde | 553 | 3,396 | 2,130 |
| | Seed Treatments | | | |
| | Flutolanil | 257 | 1,127 | 1,127 |
| | Imazalil | 56 | 2,114 | 2,114 |
| | Imazalil/pencycuron | 809 | 1,463 | 1,463 |

Acknowledgements

This report was compiled by James Quirke, with invaluable assistance from Michael Lavery, AFBI, Mr. David Williams, AFBI, Mr David Mathews AFBI, Joanne Kirby, AFBI and Amanda Patton, AFBI. Thanks are due to all the farmers who co-operated willingly with the survey and provided the detailed information needed to compile this report. Data collection for the survey was carried out by staff of the Agricultural Environment and Structures (AES) Division of the Department of Agriculture, Food and the Marine. Interviews with growers were conducted Pat Lordan, Jim Connolly, Gerry McPhilips, Hugh Gillespie, James Ryan, Richard McHugh, John Ryan, Pat Delaney, Tom Scanlon by and James Quirke.

References

- Pesticide Control Service, Department of Agriculture and Food Laboratories (2007).**
Pesticide Usage Survey, Report Number 2, Arable Crops, 2004.
- Pesticide Control Service, Department of Agriculture and Food Laboratories (2015).**
Pesticide Usage Survey, Report Number 3, Arable Crops, 2012

