

OPEN HERE

UFI: FGH0-936A-JN4Y-13WN
**RISK AND SAFETY
INFORMATION**

ALLY® MAX SX®
Herbicide

Contains tribenuron methyl. May produce an allergic reaction.

May cause damage to organs through prolonged or repeated exposure.

Very toxic to aquatic life with long lasting effects.

Do not breathe dust/ fume/ gas/ mist/ vapours/ spray.

Avoid release to the environment.

Get medical advice/ attention if you feel unwell.

Collect spillage.

Dispose of contents / container to a licensed hazardous-waste disposal contractor or collection site

except for triple rinsed empty clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions of use

PCS No. 03435



WARNING

FMC

ALLY® MAX SX®
Cereal herbicide

GROUP 2 HERBICIDE

PCS No: 03435

A water soluble granule formulation containing 143 g/kg metsulfuron-methyl and 143 g/kg tribenuron-methyl for weed control in wheat and barley, oats and triticale

FOR PROFESSIONAL USE ONLY

Registered trademark of FMC or an affiliate

Batch N°:

Contents: **84 g**

FMC Agro Limited, Rectors Lane, Pentre, Flintshire, CH5 2DH. Tel: 01244 537370. Email: fmc.agro.uk@fmc.com. Website: www.fmc-agro.co.uk. Technical Helpline: 01423 205011.

Emergency No.s - Medical Incidents Involving Crop Protection Products: England & Wales: 111, Scotland: 09454 24 24 24, Northern Ireland: Local GP or Pharmacist, Republic of Ireland: 01 837 9964.

Healthcare professionals seeking poisons information: www.toxbase.org.

Emergency No.s - Transportation, Warehousing & After Sale Incidents: CHEMTREC help centre Dialling from the UK and NI: 0870 820 0418, Dialling from Ireland: 01 901 4670



11005816_22072105_IRELAND

ALLY[®] MAX SX[®]

Cereal herbicide

GROUP 2 HERBICIDE

PCS No: 03435

A water soluble granule formulation containing 143 g/kg metsulfuron-methyl and 143 g/kg tribenuron-methyl for weed control in wheat and barley, oats and triticale.

FOR PROFESSIONAL USE ONLY

FMC Agro Limited, Rectors Lane, Pentre, Flintshire, CH5 2DH.
Tel: 01244 537370. Email: fmc.agro.uk@fmc.com. Website: www.fmc-agro.co.uk. Technical Helpline: 01423 205011.

Emergency No.s - Medical Incidents Involving Crop Protection Products: England & Wales: 111, Scotland: 08454 24 24 24, Northern Ireland: Local GP or Pharmacist, Republic of Ireland: 01 837 9964.

Healthcare professionals seeking poisons information: www.toxbase.org.
Emergency No.s - Transportation, Warehousing & After Sale Incidents: CHEMTRAC help centre Dialling from the UK and NI: 0870 820 0418, Dialling from Ireland: 01 901 4670

* Registered trademark of FMC or an affiliate

IMPORTANT INFORMATION FOR USE ONLY AS A PROFESSIONAL HERBICIDE				
Crops	Maximum individual and total dose (g product / ha)	Maximum number of treatments (per crop)	Latest time of application	Method of application
Wheat, barley, oats and triticale	42	1	Up to and including flag-leaf ligule / collar just visible (GS39)	Tractor mounted sprayer

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS

SAFETY PRECAUTIONS

ENVIRONMENTAL PROTECTION

Do not contaminate water with the product or its container (Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads).

Do not allow direct spray from ground crop sprayers to fall within 5m of the top of a bank of a static or flowing water body or within 1m of the top of a ditch that is dry at the time of application. Direct spray away from water.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

RESTRICTIONS

- ALLY® MAX SX® should not be applied to any crop suffering from stress as a result of drought, waterlogging, low

temperatures, pest or disease attack, nutrient or lime deficiency or other factors reducing crop growth.

- Do not use ALLY® MAX SX® on cereal crops undersown with grasses, clover or other legumes or any other broad-leaved crop.
- Due to the high level of activity of the herbicide, special care must be taken to avoid damage by drift onto broad-leaved plants outside the target area, or onto ponds, waterways or ditches. Thorough cleansing of equipment is also very important - see below.
- Contract agents should be consulted before using on crops grown for seed.
- ALLY® MAX SX® should not be applied within 7 days of rolling the crop.
- Do not apply ALLY® MAX SX® more than once to any cereal crop.
- All requirements or restrictions on other product labels must be adhered to when applied to the same crop as ALLY® MAX SX®. Contact your FMC distributor for further information.
- Do not tank-mix ALLY® MAX SX® with chlorpyrifos. Allow at least 14 days between application of ALLY® MAX SX® and chlorpyrifos treatments.

WEED CONTROL

ALLY® MAX SX® contains metsulfuron methyl and tribenuron methyl, sulfonyleurea (ALS inhibitor) herbicides which work mainly by contact action and is most effective if applied when the weeds are small and actively growing. Good spray cover of the weeds must be obtained for best results. Susceptible plants cease growth almost immediately after application and symptoms can be seen about two weeks after application. Weed control may be reduced when soil conditions are very dry. The susceptibility rating of weeds in the following table refer to good spray cover and good growing conditions.

Weed Species	Plants up to 2 expanded true leaves	Plants up to 6 expanded true leaves	Plants up to 15 cm across/high
Black-bindweed	MS	MS	
Charlock	S	S	S
Chickweed, Common	S	S	S
Cranes-Bill, Dove's Foot	S	S	-
Dead-nettle, Red	S	S	MS

Weed Species	Plants up to 2 expanded true leaves	Plants up to 6 expanded true leaves	Plants up to 15 cm across/high
Docks	S	S	S
Fat-hen	S	R	R
Field-speedwell, Common	S	S	MS
Forget-me-not, Field	S	MS	MS
Fool's Parsley	S	S	MS
Hemp-nettle, Common	S	S	S
Knotgrass	S	MS	MS
Mayweeds	S	S	S
Nettle, Small	S	S	-
Pale Persicaria	S	S	S
Pansy, Field	S	MS	MS
Parsley piert	S	S	S
Poppy, Common	S	S	MS

Weed Species	Plants up to 2 expanded true leaves	Plants up to 6 expanded true leaves	Plants up to 15 cm across/high
Redshank	S	S	S
Shepherd's-purse	S	S	S
Sowthistle, Smooth	S	–	–
Thistle, Creeping*	S	S	S
Volunteer Rape	S	S	–
Volunteer Sugar Beet	S	S	S

S = Susceptible, MS = Moderately susceptible,
MR = Moderately resistant, R = Resistant

* Best control of Creeping thistle is achieved from later application when thistles have emerged.

RESISTANCE

This product contains tribenuron-methyl and metsulfuron-methyl, which are ALS inhibitors, also classified by the Herbicide Resistance Action Committee as Group 2.

When herbicides with the same mode of action are used repeatedly over several years in the same field, selection of resistant biotypes can take place. These can propagate and may become dominating. A weed species is considered resistant to a herbicide if it survives a correctly applied treatment at the recommended dose. Development of resistance with a weed species can be avoided or delayed by using cultural methods of control; using herbicides with non-ALS modes of action within the cropping season and/or throughout the cropping rotation and sequencing or tank-mixing with suitable products having a different mode of action where two or more components are active against the target weeds. For the control of Chickweed, this product must always be applied in a mixture with an alternate mode of action herbicide at recommended rates.

A strategy for preventing and managing resistance should be adopted. Further details and advice on how to implement such a strategy may be obtained from your crop adviser or supplier. The Herbicide Resistance Action Committee (HRAC) also produces guidelines that may be consulted for additional information.

CROPS

ALLY® MAX SX® can be used on all varieties of wheat, barley, oats and triticale between the growth stages given below.

DOSE

Apply ALLY® MAX SX® at 42 g/ha.

TIMING

ALLY® MAX SX® should be applied in the spring after 1st February from the three-leaf stage (GS 13) up to and including the flag leaf ligule/collar just visible (GS 39).

FOLLOWING CROPS

Only cereals, oilseed rape, field beans or grass may be sown in the same calendar year as harvest of a cereal crop treated with ALLY® MAX SX®.

VOLUME AND APPLICATION

BEFORE USING ALLY® MAX SX®, SPRAYING EQUIPMENT MUST BE CLEAN AND FREE FROM CONTAMINATION WITH OTHER PESTICIDES.

Application should be made in 100-200 litres of water per hectare, using suitable ground equipment to give good spray cover of the weeds. When crops are thick or weed growth is dense, use 400 litres of water per hectare. Care should be taken not to overlap spray swaths.

MIXING

Quarter fill the spray tank with clean water, start the agitation and add the required quantity of ALLY® MAX SX® directly to the tank without prior creaming. Continue agitation while topping up the tank and while spraying.

COMPATIBILITY

Always add ALLY® MAX SX® to the spray tank first when using in tank-mix and ensure it is fully dispersed before adding the partner product. Do not allow ALLY® MAX SX® to come into contact with undiluted pesticide concentrate. Products should only be tank-mixed if each product can be applied within the label recommendations for its use. For further information contact your FMC distributor.

SPRAY TANK CLEAN-OUT

POOR CLEANOUT PRACTICES AND INSUFFICIENT WATER VOLUMES USED FOR THE RINSE PROCEDURE MAY RESULT IN INADEQUATE REMOVAL OF PRODUCT DEPOSITS. SUBSEQUENT USE OF APPLICATION EQUIPMENT IN THESE CIRCUMSTANCES MAY RESULT IN DAMAGE TO NON-CEREAL CROPS. TO AVOID SUBSEQUENT DAMAGE TO CROPS OTHER THAN CEREALS, IMMEDIATELY AFTER SPRAYING ALLY® MAX SX® THOROUGHLY CLEAN ALL SPRAY EQUIPMENT, INCLUDING INSIDE AND OUTSIDE OF LID, USING THE FOLLOWING PROCEDURE:

Always start with a clean tank and spray system. Clean spray equipment thoroughly immediately after use.

1. Thoroughly and completely rinse all interior tank surfaces (including lid) with water (use at least 10% of the tank capacity), taking care to remove any visible deposits. Flush pump, filters and boom after removing in-line strainers, nozzle tips and screens (clean these parts separately). Drain the remainder of the rinsate from the tank.
2. Repeat the rinse, flush and drain.

3. Dispose of washings safely. Do not spray onto sensitive crop or land intended for cropping with sensitive crop.

Note:

Consult label tank cleanup procedures for all tank mix partners and be sure to use the most rigorous procedure recommended.

WARNING

EXTREME CARE SHOULD BE TAKEN TO AVOID DAMAGE BY DRIFT OF SPRAY ONTO BROAD-LEAVED PLANTS OUTSIDE THE TARGET AREA OR ONTO PONDS, WATERWAYS OR DITCHES. SPRAYING EQUIPMENT SHOULD NOT BE DRAINED OR FLUSHED ONTO LAND PLANTED WITH OR INTENDED FOR PLANTING WITH TREES OR CROPS OTHER THAN CEREALS.

NOTICE TO BUYER

FMC as the manufacturer warrants that the composition of the product in sealed original FMC packaging complies with the information given on this label. FMC is not liable for any direct or indirect consequences resulting from improper storage, handling or use of the product contrary to or otherwise not in compliance with this label, law and regulations. Various factors, especially

local conditions, such as soil condition, plant varieties and weather conditions, may either cause the product not to have the full desired effect or result in damage to the crops treated. FMC does not assume any liability for such results.

FMC, the FMC logo, ALLY® MAX SX® are trademarks of FMC Corporation and/or an affiliate. ©2016-2026 FMC Corporation. All rights reserved.

Specimen-
2026 to date